



Bourne Worster, S., Stross, C., Vaughan, F. M. W. C., Linden, N., & Manby, F. R. (2019). Structure and Efficiency in Bacterial Photosynthetic Light Harvesting. *Journal of Physical Chemistry Letters*, 10, 7383-7390. [23].  
<https://doi.org/10.1021/acs.jpcllett.9b02625>

Peer reviewed version

Link to published version (if available):  
[10.1021/acs.jpcllett.9b02625](https://doi.org/10.1021/acs.jpcllett.9b02625)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the final published version of the article (version of record). It first appeared online via American Chemical Society at <https://pubs.acs.org/doi/10.1021/acs.jpcllett.9b02625> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

# Structure and Efficiency in Bacterial Photosynthetic Light-Harvesting

Susannah Bourne Worster<sup>a,†</sup> Clement Stross<sup>a,b,†</sup> Felix M. W. C. Vaughan<sup>a,b,c,†</sup>

Noah Linden<sup>b,†</sup> and Frederick R. Manby<sup>a,1\*,†</sup>

<sup>†a</sup>: *Centre for Computational Chemistry, School of Chemistry, University of Bristol,  
Bristol BS8 1TS, UK*

<sup>‡b</sup>: *School of Mathematics, University of Bristol, Bristol BS8 1TW, UK*


<sup>¶c</sup>: *Bristol Centre for Complexity Sciences, University of Bristol, Bristol, BS2 8BB, UK*

E-mail: [fred.manby@bristol.ac.uk](mailto:fred.manby@bristol.ac.uk)

## Abstract

Photosynthetic organisms use networks of chromophores to absorb and deliver solar energy to reaction centres. We present a detailed model of the light-harvesting complexes in purple bacteria, including explicit interaction with sunlight; radiative and non-radiative energy loss; and dephasing and thermalizing effects of coupling to a vibrational bath. We capture the effect of slow vibrations by introducing time-dependent disorder. Our model describes the experimentally observed high efficiency of light harvesting, despite the absence of long-range quantum coherence. The one-exciton part of the quantum state fluctuates continuously, but remains highly mixed at all times. These results suggest a relatively minor role for structure in determining efficiency. We build hypothetical models with randomly arranged chromophores, but still observe high efficiency when nearest-neighbour distances are comparable to those in nature. This helps explain the high transport efficiency in organisms with widely differing antenna structures, and suggests new design criteria for artificial light-harvesting devices.

## Graphical TOC Entry



Photosynthesis is the process by which organisms absorb energy in the form of sunlight and convert it into chemical energy, and it is the engine that produces almost all global biomass. Since the molecular architecture needed to carry out photosynthesis is costly to produce and maintain, sunlight is initially captured by an extensive antenna network of chlorophyll molecules (chromophores), through which absorbed energy must then be transported to specialised reaction centres to begin the photosynthetic reaction cycle. Experiments show that transport through the antenna can be highly efficient: absorption of a sufficiently energetic photon almost always leads to the charge-separation process that is a prerequisite for chemical storage of the energy.<sup>1-3</sup>

Two-dimensional spectroscopic investigations present the possibility of a substantial role for quantum coherence in achieving high efficiency,<sup>4,5</sup> an idea that was initially supported by a variety of theoretical and computational studies.<sup>6-11</sup> More recently this has been contested on the grounds that the oscillatory signature observed in the spectroscopy measurements might have been incorrectly interpreted,<sup>3,12-14</sup> or might be a result of the (unnatural) laser pulses used to excite the sample.<sup>15,16</sup> Furthermore, even if the oscillatory pattern truly revealed quantum coherence, it might not necessarily have an important functional role.<sup>17,18</sup> The research nevertheless sparked considerable interest in characterising the quantum state of the antenna over time and identifying criteria for efficiency. This has proved a challenging undertaking due to the complexity of the system and the number of factors influencing its behaviour. In particular, it has been shown that the nature of the light source,<sup>15,16</sup> the method used to model the vibrational environment,<sup>19</sup> the inclusion (or not) of energetic and spatial disorder,<sup>12,20</sup> and the consideration of short-time versus steady-state dynamics<sup>16,17,21-23</sup> all influence the way the system evolves. We aim to bring all these ideas together in a comprehensive and realistic model that includes detailed descriptions of each step of the process; from the continuous absorption of weak, incoherent sunlight to the final consumption of excitons at a reaction centre. We apply this model to a larger system than has yet been considered, including both of the major purple-bacteria light harvesting systems (LHI and

LHII) and the branched chromophore chains of the reaction centre (RC).

In addition, the suggestion that coherence may play a role in exciton transport inspired several attempts to design artificial systems that exploited the same principles. These systems, both hypothetical and real, require exquisite control over the placement and orientation of pigment chromophores in order to create and preserve the necessary coherence.<sup>20,24–26</sup> This makes them difficult to realise and those that have been created are small and delicately constructed. In the second part of this paper we consider an artificial antenna system of much simpler construction: a box of chromophores placed randomly (as if in solution) around a central reaction centre. Remarkably, the high efficiencies we observe demonstrate that structure (in the sense of a detailed arrangement of chromophores) does not play a significant role in determining the efficiency of an antenna. This could offer a pathway to designing cheaper, more scalable synthetic light harvesting devices.

First, we consider that the flow of energy through a photosynthetic antenna system is governed by a complex interplay of several physical processes: coherent evolution of non-stationary states; relaxation due to coupling with vibrational modes in the environment; absorption and emission of light; non-radiative decay, which results in exciton loss; and, finally, consumption of excitons at the reaction centre, triggering the biochemical cascade that ultimately leads to production of adenosine triphosphate. For reasons justified below, we model the system using a Lindblad master equation with a time-dependent Hamiltonian. Thus the evolution of the system density matrix is described by the Liouville-von Neumann equation

$$\frac{d\rho(t)}{dt} = -\frac{i}{\hbar} \left[ \hat{H}(t), \rho(t) \right] + (\mathcal{L}_{\text{rad}} + \mathcal{L}_{\text{bath}} + \mathcal{L}_{\text{nr}} + \mathcal{L}_{\text{sink}}) \rho(t). \quad (1)$$

Although each chromophore is a large, complex molecule, its participation in exciton transport is dominated by its electronic ground state  $|\psi_0\rangle$  and first excited state  $|\psi_1\rangle$  under biologically relevant conditions. Consequently, we model each chromophore site as a two-level system, with the states of a network of chromophores modelled as Hartree products of the

single-site states. We consider only the  $n + 1$  states with either zero ( $|0\rangle = |0_1, 0_2, \dots, 0_n\rangle$ ) or one ( $|i\rangle = |0_1 \dots 1_i \dots 0_n\rangle$ ) exciton.<sup>18</sup> Rather than focusing on a single component, as has been done previously, we consider transport across the whole antenna using a model system consisting of one LHII and one LHI/RC complex (*SI Appendix, Fig. S1*). The system contains a total of  $n = 65$  chromophores.

The unitary dynamics of the exciton are governed by a Hamiltonian of the form

$$\hat{H} = \sum_i E_i |i\rangle\langle i| + \sum_{i \neq j} V_{ij} |j\rangle\langle i| \quad (2)$$

where  $E_i$  are the excitation energies at each site and  $V_{ij}$  is the strength of the interaction between the transition dipole moments  $\boldsymbol{\mu}_i$  on sites  $i$  and  $j$ , calculated using the point dipole approximation. The point dipole approximation overestimates the nearest neighbour coupling in the LHI and RC complexes so these values were corrected to match the results of more detailed calculations (*SI Appendix*). Realistic disorder is introduced into the Hamiltonian by randomly drawing both  $E_i$  and  $|\boldsymbol{\mu}_i|$  from distributions computed using time-dependent density functional theory (TDDFT) on snapshots of a molecular dynamics (MD) simulation. The unit vectors  $\hat{\boldsymbol{\mu}}_i$  are taken separately from randomly selected MD snapshots. The data used to generate each Hamiltonian snapshot, including the molecular dynamics simulations and the distribution of  $E_i$  and  $|\boldsymbol{\mu}_i|$  were taken from calculations performed by some of us in ref. 12, where they were verified by reproducing the absorption spectrum of LHII.

The passage of excitations through the antenna is not determined solely by the antenna itself and it has been shown that different light sources induce very different dynamics.<sup>15,16,22</sup> Coherent pulses, localised in time and space, excite coherent superpositions of energy eigenstates, which then evolve coherently. Continuous incoherent illumination excites stationary eigenstates with well-defined energies. To understand the dynamics of photosynthesis it is therefore crucial to use a realistic model of the interaction between the system and incident radiation.

We capture this interaction using a Lindblad operator ( $\mathcal{L}_{\text{rad}}$ ) derived from the Born-Markov quantum optical master equation, including terms for both absorption ( $\mathcal{L}_{\text{rad}}^{\text{abs}}$ ) and spontaneous and stimulated emission ( $\mathcal{L}_{\text{rad}}^{\text{emit}}$ ). The rates of each process are parameterised to correctly account for the initial blackbody temperature of the sun, the attenuation of the radiation as it passes through the atmosphere, and the anisotropy of absorption into different eigenstates (*SI Appendix*).

At  $10^{-4}$  ps $^{-1}$  the rate of absorption is orders of magnitude slower than any of the other processes governing the dynamics of the antenna. As a result, the system remains almost entirely in the zero-exciton ground state for all time. We find that the amount of excited state present at any given time is on the order  $p = 10^{-9}$ .<sup>1</sup> To accelerate equilibration to the correct excited-state population, simulations are started from an initial state  $\rho_{\text{init}} = (1-p)|0\rangle\langle 0| + p|\psi\rangle\langle\psi|$  with the excited state manifold in a completely delocalised pure state  $|\psi\rangle = 1/\sqrt{n} \sum_j |j\rangle$ . As will be seen, the dynamics soon washes out any specific signatures of this initial state.

In addition to emission processes, exciton density is lost at the reaction centre, where the energy is used to trigger a series of electron transfers. As this process is effectively irreversible,<sup>27</sup> we treat the reaction centre as a sink through which excitons are continuously lost. It is described by the Lindblad operator

$$\mathcal{L}_{\text{sink}}\rho = \Gamma_{\text{sink}} \sum_{j \in \text{SP}} \left( |0\rangle\langle j|\rho|j\rangle\langle 0| - \frac{1}{2}\{ |j\rangle\langle j|, \rho \} \right), \quad (3)$$

where the sum runs over the chromophores in the ‘special pair’, the rate of exciton loss  $\Gamma_{\text{sink}}$  is taken as the rate of the first step in the electron transfer chain,<sup>28,29</sup> and  $\{ \cdot, \cdot \}$  denotes anticommutator.

Finally, the vibrational modes of the light-harvesting complexes and the surrounding

---

<sup>1</sup>A simple kinetics calculation gives the steady-state probability of being in the excited state as  $p = k_{\text{abs}}/(\Gamma_{\text{sink}} + \Gamma_{\text{nr}} + k_{\text{spont}} + k_{\text{stim}} + k_{\text{abs}})$ , where  $k_{\text{spont}}$ ,  $k_{\text{stim}}$  and  $k_{\text{abs}}$  are, respectively, the rates of spontaneous emission, stimulated emission and absorption. An order of magnitude estimate gives  $p = 10^{-9}$ .

medium have a huge impact on how excitons are transported through the antennae. They can act both to reduce the amount of excited state in the system by offering a pathway for non-radiative decay back to the ground state, and to redistribute remaining excitations by acting as a driving force for relaxation within the excited state manifold. Vibrations that are slow on the timescale of dynamics relevant for photosynthetic energy transport disrupt the symmetrical ring structure of the individual light-harvesting complexes, introducing variations in the excitation energies of individual chromophores, and in the interactions between them. Comparisons between exciton transfer rates calculated with multi-chromophoric Förster theory (MCFT) and basic Förster theory highlight the importance of treating the effects of both static and dynamic disorder.<sup>36,37</sup> We include each of these effects separately in our model.

The symmetry-breaking effects of slow vibrations are captured by introducing static disorder into the Hamiltonian, as described above. The exact form of the disorder will vary slowly with the motion of the vibration over the lifetime of an exciton. We approximate this time-dependence by generating a new, random Hamiltonian at regular intervals of 0.1 ps, corresponding approximately to a vibrational frequency below which higher harmonic levels become significantly occupied at 300 K. By updating the entire Hamiltonian periodically, we extend the example of previous works that use atomistic approaches to include a time-dependent description of disorder in the site excitation energies.<sup>38</sup> A similar technique has also been used to successfully describe the photochemistry of small organic molecules.<sup>39</sup>

Non-radiative loss of excitons competes with their biochemical utilisation at the reaction centre. We model it using a Lindblad operator  $\mathcal{L}_{\text{nr}}$  that has an identical form to  $\mathcal{L}_{\text{sink}}$  (equation 3), with  $j$  summed over every site in the system and a non-radiative decay constant

**Table 1: Rate constants and timescales for Lindblad terms**

	$\Gamma_{\text{sink}}$	$\Gamma_{\text{nr}}$	$\Gamma_{\text{deph}}$
Value / ps <sup>-1</sup>	0.125	0.001	11
References	28,29	30–34	7,35
Timescale	8 ps	1 ns	90 fs



$\Gamma_{\text{nr}}$ .

We consider two different models for the Lindblad term  $\mathcal{L}_{\text{bath}}$  describing the relaxation processes. The first is a local dephasing model

$$\mathcal{L}_{\text{deph}}\rho = \Gamma_{\text{deph}} \sum_j \left( |j\rangle\langle j| \rho |j\rangle\langle j| - \frac{1}{2} \{ |j\rangle\langle j|, \rho \} \right), \quad (4)$$

in which the coherences between the states  $|j\rangle$  localised on each site decay exponentially with a decay constant  $\Gamma_{\text{deph}}$ . Our second model is a global thermalizing model<sup>40</sup>

$$\mathcal{L}_{\text{therm}}\rho = \sum_{a \neq b} \frac{k_{b \rightarrow a}}{2\pi} \left( |a\rangle\langle b| \rho |b\rangle\langle a| - \frac{1}{2} \{ |b\rangle\langle b|, \rho \} \right), \quad (5)$$

in which exciton population is transferred between eigenstates  $|b\rangle$  and  $|a\rangle$  at a rate  $k_{b \rightarrow a}$ . The local model provides an intuitive picture of the effect of random environmental fluctuations but forces the system to relax to the maximally mixed state  $\rho_{\text{mm}} = \mathbb{I}/n$  (*SI Appendix*). By contrast, the global approach relaxes the system to the correct instantaneous thermal equilibrium state but does not explicitly account for the proximity and strength of coupling between chromophores. In a regime where inter-chromophore couplings are small on the scale of the system-bath coupling, this can result in an unphysical flow of energy between spatially well-separated and uncoupled chromophores.<sup>41</sup> A more in-depth analysis of the performance of the two models can be found in the *SI Appendix* and in ref. 42.

A further concern is the possible role of non-markovianity in the system-bath coupling.<sup>43,44</sup> However we find that the global Lindblad approach gives good qualitative agreement with preliminary calculations on small model systems using the exact hierarchical-equations-of-motion (HEOM) approach.<sup>45</sup>

The rate constants in equation 5 are determined using the Redfield theory expression<sup>46</sup>

$$\begin{aligned} k_{a \rightarrow b} &= \exp\left(\frac{\omega_{ab}}{k_{\text{B}}T}\right) k_{b \rightarrow a} \\ &= ((1 + n(\omega_{ab}))J(\omega_{ab}) + n(\omega_{ba})J(\omega_{ba})), \end{aligned} \quad (6)$$

where the relationship between  $k_{a \rightarrow b}$  and  $k_{b \rightarrow a}$  ensures that detailed balance is obeyed. The angular frequency  $\omega_{ab}$  is that of the energy gap between states  $|a\rangle$  and  $|b\rangle$  and  $n(\omega) = 1/(\exp(\omega/k_{\text{B}}T) - 1)$  is the Bose-Einstein distribution.

The spectral density  $\omega^2 J(\omega)$  describes the coupling of the excitonic system to the vibrational modes of the environment, under the assumption that all modes can be modelled as harmonic oscillators. This assumption breaks down for low-frequency modes, whose effect, however, we capture separately by periodically changing the Hamiltonian, as described above. We use a Debye spectral density, multiplied by a scaling factor of  $f = 11.87 \text{ rad ps}^{-1}$ , such that the thermalization timescale of the global Lindblad approximately matches that of the local dephasing model and the experimentally measured thermalization timescale of LHII<sup>47–49</sup> (see *SI Appendix*, Fig. S2):

$$\omega^2 J(\omega) = f \frac{2\omega_c \omega}{\omega^2 + \omega_c^2}. \quad (7)$$

The cut-off frequency  $\omega_c$  is taken as  $1/0.166 \text{ rad ps}^{-1}$ .<sup>50,51</sup>

Under the influence of  $\mathcal{L}_{\text{bath}}$ , the excited-state manifold relaxes rapidly towards a steady-state. When  $\mathcal{L}_{\text{bath}} = \mathcal{L}_{\text{deph}}$ , the dynamics for every realization of  $\hat{H}$  equilibrates to  $\rho_{\text{mm}}$ .

When  $\mathcal{L}_{\text{bath}} = \mathcal{L}_{\text{therm}}$  the steady-state is the thermal state of the current Hamiltonian, in which eigenstates are populated according to the Boltzmann distribution. The thermal state changes when the Hamiltonian changes, with an average trace distance of 0.4 between instantaneous thermal states. Following a change in the Hamiltonian, the exciton manifold reequilibrates to the new thermal state within a few hundreds of femtoseconds (see figure




Figure 1: Trace distance between the time-evolving state and the thermal state  $\rho_{\text{therm}} = \sum_{\alpha} \exp(E_{\alpha}/k_{\text{B}}T)/Z$  of the current Hamiltonian as a function of time for the LHI-LHII model system. The Hamiltonian changes at 1 ps, as does the corresponding  $\rho_{\text{therm}}$ . It takes several hundred femtoseconds for the system to requilibrate, under the influence of the Hamiltonian and the thermalizing Lindblad operator  $\mathcal{L}_{\text{therm}}$ .

1), in keeping with experimentally measured equilibration times in individual LHCs.<sup>47</sup> The Hamiltonian changes on a similar timescale (every 100 fs), such that the quantum state of the antenna is constantly evolving (see figure 2a). However, since individual, instantaneous thermal states are not too dissimilar, the system spends much of its time in states that are close to thermal and the time-averaged state  $\rho_{\text{av}}$ , shown in figure 2b, is close (trace distance  $\approx 0.18$ ) to the average of the thermal states for all realisations of the Hamiltonian  $\langle \rho_{\text{therm}} \rangle_{\text{av}}$ . The longer the period between changes of the Hamiltonian, the more closely  $\rho_{\text{av}}$  resembles  $\langle \rho_{\text{therm}} \rangle_{\text{av}}$ .

Thermal states are, by nature, mixed states. The extent to which the state of a system is mixed is quantified by  $\text{tr } \rho^2$ , which is 1 for a pure state and  $1/N$  for the maximally mixed state. It may be shown analytically (*SI Appendix*) that the full average over the thermal states of all possible Hamiltonians is exactly the maximally mixed state. Correspondingly, the values of  $\text{tr } \rho^2$ , given in table 2, for the average states of both the full system and each component subsystem are small and close to their values for the maximally mixed state. On average then, the system is not only in a highly mixed state but in one that equally favours equivalent sites within a ring.

**Table 2: Properties of subsystem states calculated for the renormalized substates of the average state**

	LHI-LHII	B875	B850	B800
$\text{tr } \rho^2$	0.073	0.087	0.135	0.114
Min. value of $\text{tr } \rho^2$	0.015	0.03	0.06	0.11
IPR	3.78	8.41	6.41	1.57
Max. value of IPR	65	32	18	9

While  $\text{tr } \rho^2$  quantifies the character (pure/mixed) of the state and sets an upper bound on the amount of coherence that it may contain (*SI Appendix*), it does not directly measure the extent of coherent delocalisation of the state. This is done using the inverse participation ratio (IPR):<sup>52</sup>

$$\text{IPR} = \frac{\left[ \sum_{jk} |\rho_{jk}| \right]^2}{n_{\text{site}} \sum_{jk} |\rho_{jk}|^2}, \quad (8)$$

which is bounded between 1 (incoherent and fully localised) and  $n_{\text{site}}$  (fully delocalised coherent pure state). The average state contains very little coherence, as evidenced by the small values of the IPR in table 2. In B800 particularly, there is essentially no coherence (IPR = 1.57), reflecting the fact that the chromophores are well-spaced and weakly coupled. Stronger coupling in the B850 and B875 rings introduces coherences between neighbouring sites, raising the value of the IPR. These short-range coherences correspond to exciton states delocalized over pairs of chromophores. Studies using MCFT have shown that these pairwise coherent states play an important role in determining the B800-B850 transfer rate, since the corresponding splitting of the energy levels improves spectral overlap with the B800 states.<sup>36,53</sup> However, when  $\rho_{\text{B850}}$  and  $\rho_{\text{B875}}$  are expressed in the reduced basis of the BChl dimer subsystem, to more accurately reflect the structure of these rings, they are also found to be essentially maximally mixed (*SI Appendix, Fig. S3*).

Overall, the excitonic subsystem is, on average, in a highly mixed, incoherent state that nevertheless places population equally on all sites. Instantaneously, the system is never in its average state but rather fluctuates continuously between instantaneous thermal states. However, since the thermal states all have similar forms (though favouring different chromophore sites), the properties of the system do not fluctuate a great deal. Plots of  $\text{tr } \rho(t)^2$  and IPR in figure 2 (panels c and d) indicate that, at all times, the state of the system remains highly mixed with no long-range coherence.

The efficiency of a photosynthetic apparatus can be defined in a number of different ways, not all of which are equivalent.<sup>21</sup> In this work, we will define the efficiency as

$$\mathcal{E}[\rho] = \frac{\langle 0 | \mathcal{L}_{\text{sink}} \rho | 0 \rangle}{\langle 0 | (\mathcal{L}_{\text{sink}} + \mathcal{L}_{\text{nr}} + \mathcal{L}_{\text{rad}}^{\text{emit}}) \rho | 0 \rangle}. \quad (9)$$

For each unit of energy that is absorbed by the antenna, the efficiency tells us what fraction is used productively by the sink. This measure reflects how well the system is adapted to transport and use the absorbed energy. Note that it does not tell us what fraction of

incident light reaches the reaction centre as this would require additional consideration of how efficiently the system absorbs light. While this is also an important question, we will not consider it here.

The efficiency of the antenna, plotted in figure 3, remains fairly constant over time, reflecting the consistent form of the quantum state throughout the evolution. The efficiencies obtained with the two variants of  $\mathcal{L}_{\text{bath}}$  span the experimentally measured efficiencies for photosystem II in higher plants (84%–91%).<sup>1,3</sup>

Using  $\mathcal{L}_{\text{bath}} = \mathcal{L}_{\text{therm}}$ , the time-averaged efficiency  $\langle \mathcal{E}[\rho(t)] \rangle_{\text{av}}$  is 96%. This is likely an overestimate, since the global model does not account for the weaker coupling between B875 and the RC, which significantly retards the rate of exciton transfer between them.<sup>54</sup> Hence, this model tends to overestimate the population of the RC (*SI Appendix, Fig. S4*), on which  $\mathcal{E}$  is heavily dependent (*SI Appendix*).

By contrast, using  $\mathcal{L}_{\text{bath}} = \mathcal{L}_{\text{deph}}$  underestimates the efficiency by failing to capture the energy gradient between LHII and LHI and consequently underestimating the population of the B875 ring and the rate of exciton transfer onto the RC. Using the local dephasing model  $\langle \mathcal{E}[\rho(t)] \rangle_{\text{av}}$  is 62%. This value is highly sensitive to the number of antenna chromophores, since exciton density is distributed equally between all chromophores under the action of  $\mathcal{L}_{\text{deph}}$ . Therefore, increasing the number of chromophores reduces the population on any single site. For example, if only LHI is modelled,  $\langle \mathcal{E}[\rho(t)] \rangle_{\text{av}} = 75\%$ .

In addition to observing minimal fluctuations in the efficiency, we find that  $\langle \mathcal{E}[\rho(t)] \rangle_{\text{av}}$  is closely reproduced (also 96% for the global model) by  $\mathcal{E}[\rho_{\text{av}}]$ , the efficiency of the time-averaged state. Together, these results indicate that, despite short-time fluctuations in the state, the system evolves continuously through states that are similar in character both to each other and to their time average.

The highly symmetric ring structures of natural photosynthetic antenna systems are often thought to imply a special physical significance for the particular spatial arrangement of chromophores (such as maintaining long-lived coherence). We investigated this hypothesis

by calculating the efficiency of a box of 32 chromophores (for comparison with LHI) placed and oriented randomly around a central reaction centre. The ‘antenna’ chromophores were not allowed within 3 Å of each other or within 45 Å of the middle of the reaction centre (*SI Appendix*). The dimensions of the box were adjusted to alter the concentration of antenna chromophores without changing the ratio of chromophores to reaction centre. Relaxation was implemented using the local dephasing model ( $\mathcal{L}_{\text{deph}}$ ), which is sensitive to chromophore spacing and particularly appropriate to the lower concentrations investigated. Each concentration was sampled 32 times with different random configurations of the chromophores. The results are shown in figure 4.

Two clear trends are visible. Firstly, as the concentration of chromophores is reduced, the mean efficiency decreases. More weakly coupled chromophores do not transfer the exciton as quickly to the reaction centre (if at all) so more excited state is lost through non-radiative decay and emission processes. Secondly, the standard deviation in the efficiency increases with decreased concentration. This can be understood by considering that in a small box, at high concentration, the chromophores will always be close to each other and to the reaction centre. At lower concentrations, the random placement of chromophores will sometimes result in clusters of closely-spaced, strongly coupled chromophores and other times in all the chromophores being far apart and only weakly coupled. Crucially, when the chromophores were closely spaced, there was very little variation in the efficiency over the 32 sampled configurations, indicating that the orientation of the chromophores was not significant.

The same behaviour was observed whether the chromophores were placed freely in 3D (figure 4a), as they might be in a liquid phase, or confined to a pseudo-2D slab (figure 4b), to mimic the layered structure of natural antenna networks.

Whilst chromophore spacing is an important factor for determining efficiency, it is equally important to balance the rates of competing processes in the system. Figure 4c illustrates the impact of varying the dephasing rate. There is a sharp drop in efficiency when the dephasing rate exceeds  $100 \text{ ps}^{-1}$ , where continual measurement by the environment prevents the system

evolving. Typically, simulations have shown that efficiency drops off at low dephasing rates too, as the exciton is trapped by Anderson localisation. However, in our calculations this effect is largely mitigated by the coupling to slow vibrations.

In conclusion, we have examined the nature and evolution of the quantum state of a photosynthetic antenna system under constant illumination by a natural, incoherent light source. When talking about the state of light harvesting systems it is common to refer only to the part of the state that contains exactly one exciton.

However, it is worth noting that the overall quantum state is, to a very good approximation, the zero-exciton ground state. This contrasts with the often-employed picture of photon absorption triggering a large change in the excitonic state of a single chromophore. While this picture is relevant for understanding ultrafast spectroscopic experiments, it has little bearing on the phenomenon of photosynthesis in sunlight.

Within the detailed model we have constructed of the many processes governing the state and dynamics of the excitonic subsystem, accurately capturing the effect of interacting with a vibrational environment is particularly challenging, since the similar strengths of inter-system and system-bath coupling places the problem uncomfortably between the ranges of validity of commonly used approximations.<sup>3,55</sup> The two approaches we have employed embody the two opposing regimes of strongly coupled chromophores interacting collectively with the bath ( $\mathcal{L}_{\text{therm}}$ ) and weakly coupled chromophores interacting independently with a local bath ( $\mathcal{L}_{\text{deph}}$ ). Both methods make the markovian approximation so will not capture the small non-markovian effects that may arise from strongly coupled discrete vibrational modes. Although neither approach can, on its own, reliably describe the effect of the vibrational environment, we obtain valuable insights by observing features that arise from both descriptions.

Firstly, we conclude that the state of the excitonic subsystem is at all times highly mixed with little or no coherence. Indeed, both descriptions indicate that, on the timescale of the sink (around 10 ps), the average state is close to the maximally mixed state within each ring subsystem. In other words, exciton density is distributed equally between equivalent sites in



an incoherent manner. Over the whole antenna, the distribution of excited state population is skewed towards LHI because of the energy gradient between successive ring components and the large number of chromophores.<sup>25,54</sup> The importance of this effect is exemplified by the low efficiency obtained using the local dephasing model, which spreads exciton density uniformly over the whole antenna system.

The coupling to a vibrational environment, and the consequent relaxation, is an important mechanism for transporting excitations through the antenna and overcoming disorder-driven localization. The role of dephasing in exciton transport has been noted before and is sometimes referred to as environment-assisted transport.<sup>21,56,57</sup> What we additionally see here is that slow vibrational modes keep the exciton density moving continuously around the antenna. This is in good agreement with atomistic simulations of LHII, which also show excitons moving perpetually around the system, with a bias towards lower energy ring components.<sup>58</sup> In the absence of the time-dependent disorder induced by slow vibrational modes, the efficiency of the antenna, under the global Lindblad model, is simply the efficiency of the thermal state. This is invariably a high value, although, depending on the particular static geometry being investigated, the efficiency in the absence of slow modes may be slightly higher or slightly lower than the value obtained in the presence of slow vibrations. However, the limitations of the global model, discussed above, mean we cannot conclusively rule out the possibility that slow vibrations play a role in preventing exciton density becoming trapped in areas of the antenna far from the reaction centre. More detailed investigations would be needed to determine whether slow vibrations play an important role in exciton transport or whether they are simply a coincidental feature of natural protein-based antennae.

Without the requirement for coherence, the rigid structural constraints that have been put forward to maintain coherence can be relaxed.<sup>24-26,59</sup> Indeed, the results of figure 4 demonstrate that the orientation of chromophores has little bearing on the antenna efficiency. Instead, the typical nearest neighbour distance appears to be the key determinant of efficiency. This is consistent with previous work by Mohseni *et al.*, showing that the rate of

exciton transfer across a randomly arranged collection of chromophores becomes less sensitive to chromophore orientation at higher chromophore concentrations.<sup>60</sup> It is also supported by the observations of Baghbanzadeh *et al.* that adjusting the spacing of chromophores in natural LHI and LHII can compensate for any loss in efficiency that might arise from randomising the chromophore orientations.<sup>61</sup> We note, however, that our more comprehensive treatment of processes affecting the dynamics, particularly the inclusion of dephasing processes, means that we observe a loss of sensitivity to chromophore orientation at higher chromophore concentrations than used in ref. 61.

So why are highly organised arrangements of chromophores found in nature? Effects such as concentration quenching and the energetic cost of synthesising chromophores (which we did not include in our simulation) would make high chromophore concentrations undesirable. We suggest that perhaps the organisation of chromophores into circular structures may simply be a means of avoiding the extremely high pigment concentration that would ensure the right separation in a completely disordered solution. It is possible that the same end could be achieved in an artificial system without the need for complex protein architecture, for example, by anchoring chromophores to a bead. However, to design an appropriate artificial matrix to support the chromophores, further investigation would be needed into the role of the environment in determining the rates of dephasing and non-radiative decay, which need to be carefully balanced to maintain efficiency.

## Supporting Information Available

Details on constructing Hamiltonians; light-matter Lindblad; benchmarking of bath models. Notes on interpreting  $\text{tr}(\rho^2)$ , and the relationship between efficiency and the population of sink sites. Mathematical proofs that local dephasing equilibrates to the maximally mixed state, and that the average of the thermal states is the maximally mixed state. Additional figures: Structure of bacterial antenna complexes; relaxation timescale of LHII under global

Lindblad; reduced density matrix of B850 in dimer basis; sample chromophore populations; properties of evolving state under local dephasing model. Tables of Hamiltonian parameters.

## Acknowledgement

We gratefully acknowledge the funding agencies that supported this work: CS was supported through the doctoral training grant of the Engineering and Physical Sciences Research Council (EPSRC); FMWCV was supported through the EPSRC Bristol Centre for Complexity Sciences; SBW is supported by a research fellowship from the Royal Commission for the Exhibition of 1851.

## References

- (1) Wientjes, E.; Van Amerongen, H.; Croce, R. Quantum yield of charge separation in photosystem II: Functional effect of changes in the antenna size upon light acclimation. *Journal of Physical Chemistry B* **2013**, *117*, 11200–11208.
- (2) Rijgersberg, C. P.; Van Grondelle, R.; Amesz, J. Energy transfer and bacteriochlorophyll fluorescence in purple bacteria at low temperature. *Biochimica et Biophysica Acta - Bioenergetics* **1980**, *592*, 53–64.
- (3) Fassioli, F.; Dinshaw, R.; Arpin, P. C.; Scholes, G. D. Photosynthetic light harvesting: excitons and coherence. *Journal of The Royal Society Interface* **2013**, *11*, 20130901–20130901.
- (4) Engel, G. S.; Calhoun, T. R.; Read, E. L.; Ahn, T. K.; Mančal, T.; Cheng, Y. C.; Blankenship, R. E.; Fleming, G. R. Evidence for wavelike energy transfer through quantum coherence in photosynthetic systems. *Nature* **2007**, *446*, 782–786.

- (5) Lee, H.; Cheng, Y.-C.; Fleming, G. R. Coherence Dynamics in Photosynthesis: Protein Protection of Excitonic Coherence. *Science* **2007**, *316*, 1462–1465.
- (6) Calhoun, T. R.; Fleming, G. R. Quantum coherence in photosynthetic complexes. *Physica Status Solidi (B) Basic Research* **2011**, *248*, 833–838.
- (7) Harel, E.; Engel, G. S. Quantum coherence spectroscopy reveals complex dynamics in bacterial light-harvesting complex 2 (LH2). *Proceedings of the National Academy of Sciences* **2012**, *109*, 706–711.
- (8) Panitchayangkoon, G.; Hayes, D.; Fransted, K. A.; Caram, J. R.; Harel, E.; Wen, J.; Blankenship, R. E.; Engel, G. S. Long-lived quantum coherence in photosynthetic complexes at physiological temperature. *Proceedings of the National Academy of Sciences* **2010**, *107*, 12766–12770.
- (9) Strümpfer, J.; Sener, M.; Schulten, K. How quantum coherence assists photosynthetic light-harvesting. *Journal of Physical Chemistry Letters* **2012**, *3*, 536–542.
- (10) Collini, E.; Wong, C. Y.; Wilk, K. E.; Curmi, P. M.; Brumer, P.; Scholes, G. D. Coherently wired light-harvesting in photosynthetic marine algae at ambient temperature. *Nature* **2010**, *463*, 644–647.
- (11) van Hulst, N. F.; Cogdell, R. J.; Brinks, D.; Hildner, R.; Nieder, J. B. Quantum Coherent Energy Transfer over Varying Pathways in Single Light-Harvesting Complexes. *Science* **2013**, *340*, 1448–1451.
- (12) Stross, C.; Van der Kamp, M. W.; Oliver, T. A. A.; Harvey, J. N.; Linden, N.; Manby, F. R. How Static Disorder Mimics Decoherence in Anisotropy Pump-Probe Experiments on Purple-Bacteria Light Harvesting Complexes. *The Journal of Physical Chemistry B* **2016**, *120*, 11449–11463.

- (13) Christensson, N.; Kauffmann, H. F.; Pullerits, T.; Mančal, T. Origin of long-lived coherences in light-harvesting complexes. *Journal of Physical Chemistry B* **2012**, *116*, 7449–7454.
- (14) Duan, H.-G.; Prokhorenko, V. I.; Cogdell, R. J.; Ashraf, K.; Stevens, A. L.; Thorwart, M.; Miller, R. J. D. Nature does not rely on long-lived electronic quantum coherence for photosynthetic energy transfer. *Proceedings of the National Academy of Sciences* **2017**, *114*, 8493–8498.
- (15) Mančal, T.; Valkunas, L. Exciton dynamics in photosynthetic complexes: Excitation by coherent and incoherent light. *New Journal of Physics* **2010**, *12*.
- (16) Brumer, P.; Shapiro, M. Molecular response in one-photon absorption via natural thermal light vs. pulsed laser excitation. *Proceedings of the National Academy of Sciences* **2012**, *109*, 19575–19578.
- (17) Kassal, I.; Yuen-Zhou, J.; Rahimi-Keshari, S. Does coherence enhance transport in photosynthesis? *Journal of Physical Chemistry Letters* **2013**, *4*, 362–367.
- (18) Tiersch, M.; Popescu, S.; Briegel, H. J. A critical view on transport and entanglement in models of photosynthesis. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **2012**, *370*, 3771–3786.
- (19) Fassioli, F.; Olaya-Castro, A.; Scheuring, S.; Sturgis, J. N.; Johnson, N. F. Energy transfer in light-adapted photosynthetic membranes: From active to saturated photosynthesis. *Biophysical Journal* **2009**, *97*, 2464–2473.
- (20) Sarovar, M.; Birgitta Whaley, K. Design principles and fundamental trade-offs in biomimetic light harvesting. *New Journal of Physics* **2013**, *15*, 013030.
- (21) Manzano, D. Quantum Transport in Networks and Photosynthetic Complexes at the Steady State. *PLoS ONE* **2013**, *8*, 1–8.

- (22) Brumer, P. Shedding (Incoherent) Light on Quantum Effects in Light-Induced Biological Processes. *The Journal of Physical Chemistry Letters* **2018**, *9*, 2946–2955.
- (23) Mančal, T.; Valkunas, L. Exciton dynamics in photosynthetic complexes: Excitation by coherent and incoherent light. *New Journal of Physics* **2010**, *12*.
- (24) Buckhout-White, S.; Spillmann, C. M.; Algar, W. R.; Khachatryan, A.; Melinger, J. S.; Goldman, E. R.; Ancona, M. G.; Medintz, I. L. Assembling programmable FRET-based photonic networks using designer DNA scaffolds. *Nature Communications* **2014**, *5*, 5615.
- (25) Hemmig, E. A.; Creatore, C.; Wunsch, B.; Hecker, L.; Mair, P.; Parker, M. A.; Emmott, S.; Tinnefeld, P.; Keyser, U. F.; Chin, A. W. Programming Light-Harvesting Efficiency Using DNA Origami. *Nano Letters* **2016**, *16*, 2369–2374.
- (26) Olejko, L.; Bald, I. FRET efficiency and antenna effect in multi-color DNA origami-based light harvesting systems. *RSC Adv.* **2017**, *7*, 23924–23934.
- (27) Blankenship, R. E.; Tiede, D. M.; Barber, J.; Brudvig, G. W.; Fleming, G.; Ghirardi, M.; Gunner, M. R.; Junge, W.; Kramer, D. M.; Melis, A. et al. Comparing Photosynthetic and Photovoltaic Efficiencies and Recognizing the Potential for Improvement. *Science* **2011**, *332*, 805–809.
- (28) Greenfield, S. R.; Seibert, M.; Govindjee,; Wasielewski, M. R. Direct Measurement of the Effective Rate Constant for Primary Charge Separation in Isolated Photosystem II Reaction Centers. *The Journal of Physical Chemistry B* **2002**, *101*, 2251–2255.
- (29) Demmig-Adams, B.; Garab, G.; Adams Iii, W. *Advances in Photosynthesis and Respiration Including Bioenergy and Related Processes Non-Photochemical Quenching and Energy Dissipation in Plants, Algae and Cyanobacteria Non-Photochemical Quenching and Energy Dissipation in Plants, Algae and Cyanobacteri*; 2014.

- (30) Palacios, M. A.; De Weerd, F. L.; Ihalainen, J. A.; Van Grondelle, R.; Van Amerongen, H. Superradiance and exciton (de)localization in light-harvesting complex II from green plants? *Journal of Physical Chemistry B* **2002**, *106*, 5782–5787.
- (31) Zankel, K. L.; Reed, D. W.; Clayton, R. K. Fluorescence and photochemical quenching in photosynthetic reaction centers. *Proceedings of the National Academy of Sciences* **2006**, *61*, 1243–1249.
- (32) Pandit, A.; Shirzad-Wasei, N.; Wlodarczyk, L. M.; Van Roon, H.; Boekema, E. J.; Dekker, J. P.; De Grip, W. J. Assembly of the major light-harvesting complex II in lipid nanodiscs. *Biophysical Journal* **2011**, *101*, 2507–2515.
- (33) Monshouwer, R.; Abrahamsson, M.; van Mourik, F.; van Grondelle, R. Superradiance and Exciton Delocalization in Bacterial Photosynthetic Light-Harvesting Systems. *The Journal of Physical Chemistry B* **2002**, *101*, 7241–7248.
- (34) Bopp, M. a.; Jia, Y.; Li, L.; Cogdell, R. J.; Hochstrasser, R. M. Fluorescence and photobleaching dynamics of single light-harvesting complexes. *Proceedings of the National Academy of Sciences of the United States of America* **1997**, *94*, 10630–5.
- (35) Fidler, A. F.; Singh, V. P.; Long, P. D.; Dahlberg, P. D.; Engel, G. S. Time scales of coherent dynamics in the light-harvesting complex 2 (LH2) of rhodobacter sphaeroides. *Journal of Physical Chemistry Letters* **2013**, *4*, 1404–1409.
- (36) Scholes, G. D.; Fleming, G. R. On the Mechanism of Light Harvesting in Photosynthetic Purple Bacteria: B800 to B850 Energy Transfer. *The Journal of Physical Chemistry B* **2000**, *104*, 1854–1868.
- (37) Beljonne, D.; Curutchet, C.; Scholes, G. D.; Silbey, R. J. Beyond förster resonance energy transfer in biological and nanoscale systems. *Journal of Physical Chemistry B* **2009**, *113*, 6583–6599.

- (38) Shim, S.; Rebentrost, P.; Valleau, S.; Aspuru-Guzik, A. Atomistic study of the long-lived quantum coherences in the Fenna-Matthews-Olson complex. *Biophysical Journal* **2012**, *102*, 649–660.
- (39) Schile, A. J.; Limmer, D. T. Simulating conical intersection dynamics in the condensed phase with hybrid quantum master equations. *arXiv:1905.0029v [cond-mat.stat-mech]* **30 Apr 2019**,
- (40) Ostilli, M.; Presilla, C. Thermalization of noninteracting quantum systems coupled to blackbody radiation: A Lindblad-based analysis. *Physical Review A* **2017**, *95*, 062112.
- (41) Hofer, P. P.; Perarnau-Llobet, M.; Miranda, L. D. M.; Haack, G.; Silva, R.; Brask, J. B.; Brunner, N. Markovian master equations for quantum thermal machines: Local versus global approach. *New Journal of Physics* **2017**, *19*.
- (42) Wu, J.; Liu, F.; Shen, Y.; Cao, J.; Silbey, R. J. Efficient energy transfer in light-harvesting systems, I: Optimal temperature, reorganization energy and spatial-temporal correlations. *New Journal of Physics* **2010**, *12*.
- (43) Caruso, F.; Chin, A. W.; Datta, A.; Huelga, S. F.; Plenio, M. B. Entanglement and entangling power of the dynamics in light-harvesting complexes. *Physical Review A - Atomic, Molecular, and Optical Physics* **2010**, *81*, 1–8.
- (44) Vaughan, F. M. W. C.; Linden, N.; Manby, F. R. How Markovian is exciton dynamics in purple bacteria? *Journal of Chemical Physics* **2017**, *146*.
- (45) Vaughan, F. M. W. C. Exciton Energy Transfer in Photosynthetic Light Harvesting Complexes. Ph.D. thesis, University of Bristol, 2017.
- (46) May, V.; Kühn, O. *Charge and Energy Transfer Dynamics in Molecular Systems*; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2011.



- (47) Stuart, T. A.; Vengris, M.; Novoderezhkin, V. I.; Cogdell, R. J.; Neil Hunter, C.; Van Grondelle, R. Direct visualization of exciton reequilibration in the LH1 and LH2 complexes of *Rhodobacter sphaeroides* by multipulse spectroscopy. *Biophysical Journal* **2011**, *100*, 2226–2233.
- (48) Pullerits, T.; Chachisvilis, M.; Jones, M. R.; Hunter, C. N.; Sundström, V. Exciton dynamics in the light-harvesting complexes of *Rhodobacter sphaeroides*. *Chemical Physics Letters* **1994**, *224*, 355–365.
- (49) Freiberg, A.; Timpmann, K.; Lin, S.; Woodbury, N. W. Exciton Relaxation and Transfer in the Lh2 Antenna Network of Photosynthetic Bacteria. *Journal of Physical Chemistry B* **1998**, *102*, 10974–10982.
- (50) Ishizaki, A.; Fleming, G. R. Theoretical examination of quantum coherence in a photosynthetic system at physiological temperature. *Proceedings of the National Academy of Sciences* **2009**, *106*, 17255–17260.
- (51) Zhu, J.; Kais, S.; Rebentrost, P.; Aspuru-Guzik, A. Modified scaled hierarchical equation of motion approach for the study of quantum coherence in photosynthetic complexes. *Journal of Physical Chemistry B* **2011**, *115*, 1531–1537.
- (52) Meier, T.; Chernyak, V.; Mukamel, S. Multiple Exciton Coherence Sizes in Photosynthetic Antenna Complexes viewed by Pump-Probe Spectroscopy. *The Journal of Physical Chemistry B* **1997**, *101*, 7332–7342.
- (53) Jang, S.; Newton, M. D.; Silbey, R. J. Multichromophoric Förster resonance energy transfer from B800 to B850 in the Light harvesting complex 2: Evidence for subtle energetic optimization by purple bacteria. *Journal of Physical Chemistry B* **2007**, *111*, 6807–6814.
- (54) Blankenship, R. E. *Molecular Mechanisms of Photosynthesis*, 1st ed.; Blackwell Science Ltd: Oxford, UK, 2002.

- (55) Olaya-Castro, A.; Fassioli, F. Characterizing quantum-sharing of electronic excitation in molecular aggregates. *Procedia Chemistry* **2011**, *3*, 176–184.
- (56) Pelzer, K. M.; Can, T.; Gray, S. K.; Morr, D. K.; Engel, G. S. Coherent transport and energy flow patterns in photosynthesis under incoherent excitation. *Journal of Physical Chemistry B* **2014**, *118*, 2693–2702.
- (57) Caruso, F.; Chin, A. W.; Datta, A.; Huelga, S. F.; Plenio, M. B. Highly efficient energy excitation transfer in light-harvesting complexes: The fundamental role of noise-assisted transport. *The Journal of Chemical Physics* **2009**, *131*, 105106.
- (58) Sisto, A.; Stross, C.; van der Kamp, M.; O’Connor, M.; Johnson, G. T.; Hohenstein, E. G.; Manby, F. R.; Glowacki, D. R.; Martinez, T. J. Atomistic absorption spectra and non-adiabatic dynamics of the LH2 complex with a GPU-accelerated ab initio exciton model. *Phys. Chem. Chem. Phys.* **2017**, *19*, 5–12.
- (59) Knee, G. C.; Rowe, P.; Smith, L. D.; Troisi, A.; Datta, A. Structure-Dynamics Relation in Physically-Plausible Multi-Chromophore Systems. *Journal of Physical Chemistry Letters* **2017**, *8*, 2328–2333.
- (60) Mohseni, M.; Shabani, A.; Lloyd, S.; Omar, Y.; Rabitz, H. Geometrical effects on energy transfer in disordered open quantum systems. *Journal of Chemical Physics* **2013**, *138*.
- (61) Baghbanzadeh, S.; Kassal, I. Geometry, Supertransfer, and Optimality in the Light Harvesting of Purple Bacteria. *Journal of Physical Chemistry Letters* **2016**, *7*, 3804–3811.




Figure 2: Evolution of the state of the exciton manifold in the full LHI-LHII model system. The Hamiltonian changes every 0.1 ps. Dephasing is implemented using the global model  $\mathcal{L}_{\text{therm}}$  (an equivalent figure for the local model can be found in *SI Appendix, Fig. S5*). Rates of other processes are given in table 1. (a) Trace distance of the instantaneous evolving state  $\rho(t)$  from the current thermal state  $\rho_{\text{therm}}$ , which changes with the Hamiltonian (only a short time sample is shown so that the details of the evolution are more clearly visible). The coloured bar along the top illustrates the amount of time spent close to a thermal state, with dark blue marking stretches of time where the trace distance is  $\leq 0.2$ , and light blue where the trace distance is  $> 0.2$ . (b) Plot of  $\log_{10} |\text{Re}(\rho_{\text{av}})|$ . The renormalized average density matrix  $\rho_{\text{av}}$  is calculated from 5 ps onward, to avoid any influence from the specific choice of initial state. (c) Degree of purity quantified by  $\text{tr} \rho(t)^2$ . (d) The inverse participation ratio of  $\rho(t)$ . For (c) – (d), the mean value of each quantity starting from 5 ps is displayed on the graph.




Figure 3: Efficiency of the LHI-LHII antenna over the same period of evolution plotted in figure 2. From top to bottom, the traces correspond to efficiency of the full antenna (one LHI and one LHII) with  $\mathcal{L}_{\text{bath}} = \mathcal{L}_{\text{therm}}$ ; efficiency of the LHI/RC complex with  $\mathcal{L}_{\text{bath}} = \mathcal{L}_{\text{deph}}$ , and efficiency of the full antenna with  $\mathcal{L}_{\text{bath}} = \mathcal{L}_{\text{deph}}$ . The mean efficiency, displayed to the right hand side of each trace, is calculated from 5 ps.




Figure 4: Efficiency as a function of chromophore concentration in a (a) cubic ( $d \times d \times d$ ) box or (b) pseudo-2D ( $d \times d \times 3\text{\AA}$ ) slab of 32 chromophores arranged randomly around a centrally placed reaction centre. Each concentration was sampled 32 times. Range bars indicate one standard deviation above and below the mean efficiency. Horizontal dotted lines indicate, for comparison, the efficiency of the LHI-RC complex in purple bacteria, using the local dephasing model. In each sample, the efficiency was averaged over the evolution from 10 ps to 100 ps. (c) Efficiency as a function of dephasing rate for one arrangement of chromophores in a cubic box with  $d = 125\text{\AA}$ . (d). Example structure of a chromophore box with  $d = 95\text{\AA}$  (corresponding to  $10^6/(d/\text{\AA})^3 = 1.17$ ).