

Fungal Genetics Reports

Volume 43

Article 9

PCR from fungal spores after microwave treatment

Adlane Ferreira

University of British Columbia

V. B. Glass

University of British Columbia

N. Louise

University of British Columbia

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Ferreira, A., V.B. Glass, and N. Louise (1996) "PCR from fungal spores after microwave treatment," *Fungal Genetics Reports*: Vol. 43, Article 9. <https://doi.org/10.4148/1941-4765.1305>

This Regular Paper is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

PCR from fungal spores after microwave treatment

Abstract

We describe a fast template preparation method for PCR amplification from fungal spores using microwave irradiation. The method is useful when the availability of fungal material is restricted or the number of processed samples is large.

PCR from fungal spores after microwave treatment

Ferreira, Adlane V.B. and Glass, N. Louise- Biotechnology Laboratory and Dept. Botany, University of British Columbia, Vancouver, Canada, V6T 1Z4.

We describe a fast template preparation method for PCR amplification from fungal spores using microwave irradiation. The method is useful when the availability of fungal material is restricted or the number of processed samples is large.

The polymerase chain reaction (PCR) has been widely used for diverse genetic analyses in fungi, including population genetics and phylogenetic studies. For some of these analyses a large number of DNA samples are needed. Although several simplifications of DNA extraction protocols have been reported, most of them involve the mechanical disruption of mycelial tissue, use of organic solvents and various centrifugation steps. Direct PCR from fungal spores is not readily suitable to all fungi mainly because of difficulties in rupturing the cell wall. Although spores from some fungi can be directly amplified after a prolonged initial step at 94°C in the PCR (Aufauvre-Brown *et al.* 1993, *Curr. Genet.* **24**:177-178), spores from different fungi can be less susceptible to heat denaturation in aqueous solution (Saupe S. and Ferreira A.V.B., unpublished results). Direct spore PCR was performed after freezing the samples at very low temperatures (Jin-Rong and Hamer, 1995 *Fungal Genet. Newsl.*, **42**:80) but it was shown that an excessive number of spores can inhibit the PCR.

Microwave irradiation has proven to be useful in DNA extraction protocols from different eukaryotes (Goodwin and Lee 1993, *Biotechniques* **15**:438-444). It is presumed that microwave irradiation acts by exposing the DNA normally protected by the cell structures. Here we describe a simple 10-minute template preparation method for PCR amplification from minute quantities of fungal spores using microwave irradiation.

Vegetative spores from five filamentous fungal species were taken from an agar slant or plate with a loop and transferred to an empty 500 µl Eppendorf tube. *Neurospora crassa* sexual spores were also tested. Each tube was closed and the dry spores were irradiated in full power (700W) for 5 min in a domestic microwave oven (Danby D811). A beaker containing water was present during the irradiation to avoid damage to the oven. Thirty microliters of TE (10 mM Tris-HCl pH 7.5, 1 mM EDTA) was added to the tube, the tube was vortexed and then spun in a microfuge for 1-5 min at 14,000 RPM. The supernatant liquid was collected and transferred to a clean Eppendorf tube. Five microliters of the supernatant liquid was used in a 50 µl PCR. An example of the results of a PCR using the internal transcribed spacer (ITS) ribosomal region primers (White *et al.* 1990, *PCR Protocols*, Academic Press.) is shown in Figure 1. DNA prepared from five different fungal species using the microwave method was amplified under the following conditions: 10 mM Tris-HCl pH 9.0, 50 mM KCl, 1.5 mM MgCl₂, 0.1% Triton X-100, 5% glycerol, 0.2 mM of dNTPs, 5 units *Taq* DNA polymerase and 0.02 mM of each of the ITS1 and ITS4 primers. After 40 cycles of 95 C, 55 C and 72 C for 1 min. each, half of the reaction mixture was subjected to electrophoresis in 1.5% agarose gels. A control PCR is shown (lane 8) performed with 100 ng of phenol:chloroform-purified *N. crassa* DNA.

We have used this method to screen *N. crassa* transformants for homologous recombination at specific sites using a variety of different primers. Amplification from samples prepared in this manner was efficient even for a single copy gene. At least 6 PCR reactions can be performed if starting with approximately 10⁷ spores (based on *N. crassa*). The time of microwave irradiation could be changed for different species as we have had satisfactory amplifications of *N. crassa* DNA after only one minute (but not 30 sec) of irradiation (not shown). This method drastically reduces manipulation and template preparation time and therefore minimizes the possibility of contamination of the samples with extraneous DNA.

Acknowledgments

We thank Dr. Sven Saupe for his useful comments. A.V.B. F. is supported by a fellowship from the Brazilian Ministry of Education.




Figure 1. PCR of ITS from filamentous fungi after microwave treatment. Lanes (1) *Fusarium oxysporum*; (2) *Penicillium expansum*; (3) *Alternaria alternata*; (4) *Cochliobolus bicolor*; (5) *N. crassa* vegetative and (6) sexual spores; (7) no DNA control and (8) *N. crassa* phenol:chloroform- extracted DNA. (M) marker, BRL 1 Kb ladder.

Last modified 7/25/96 KMC