

Fungal Genetics Reports

Volume 44

Article 6

Why are nit-2 transformants of *Aspergillus nidulans* partially derepressed?

M. A. Davis
University of Melbourne

M. J. Hynes
University of Melbourne

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Davis, M. A., and M.J. Hynes (1997) "Why are nit-2 transformants of *Aspergillus nidulans* partially derepressed?," *Fungal Genetics Reports*: Vol. 44, Article 6. <https://doi.org/10.4148/1941-4765.1274>

This Regular Paper is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Why are nit-2 transformants of *Aspergillus nidulans* partially derepressed?

Abstract

The major nitrogen metabolite repression control genes of *Aspergillus nidulans* and *Neurospora crassa* are the *areA* and *nit-2* genes respectively. Both genes encode transcriptional activators with highly conserved C2-X17-C2 zinc finger DNA binding domains which recognize similar WGATAR DNA sequences (Kudla *et al.*, 1990 EMBO J. 9: 1355-1364; Fu and Marzluf, 1990 Proc. Natl. Acad. Sci USA 87: 5331-5335). We have shown previously that the *N. crassa nit-2* gene can fully complement an *areA* loss-of-function mutation in *A. nidulans*, restoring growth on a wide variety of nitrogen sources. However, the nit-2 transformants are partially derepressed (Davis and Hynes, 1987 Proc. Natl. Acad. Sci. USA 84: 3753-3757).

Why are *nit-2* transformants of *Aspergillus nidulans* partially derepressed?

Meryl A. Davis and Michael J. Hynes- Department of Genetics, University of Melbourne, Parkville 3052, Australia

The major nitrogen metabolite repression control genes of *Aspergillus nidulans* and *Neurospora crassa* are the *areA* and *nit-2* genes respectively. Both genes encode transcriptional activators with highly conserved C2-X17-C2 zinc finger DNA binding domains which recognize similar WGATAR DNA sequences (Kudla *et al.*, 1990 EMBO J. **9**: 1355-1364; Fu and Marzluf, 1990 Proc. Natl. Acad. Sci USA **87**: 5331-5335). We have shown previously that the *N. crassa nit-2* gene can fully complement an *areA* loss-of-function mutation in *A. nidulans*, restoring growth on a wide variety of nitrogen sources. However, the *nit-2* transformants are partially derepressed (Davis and Hynes, 1987 Proc. Natl. Acad. Sci. USA **84**: 3753-3757).

We have considered a number of possible explanations for the apparent inability of the Nit-2 protein to respond fully to nitrogen control in *A. nidulans*.

1. One possibility is that the Nit-2 protein is overexpressed in *A. nidulans* and that elevated levels of Nit-2 lead to derepression. Expression of *nit-2* is from the *N. crassa* promoter and transformants were generated by random integration into the *A. nidulans* genome. However, no correlation was observed between plasmid copy number and the degree of derepression of individual transformants. In addition, we have generated random multicopy *areA* transformants and found that these retain nitrogen control. The fact that overexpression of AreA protein does not lead to equivalent derepression also argues against this possibility.
2. A second possibility is that the Nit-2 protein is not interacting efficiently with the product of the *A. nidulans* homolog of the *nmr-1* gene. In *N. crassa*, there is clear evidence that the Nmr-1 protein is required for repression. *nmr-1* mutants are derepressed and the Nit-2 and Nmr-1 proteins have been shown to interact physically via sequences in the DNA-binding region and the C-terminal regions of Nit-2 (Xiao *et al.*, 1995 Biochemistry **34**:8861-8868). The *nit-2* and *areA* genes show considerable divergence outside these regions and the overall conformation of the two proteins may differ such that the presumptive *A. nidulans* Nmr-1 equivalent protein may not interact effectively with Nit-2 to prevent its activity. We have found that this possibility is also not a sufficient explanation for the partial depression phenotype. We have co-transformed the *N. crassa nmr-1* gene, together with the *nit-2* gene, into *areA* mutants of *A. nidulans* and find no difference between transformants which contain *nmr-1* and those that do not contain *nmr-1*.
3. A third possibility is that the *nit-2* clone used in the initial experiments lacks key sequences in the 3' untranslated region (3'UTR) of the gene. Recent studies (Platt *et al.*, 1996 EMBO J. **14**:2791-2801) have shown that the *areA* gene encodes sequences present in the 3'UTR of the *areA* mRNA which have a role in determining rates of mRNA turnover. Deletion of these sequences results in a more stable mRNA and a partially derepressed phenotype. The original *nit-2* clone used was a 6kb *EcoRI* clone (pnit-2). Inspection of the *nit-2* sequence shows an *EcoRI* site only 170bp downstream from the translation stop codon and hence the 3' end of the *nit-2* gene may have been truncated with respect to these key sequences.

While neither Nit-2 overexpression nor Nit-2/Nmr-1 protein interactions appear to be sufficient explanation for the partial derepression of *nit-2* transformants, the possibility of differences in RNA turnover needs further investigation. The primary experiment would be to transform an *A. nidulans areA* loss of function mutant with a longer *nit-2* clone. If the results of this experiment support the prediction that such transformants would regain full nitrogen control, a number of important conclusions can be drawn. The results would indicate that *areA* and *nit-2* share this additional level of control. Furthermore, it would help to localise the key sequences as they must lie downstream of the 3' *EcoRI* site of the *nit-2* gene. Platt *et al.* (1996) identified a region of conserved sequence 3' to the *A. nidulans areA* gene and the *Penicillium chrysogenum nre* gene. Comparison of this sequence with the 3' untranslated sequences of *nit-2* suggests there may be a less well conserved sequence (Figure 1) downstream of the *EcoRI* site (nt 4389). Conserved sequences in the 3' UTR region of these genes may therefore indicate functionally important sites for nitrogen source-regulated turnover of both *areA* and *nit-2* mRNAs.

Acknowledgments: We acknowledge the support of the Australian Research Council and we thank Dr. G. Marzluf for providing the plasmid pnmr-1.

```

Nit2UTR      4454 TGCTTTTTACTCTGGCCTTTTTTAAACGGTACGGCTGTTGAAGCTGTCAAGTTTCGA
areAUTR      3035 TGTGCTTGACT-TGTGATTTTTGCTTGATAACAACCGC---GTCTGTCCGGTACCTG
          **  ** *** **  *****  * *** * *  ***** ** *

Nit2UTR      AGGACAGCTTTTCTAAACATTTCTCCCCGCACGTCTTATTTCTGGTACACAAAG
areAUTR      GCCTTCGTCTGTCCGG-TACCTGGCCTTCGTCTGCTTTTTTTGTTTGTGTTTCATG
          *  * **  *  * ** **  * ** ***  * **  * **

Nit2UTR      CGTTGATAGAAGAAGGAGCCGGGCTGATGAGGATGGGAAGAGGGGCCAGGAG
areAUTR      TGTT-ATGTTTGTGTTGCATTTCGAGGTCAGGCCGGGCAGAAGTGCGCATCTC
          *** **  *  * **  * * ***  *** *** * ** **

```

Figure 1. Alignment of *nit-2* (Nit2UTR) and *areA* (areAUTR) 3' sequences using Clustal W (Thompson *et al.* 1994 Nucl. Acid Res. **22**: 4673-4680) accessed via ANGIS (Australian National Genome Information Service).

[Return to the FGN 44 Table of Contents](#)

[Return to the FGSC home page](#)