

Fungal Genetics Reports

Volume 14

Article 21

Kynureninase determination

M. Vukosavovitch

J. R. Turner

W. K. Matchett

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Vukosavovitch, M., J.R. Turner, and W.K. Matchett (1969) "Kynureninase determination," *Fungal Genetics Reports*: Vol. 14, Article 21. <https://doi.org/10.4148/1941-4765.2052>

This Enzyme Methodology is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Kynureninase determination

Abstract

Kynureninase determination

Vukosavovich, M., J. R. Turner and W. K.

Matchett. A direct method of assaying
kynureninase.

A rapid method for assaying kynureninase has been developed. As in the procedure described by Jakoby and Bonner (1953 J. Biol. Chem. 205:699), the method involves detection of anthranilic acid produced by measuring the intensity of its fluorescence. Extracts prepared from tryptophan-induced wild type strain 74A were used as a source of kynureninase. The activity was purified approximately 4-fold by ammonium sulphate precipitation and Sephadex chromatography.

The reaction mixture contained 0.16 μ mole pyridoxal phosphate, 3 μ mole magnesium sulfate, 0.06 μ mole kynurenine, 50 μ mole Tris-(hydroxymethyl) aminomethan pH 8.0, and protein (50-500 μ g) in a final volume of 1.2 ml. The reaction was conducted at 30°C in the cuvette of an Aminco Bowman spectrofluorimeter equipped with thermal spacer. Anthranilic acid which appeared during the reaction was detected by measuring the intensity of its fluorescence at 409 nm. Activation was at 315 nm.

The amount of anthranilic acid produced is linear with time for about 5 minutes, after which the rate decreases. The velocity of the reaction is proportional to protein concentration from 0.045 to 0.54 mg/ml and extrapolates to zero at zero protein concentration. The velocity of the reaction approached a maximal value when the kynurenine concentration was 0.05 μ mole/ml. Kynur-

enine inhibits the reaction in concentrations greater than $0.16 \mu\text{mole/ml}$. The rate of conversion of kynurenine to anthranilic acid was measured at kynurenine concentrations from $5 \times 10^{-3} \mu\text{mole/ml}$ to $1 \times 10^{-1} \mu\text{mole/ml}$ using the direct method. A standard Lineweaver-Burke plot of the data describes a line with an intercept = 0.133, slope = 1.88×10^{-6} , correlation coefficient = 0.99. The K_m calculated from these data was 1.42×10^{-5} , which is essentially the same as that published by Jakoby and Bonner.

It has previously been reported that magnesium ions activate the enzyme at all stages of purification. The assays reported here have no magnesium requirement. Traces of magnesium present in the water could activate the reaction; however, EDTA had no effect on the rate. The reaction proceeds without pyridoxal phosphate; however, the amount of anthranilic acid produced is not linear with time. Successive additions of pyridoxal phosphate decrease the initial rate slightly, but the amount of anthranilic acid produced becomes a linear function of time. At $0.13 \mu\text{M/ml}$ the rate was constant for about 5 minutes. Increasing the concentration to $0.53 \mu\text{M/ml}$ decreased the rate to $1/3$ of the value observed at $0.13 \mu\text{M/ml}$.

The time required for the assay by this method is less than 10 minutes. - - - Biology Department, Pacific Northwest Laboratory, Battelle Memorial Institute, Richland, Washington 99352.