

Isolation of g-amylase

D. N. Fass

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Fass, D. N. (1969) "Isolation of g-amylase," *Fungal Genetics Reports*: Vol. 14, Article 24. <https://doi.org/10.4148/1941-4765.2055>

This Enzyme Methodology is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Isolation of g-amylase

Abstract

Isolation of g-amylase

Fass, D. N. Isolation of γ -amylase (gluc-
amylase) from the culture filtrate of N. crassa.

An amylase-super-producer strain, e.g., inos (89601) a, (FGSC#498) is grown 84-96 hours in 1% sucrose Vogel's medium. Good yields may be expected from 1 liter cultures in 2.8 l Fernbach flasks incubated at 25°C on a shaker.

Adequate aeration is essential for production of the enzyme. At the end of this time, the mycelia are removed by filtration and the medium is chilled to 4°C and cold ethanol is added to 40%. The solution is allowed to stand overnight at 4°C and the resulting precipitate is removed by centrifugation at 25000 X g for 10 min. To the alcoholic supernatant is added a water solution of 2% glycogen in the proportion of 25 ml/l of original medium. The white precipitate which forms is centrifuged immediately at 4000 X g for 10 min and redissolved in a small volume of Vogel's salts. This mixed enzyme solution is incubated for 1 hr at 37°C and then dialyzed twice at 4°C against 50 vols of citrate buffer 0.01 M in Na⁺, pH 5.0, for 4 hrs and 8 hr. The ~~enzyme~~ sample is applied to a 2 x 15 cm column of Amberlite CG-50 equilibrated with the same buffer. Elution is carried out at 4°C with a 500 ml linear gradient from 0.01 to 1.1 N Na⁺ at approximately 40-50 ml/hr. Citrate is the counter-ion.

The amylolytic activity recovered at about 0.4 N Na⁺ shows an E-fold increase in specific activity, no invertase or α -amylase activity, and a single band in acrylamide gel electrophoresis. This work was supported in part by the NSF and NIH Training Grant in Genetics (T01-GM01316) to Florida State University. ■ ■ -Genetics Laboratories, Department of Biological Science, Florida State University, Tallahassee, Florida 32306.