

## Determination of total phosphorus in extracts

S. C. Hedman

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

---

### Recommended Citation

Hedman, S. C. (1969) "Determination of total phosphorus in extracts," *Fungal Genetics Reports*: Vol. 14, Article 13. <https://doi.org/10.4148/1941-4765.2044>

This Technical Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact [cads@k-state.edu](mailto:cads@k-state.edu).

---

## Determination of total phosphorus in extracts

### Abstract

Determination of total phosphorus in extracts

Hedman, S. C. Determination of total phosphorus in *Neurospora* extracts.

Often one desires to ascertain the total phosphorus content of various *Neurospora* extracts. The following method has been found applicable for a wide variety of such extracts. This method incorporates various features of previously published procedures as well as some new modifications. There are two parts to the procedure: acid hydrolysis and phosphate determination.

**Acid hydrolysis:** 0.2 ml of extract (containing 1-10  $\mu\text{g}$  of phosphorus) is placed in an acid-cleaned 15 x 150 mm Kimax test tube. 0.3 ml of 5 N  $\text{H}_2\text{SO}_4$  and 0.9 ml of  $\text{H}_2\text{O}$  are added. The contents are slowly heated over a Bunsen burner until dense white fumes of  $\text{SO}_3$  are given off. At this point, the contents of the test tube may be dark-brown to black in color. After cooling the mixture, 0.1 ml of 2 N  $\text{HNO}_3$  is added and heat is applied until  $\text{SO}_3$  is again given off. This  $\text{HNO}_3$  treatment is repeated until the contents of the test tube are colorless. The volume is then brought to 1.5 ml by the addition of  $\text{H}_2\text{O}$  and the tube is heated in a 100°C water bath for 5 minutes to hydrolyze pyrophosphates. An acid-cleaned glass marble is placed over the top of the test tube to prevent excessive evaporation.

**Phosphate determination:** To 1.5 ml of hydrolyzed extract are added 1.2 ml of phosphate reagent. The phosphate reagent is made as follows: (a) Stock solution: 50 g of ammonium molybdate.  $4\text{H}_2\text{O}$  are dissolved in 400 ml of 10 N  $\text{H}_2\text{SO}_4$  with constant stirring. After all is in solution, the volume is brought to 500 ml with additional 10 N  $\text{H}_2\text{SO}_4$ . This stock solution can be stored for several months at room temperature. (b) Preparation of reagent: The phosphate reagent must be made up fresh for each series of assays. To make 20 ml of such reagent, 2.0 ml of stock solution are added to 14 ml of  $\text{H}_2\text{O}$  containing 1.0 mg of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ . After the ferrous sulfate is in solution, the volume is brought to 20 ml with distilled water.

After five to ten minutes, the absorbance of each tube is read at 710  $\text{m}\mu$  in 1 ml cuvettes of 1.0 cm path length. A reagent blank is used as a reference. Under these conditions, linearity is observed between absorbance and phosphorus content over the range of 1-10  $\mu\text{g}$  of phosphorus routinely gives an optical density of  $0.464 \pm 0.010$ .

The phosphate determination by itself can also be utilized to determine the total orthophosphate content as, for example, when assaying for phosphatase activity. The following compounds do not appear to interfere with this method: tris buffer (0.2 M), trichloroacetic acid (20% w/v), bovine serum albumin (400  $\mu\text{g}/1.5$  ml),  $\text{CHCl}_3$ ,  $\text{C}_2\text{H}_5\text{OH}$ , Cleland's reagent ( $10^{-4}$  M), or sucrose (0.3 M). ■ ■ ■ Department of Biology, University of Minnesota, Duluth, Duluth, Minnesota 55812.