

Fungal Genetics Reports

Volume 14

Article 9

Effects of "No-Pest" Strip Insecticide

S. M. Michalek

H. E. Brockman

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Michalek, S. M., and H.E. Brockman (1969) "Effects of "No-Pest" Strip Insecticide," *Fungal Genetics Reports*: Vol. 14, Article 9. <https://doi.org/10.4148/1941-4765.2039>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Effects of "No-Pest" Strip Insecticide

Abstract

Effects of "No-Pest" Strip Insecticide

A test for mutagenicity of Shell "No-Pert Strip Insecticide" in Neurospora crassa.

small flying insects for up to three months. Use one strip for each 1000 (10'x10'x10') cubic feet of enclosed area. Note: No-Pert strips are not designed to kill insects instantaneously. They work by continuous action affecting the insect over a short period of time. Caution: Keep out of reach of children. Do not get in mouth: harmful if swallowed. After prolonged storage, a small amount of liquid may form on the strips. Do not get liquid in eyes. Wash hands thoroughly with soap and water after handling strip. Do not contaminate feed, water, and food stuffs."

An interesting aspect of this product is that the occupants of a room containing No-Pert Strip are exposed to the insecticide continuously. We decided that the possible mutagenic effects of No-Pert Strip should be investigated.

Forward mutation at the adenine-3 (ad-3) region of a standard wild-type strain, 74-OR23-1A (Core, Brockman and de Serrer 1965 Neurospora Ned. 8: 25), and of a heterokaryon, 74-OR31-16A + 74-OR60-29A (de Serrer and Webber 1963 Neurospora Newsl. 3:3), of Neurospora crassa was studied. The heterokaryon was selected because multilocus deletions arising from chromosome breakage and intragenic mutations can be detected (de Serres 1968 Genetics 58:69). Slants of Fries' basal medium (Horowitz and Beadle 1943 J. Biol. Chem. 150:325) supplemented with 1% sucrose and 1.5% agar were inoculated and grown under three different conditions for 6 days. One set of slants (control) was placed in a 10-gallon aquarium that had been covered with plate glass and sealed with petroleum jelly 24 hr before addition of the slants. A second set of slants (experimental) was placed under identical conditions except a No-Pert Strip was placed in the aquarium 24 hr before addition of the slants. Both aquaria were re-sealed after addition of the slants and remained sealed for 6 days. A third set of slants (room control) was placed in the room (23°C) containing the aquaria. Vegetative growth and conidiation occurred at a lower rate in the experimental cultures than in the control cultures, and the amount of conidiation at 6 days was much less in the experimental cultures than in the control cultures. The viability of conidia from cultures grown under the three conditions was the same.

Conidia were harvested from the vegetative cultures after 6 days of growth and arrayed for ad-3 mutants by a direct method (de Serres and Kålmark 1958 Nature 182: 1249) or previously described (Brockmon and de Serres 1963 Genetics 48:597; Webber and de Serrer 1965 Proc. Natl. Acad. Sci. U.S.A. 53:430). The results are given in the Table. The spontaneous mutation frequency in a previous experiment was 0.4 ad-3 mutants per 1×10^6 colonies (Brockman and de Serrer 1963).

Results of forward-mutation experiment.

Strain	Condition	Colonies assayed	<u>ad-3</u> mutants
74-OR23-1A	Room control	1.1×10^6	0
	Control	0.9×10^6	0
	Experimental	1.7×10^6	0
74-OR31-16A	Room control	3.3×10^6	0
	Control	2.0×10^6	0
74-OR60-29A	Experimental	2.7×10^6	0

We conclude that Shell "No-Pert Strip Insecticide" does not induce forward mutation at the ad-3 region or lethality in N. crassa under the conditions of our experiments. It inhibits the rate of vegetative growth and conidiation and decreases the final amount of conidiation. (Research supported by the U.S. Atomic Energy Commission under Contract No. AT(11-1)-1314 with Illinois State University; Report number COO-1314-11). - - - Department of Biological Sciences, Illinois State University, Normal, Illinois 61761.