

Fungal Genetics Reports

Volume 20

Article 7

Revertibility of me-1 alleles

R. Aiuto

Albion College

L. MacPhail

Albion College

C. A. Hall

Albion College

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Aiuto, R., L. MacPhail, and C.A. Hall (1973) "Revertibility of me-1 alleles," *Fungal Genetics Reports*: Vol. 20, Article 7. <https://doi.org/10.4148/1941-4765.1815>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Revertibility of *me-1* alleles

Abstract

Revertibility of *me-1* alleles

Differential revertibility of me-1 alleles.

allelic although functionally related (Aiuto 1969 Genetics 61:s53), was also studied. The me-1 gene is relatively small and its alleles are largely non-complementing (Aiuto, unpublished data).

Table 1: Spontaneous and induced reversion rates of five methionine-requiring strains (prototrophs per 10^6 surviving conidia).*

Strain	Spont.	NA	MNNG	ICR-170	EMS	OMHA
<u>me</u> (35599)	0.00	0.00	0.00	0.00	0.00	0.00
<u>me-1</u> (2-3)	0.29	0.27	0.20	0.75	0.80	0.66
<u>me-1</u> (2-4)	0.00	0.13	0.21	0.00	0.00	0.00
<u>me-1</u> (2-65)	0.22	13.30	21.40	2.77	12.50	12.40
<u>me-1</u> (17-M154)	0.00	0.00	1.80	7.86	5.21	0.81

*complete data available from senior author.

This note reports chemical mutagenesis experiments similar to those of Malling, de Serres, and co-workers with ad-3B. The four alleles of me-1 studied (2-3, 2-4, 2-65, 17-M154) were induced by Barry with nitrous acid. A presumed me-1 allele, 35599, which has been shown to be non-

Methods used were those described by Malling and de Serres (1966 Mutat. Res. 3:470; 1968 Mutat. Res. 5:359; 1968 Mutat. Res. 6:181; and pers. commun.) The sample of ICR-170 was obtained from H. J. Creech, Institute for Cancer Research, Philadelphia, Pa. The results are presented in Table 1.

As in the ad-3B work (Kilbey, et al. 1971 Mutat. Res. 12:47), some indications of the original type of mutant lesion can be made: possible gene deletion (35599, UV-induced), non-revertible alleles (2-3 and 2-4), base substitution (2-65), and base addition or deletion (17-M154). Twenty additional me-1 alleles are being tested with the five mutagens.