

Fungal Genetics Reports

Volume 37

Article 9

Designation of newly identified mutagen-sensitive mutations in *Neurospora crassa* and their linkage data

H. INOUE
Saitama University

C. ISHII
Saitama University

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

INOUE, H., and C. ISHII (1990) "Designation of newly identified mutagen-sensitive mutations in *Neurospora crassa* and their linkage data," *Fungal Genetics Reports*: Vol. 37, Article 9. <https://doi.org/10.4148/1941-4765.1474>

This Regular Paper is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Designation of newly identified mutagen-sensitive mutations in *Neurospora crassa* and their linkage data

Abstract

Two years ago, Dr. Käfer and Dr. Schroeder independently summarized genetic information on mutagen-sensitive mutants in *Neurospora crassa* (Fungal Genetics Newsletter 35:11-13, 1988; DNA Repair 3:77-98, 1988). So far, 30 genetic loci related to mutagen-sensitivity have been identified. The mutants are more sensitive to UV, X-rays and/or chemical mutagens than the wild-type strain. We have studied DNA-repair mechanisms of *Neurospora* for several years. During the study, several mutagen-sensitive mutants were isolated and the mutations were mapped. To confirm that they are mutations in new genes, we tested allelism between these mutations and previously reported mutagen-sensitive mutations. Here, new mutations are listed and new map numbers are given to them as follows.

Designation of newly identified mutagen-sensitive mutations in *Neurospora crassa* and their linkage data

Inoue, H. and C. Ishii Laboratory of Genetics, Faculty of Science, Saitama University, Urawa 338, Japan.

Two years ago, Dr. Käfer and Dr. Schroeder independently summarized genetic information on mutagen-sensitive mutants in *Neurospora crassa* (Fungal Genetics Newsletter 35:11-13, 1988; DNA Repair 3:77-98, 1988). So far, 30 genetic loci related to mutagen-sensitivity have been indentified. The mutants are more sensitive to UV, X-rays and/or chemical mutagens than the wild-type strain. We have studied DNA-repair mechanisms of *Neurospora* for several years. During the study, several mutagen-sensitive mutants were isolated and the mutations were mapped. To confirm that they are mutations in new genes, we tested allelism between these mutations and previously reported mutagen-sensitive mutations. Here, new mutations are listed and new mus numbers are given to them as follows.

allele (isolation #)	gene#	linkage group (near marker)	sensitivity	
			UV	MMS
SA11	mus-31	IR (ad-3B)	0	S
SA32	mus-32	IR (ad-3B)	S	S
SA33	mus-33	VII (met-7)	S	S
SA18	mus-34	VR (am)	S	S
SA50	mus-35	VII (nic-3)	0	S
SA51	mus-36	IV (pan-1)	0	S
SA52	mus-37	V (cyh-2)	0	S
SA56	mus-38	IL (leu-3)	S	0

0 - similar to wild type

S - more sensitive than wild type

In Fungal Genetics Newsletter 35:11-13, 1988, *mus(FK131)*, *mus(FK132)* and *mus(FK133)* were not yet mapped. We have mapped the mutations and tested allelism with other mutagen-sensitive mutations. The results showed that both *mus(FK131)* and *mus(FK132)* are allelic to *mus-21*. The *mus(FK133)* mutant has a mutation in a new gene, located in linkage group VI (8% to *ylo-1*). Thus *mus(FK133)* has been designated *mus-39*. These new mutants will be characterized for mutagen-sensitivity, mutagenesis, meiosis and fertility.