

Fungal Genetics Reports

Volume 27

Article 10

Cellobiose-induced B-galactosidase and B-glucosidase activities of *Neurospora crassa*.

P. J. Russell

C. B. Perry

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Russell, P. J., and C.B. Perry (1980) "Cellobiose-induced B-galactosidase and B-glucosidase activities of *Neurospora crassa*," *Fungal Genetics Reports*: Vol. 27, Article 10. <https://doi.org/10.4148/1941-4765.1673>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Cellobiose-induced B-galactosidase and B-glucosidase activities of *Neurospora crassa*.

Abstract

Cellobiose-induced B-galactosidase and B-glucosidase activities of *Neurospora crassa*.

Cellobiosr-induced β -galactosidase and
 β -glucosidase activities of Neurospora crassa.

Three β -galactosidase (β -D-galactoside galactohydrolase: EC 3.2.1.23) activities have been shown in wild type *Neurospora*. One of them is induced by D-cellobiose, has an optimum activity at pH 6 and is precipitated at 50-to-75% saturation with ammonium sulfate (AS) (Perry and Lester 1973 *Biochem Biophys. Res. Commun.* 54: 1476). Since D-cellobiose is an effective inducer for β -glucosidase (β -D-glucoside glucohydrolase: EC 3.2.1.21) activity in *Neurospora*, and since one of the known β -glucosidases, "cellobiase," has optimum activity at pH 6 (Eberhart and Beck 1970 *J. Bacteriol.* 101: 408), we investigated whether or not the pH 6 form of β -galactosidase has any β -glucosidase activity.

Beta-galactosidase and β -glucosidase activities were determined by use of the chromogenic substrates, *o*-nitrophenyl- β -D-galactopyranoside (ONPG) and *p*-nitrophenyl- β -D-glucopyranoside (PNPG). These and the other procedures used in this study (e.g. growth conditions, medium cellobiose concentration, etc.), were described by Perry and Lester (1973 *Biochem Biophys. Res. Commun.* 54: 1476). A unit of enzyme activity is that amount which releases 1.0 μ mole of ONP or PNP per hour at 37°C under the assay conditions. For ion exchange chromatography, diethylaminoethyl (DEAE) cellulose (Celltex D), was equilibrated with 0.01 M potassium phosphate, pH 6.8 containing 0.001 M EDTA, and poured into a 2.2 x 15-cm column (bed volume, 50 ml). Ammonium sulfate (AS) fractions were applied in volume of less than 5 ml and elution was carried out with a linear NaCl gradient (0.025 to 0.25 M) made in 0.01 M potassium phosphate buffer.




Figure 1. DEAE-cellulose chromatography of a resuspended 50-to-75% AS fraction from cellobiose-grown mycelia. The sample applied contained 101 units of β -galactosidase activity and 135 units of β -glucosidase activity. (●) β -galactosidase activity; (○) β -glucosidase activity; (x) absorbance at 280 nm; (----) NaCl concentration.

When extracts of cellobiose-grown mycelia were assayed at pH 6, both β -galactosidase and β -glucosidase activities were detected. The extracts were then subjected to AS fractionation. Very little of the enzyme activities (1% of the β -galactosidase and 4% of the β -glucosidase present in the crude extract) was precipitated by 0.50% AS; whereas, much more (75% of the β -galactosidase and 40% of the β -glucosidase) was precipitated by 50-75% AS. The 50.75% AS-fraction was then separated by chromatography on DEAE-cellulose. The fractions obtained were assayed for both activities (see Fig. 1). β -glucosidase activity appeared in two well-defined peaks (I and II), while β -galactosidase activity appeared in only one peak, which closely coincided with peak I of the β -glucosidase activity. There was, essentially, total recovery of both activities applied to the column. When activities were normalized to the peak maxima, the β -galactosidase peak and the β -glucosidase peak I were superimposable.




Figure 2. Effect of pH on DEAE-cellulose peak-I β -galactosidase activity (\bullet), peak-I β -glucosidase activity (o), and peak-II β -glucosidase activity (x). The number of units used were: peak-I β -galactosidase, 1.30; peak-I β -glucosidase, 1.30; and peak-II β -glucosidase, 1.84. Activities were measured at pH optima.

Fractions corresponding to the two peaks were used to characterize further the β -galactosidase and β -glucosidase enzyme activities. Effects of pH on these activities are shown in Fig. 2. The pH optima of 5 and 6 for peak-II and peak-I β -glucosidase activities are similar to the optima reported for aryl- β -glucosidase (Mahadevan and Eberhart 1964 Arch. Biochem. Biophys. 108: 22) and cellobiase (Eberhart and Beck 1970), respectively. The peak-I β -glucosidase and β -galactosidase cannot be distinguished by the effect of pH upon their activities nor by the use of inhibitors. D-cellobiose or mercuric chloride. Finally, the peak-I activities were tested for their thermal stabilities. At 50°C, β -galactosidase and β -glucosidase activities showed very similar kinetics of inactivation, with half-lives of 3 to 4 min.

In summary, these data show that the cellobiose-induced, pH 6- β -galactosidase of *Neurospora* is associated with a β -glucosidase activity, possibly cellobiase. It is not known at this time whether this is because *Neurospora* possesses a single enzyme activity that has dual specificity for β -galactoside and β -glucoside substrates, or whether there are two separate enzymes that copurify in the procedures used.