

Fungal Genetics Reports

Volume 11

Article 17

A method for disrupting conidia of Neurospora

M. Kapoor

D. Bray

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Kapoor, M., and D. Bray (1967) "A method for disrupting conidia of Neurospora," *Fungal Genetics Reports*: Vol. 11, Article 17. <https://doi.org/10.4148/1941-4765.1982>

This Spores is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

A method for disrupting conidia of Neurospora

Abstract

Method for disrupting conidia

Kapoor, M. and D. Bray. A method for disrupting conidio of Neurospora crassa.

The number of conidia is estimated by determining the absorbance of the suspension with a Klett-Summerson photoelectric colorimeter, using a blue filter. A standard was prepared initially by correlating the Klett reading with the number of conidia, calculated from colony counts obtained by serial dilution followed by plating on sorbose media. A linear relationship between the number of conidio and Klett reading is realized up to an absorbance of 300 Klett units (equivalent to about 6×10^7 conidia). Suspensions at concentrations higher than this do not fall within the linear portion of the standard curve. Our standard has been prepared for conidio harvested from cultures grown at 28°C, for 6 days, using strains pe (Y8743m) (FGSC#38) and wild type 79a (FGSC#533).

Conidio are disrupted by treating a suspension containing not more than 2.1×10^7 conidia per ml (Klett reading of 100) with a Branson, model S-75 sonifier cell disruptor. Figure 1 illustrates the quantity of soluble protein liberated as a function of time of insonation of conidia of strain Y8743m. The sigmoid shape of the curve is due to the release of one fraction of protein immediately following the disruption of the conidial wall, while that associated with the separation of protoplasmic material from the wall is released at a subsequent stage. As much as 95% of the soluble protein is liberated within 15 minutes.

Conidial ATPase (Mg^{++} -requiring) is liberated within the first 5 minutes of insonation, whereas this treatment yields only about 40% of the total soluble protein. Department of Biology, University of Calgary, Calgary, Alberta, Canada.

The following method is routinely used by us for counting and disrupting the conidio for enzyme extraction. A conidial suspension containing $2 - 3 \times 10^7$ conidio per ml is prepared in buffer (phosphate or tris 0.05 M, 5×10^{-4} M in EDTA and 10^{-4} M in β -mercaptoethanol).

