

# Kansas Agricultural Experiment Station Research Reports

Volume 0  
Issue 1 *Cattleman's Day (1993-2014)*

Article 918

1988

## Influence of supplemental grain type on forage utilization by beef steers consuming early summer bluestem

E.S. Vanzant

R.C. Cochran

A.A. Beharka

*See next page for additional authors*

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

 Part of the [Other Animal Sciences Commons](#)

### Recommended Citation

Vanzant, E.S.; Cochran, R.C.; Beharka, A.A.; and Avery, T.B. (1988) "Influence of supplemental grain type on forage utilization by beef steers consuming early summer bluestem," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2321>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1988 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



---

## Influence of supplemental grain type on forage utilization by beef steers consuming early summer bluestem

### Abstract

Supplementation of early summer bluestem with low levels of corn, wheat, or sorghum grain had no effect on forage intake, forage digestibility, or total dry matter digestibility in beef steers, compared with a regimen of no supplementation. Increased total dry matter intakes for the supplemented steers reflected supplement consumption.

### Keywords

Kansas Agricultural Experiment Station contribution; no. 88-363-S; Cattlemen's Day, 1988; Report of progress (Kansas State University. Agricultural Experiment Station and Cooperative Extension Service); 539; Beef; Forage; Steers; Bluestem

### Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

### Authors

E.S. Vanzant, R.C. Cochran, A.A. Beharka, and T.B. Avery

**K****S****U**

---

Influence of Supplemental Grain Type on  
Forage Utilization by Beef Steers Consuming  
Early Summer Bluestem<sup>1</sup>

E.S. Vanzant, R.C. Cochran,<sup>2</sup>  
A.A. Beharka, and T.B. Avery<sup>2</sup>

---

### Summary

Supplementation of early summer bluestem with low levels of corn, wheat, or sorghum grain had no effect on forage intake, forage digestibility, or total dry matter digestibility in beef steers, compared with a regimen of no supplementation. Increased total dry matter intakes for the supplemented steers reflected supplement consumption.

### Introduction

Intensive-early stocking has become an important and common practice among Flint Hills producers. Previous research at the Fort Hays Branch Station has shown that low levels of grain supplementation to cattle under intensive-early stocking programs may produce efficient gains. Additionally, research at Manhattan has shown that steers may be supplemented with up to 4 pounds of sorghum grain per day with only minimal effects on forage utilization. Other research, however, indicates that more rapidly fermentable grains such as corn and wheat have a greater potential for disrupting forage utilization. The purpose of this study was to determine the effects of supplemental corn, wheat, and sorghum grain on the utilization of early summer bluestem by beef cattle.

### Experimental Procedures

Sixteen ruminally cannulated steers (avg. wt., 795 lbs.) of primarily Hereford x Angus breeding were randomly assigned to one of four treatments: 1) Control (no supplement), 2) corn, 3) wheat, or 4) sorghum grain supplements. All supplements were fed at 0.36% of body weight on an as fed basis. Soybean meal was added to the milo and corn supplements to equalize their crude protein contents with that of the wheat supplement. Animals were housed in individual pens. Fresh bluestem range grass, cut and chopped daily, was fed at 15% over each animal's previous 7-days average intake from June 11 to July 9, 1987. Forage and grain offered and forage refusals were weighed and sampled daily, analyzed for dry matter, and stored for future analyses. The trial consisted of a 14-day adaptation period, a 7-day intake measurement period, and a 7-day total fecal collection period.

---

<sup>1</sup>The authors express their sincere appreciation to Mr. Wayne Adolph and Mr. Gary Ritter for their invaluable assistance in conducting this trial.

<sup>2</sup>Dept. of Surgery and Medicine.

## Results and Discussion

There was no difference ( $P>.10$ ) in forage dry matter intake between the control treatment and any of the supplemented treatments (Table 10.1). Therefore, the trend toward increased total dry matter (DM) intakes for the supplemented groups are simply a result of the increased DM fed in the form of a supplement. There were no differences ( $P>.10$ ) in either total DM digestibility or acid detergent fiber digestibility. These results indicate that low levels of corn and wheat, as well as sorghum grain, may be used as supplements for stocker cattle grazing early summer bluestem range without adversely affecting forage utilization. Additional research is needed to determine the efficiency with which supplemental energy is converted to tissue gain in steers grazing early summer bluestem.

Table 10.1. Influence of Supplemental Grain Type on Dry Matter (DM) Intake and Digestibility in Beef Steers Consuming Early Summer Bluestem

Item	Supplement				SE <sup>a</sup>
	None	Corn	Wheat	Sorghum Grain	
Forage DM intake (lb/d)	19.0	19.5	19.2	19.9	.73
Total DM intake (lb/d)	19.0	22.3 <sup>b</sup>	22.0 <sup>b</sup>	22.7 <sup>b</sup>	.74
Forage DM intake (% body wt.)	2.5	2.5	2.4	2.5	.18
Grain DM intake (% body wt.)	--	0.35	0.34	0.35	--
Total DM intake (% body wt.)	2.5	2.8 <sup>b</sup>	2.7 <sup>c</sup>	2.8 <sup>b</sup>	.18
Total DM digestibility (%)	51.4	54.8	56.0	52.1	2.73
Acid Detergent Fiber Digestibility (%)	41.1	43.8	40.5	39.6	4.00

<sup>a</sup>SE = standard error (n=4)

<sup>b</sup>Row means differ from control ( $P<.10$ )

<sup>c</sup>Row mean tends to differ from control ( $P<.15$ )