

# Kansas Agricultural Experiment Station Research Reports

---

Volume 3  
Issue 4 *Turfgrass Research*

Article 4

---

7-2017

## Native Turfgrass Tolerance to Indaziflam Applications

J. Hoyle

*Kansas State University*, jahoyle@ksu.edu

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Horticulture Commons](#)

---

### Recommended Citation

Hoyle, J. (2017) "Native Turfgrass Tolerance to Indaziflam Applications," *Kansas Agricultural Experiment Station Research Reports*: Vol. 3: Iss. 4. <https://doi.org/10.4148/2378-5977.7153>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 7-2017 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



---

## Native Turfgrass Tolerance to Indaziflam Applications

### Abstract

Multiple research trials were conducted from 2014 to 2016 in the Manhattan, KS, area to determine the use of indaziflam in native turfgrass systems. Results indicated that evaluated species are tolerant to indaziflam applications throughout the year.

### Keywords

Native turfgrass, indaziflam, Specticle

### Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

# TURFGRASS RESEARCH 2017



JULY 2017

**K-STATE**  
Research and Extension

Kansas State University  
Agricultural Experiment Station  
and Cooperative Extension Service

K-State Research and Extension is an equal  
opportunity provider and employer.

## Native Turfgrass Tolerance to Indaziflam Applications

*Jared A. Hoyle*

**Summary.** Multiple research trials were conducted from 2014 to 2016 in the Manhattan, KS, area to determine the use of indaziflam in native turfgrass systems. Results indicated that evaluated species are tolerant to indaziflam applications throughout the year.

**Rationale.** Native turfgrass systems contain vegetation of many species and can reduce the need for fertilizer, fuel, and water inputs. Due to diverse flora, weed control applications in native turfgrasses can be problematic.

**Objective.** The objective of the collaborative study is to determine the best use of indaziflam in native turfgrass systems.

**Study Description.** Multiple research trials were conducted in the greater Manhattan, KS, area from 2014 to 2016 to determine native and naturalized turfgrass species' tolerance to indaziflam applications. Evaluated species included little bluestem (*Schizachyrium scoparium*), big bluestem (*Andropogon gerardii*), indian-grass (*Sorghastrum nutans*), and tall fescue (*Schedonorus arundinaceus*). Evaluated Specticle (active ingredient, indaziflam) formulations (200 lb/a) applications included: formulation (Specticle FLO and Specticle G), application method (broadcast, over-the-top and directed spray), application timing (spring, summer, fall, and winter), and vegetation management (mown, non-mown, burned, and non-burned). Additional treatments were Specticle Total (indaziflam + diquat + glyphosate) at 10.6 lb ai/a and 21.2 lb ai/a, Snapshot 2.5 (4 pounds trifluralin per acre + 1 pound isoxaben per acre), Barricade 65WG (0.5 pounds prodiamine per acre), and a non-treated control was included for comparison. Liquid applications were applied using a CO<sub>2</sub> pressurized backpack sprayer calibrated to deliver 40 GPA at 32 psi. Each

View all turfgrass research reports online at: <http://newprairiepress.org/kaesrr>



study site was arranged in a randomized complete block with 4 replications. Plant tolerance was evaluated on a 1-10 scale (1 = significant damage, 6 = minimal acceptable quality, and 10 = quality equal to or better than the non-treated). Phytotoxicity was evaluated on a 0-100% scale. Evaluations were conducted bi-weekly for 6 weeks after each application. Analysis of variance (ANOVA) was performed in SAS 9.4 (SAS Institute Inc., Cary, NC) and means were separated according to Fisher's protected least significant difference (LSD) level at 0.05.

**Results.** Results indicated that evaluated species are tolerant to indaziflam applications throughout the year. Specticle Total applications should be applied in the winter only to little bluestem, big bluestem, and indiagrass. Future research will evaluate additional native species and weed control.



**Figure 1.** Native turfgrass research trial located at Colbert Hills Golf Course in Manhattan, KS.

**K-STATE**  
Research and Extension

Kansas State University  
Agricultural Experiment Station  
and Cooperative Extension Service

