

UNIVERSITY OF BIRMINGHAM

University of Birmingham
Research at Birmingham

Shared decision-making within goal-setting in intermediate care

Rose, Alice; Soundy, Andrew; Rosewilliam, Sheeba

License:

Creative Commons: Attribution-ShareAlike (CC BY-SA)

Document Version

Early version, also known as pre-print

Citation for published version (Harvard):

Rose, A, Soundy, A & Rosewilliam, S 2016, 'Shared decision-making within goal-setting in intermediate care', SRR Summer 2016 Meeting, Coventry, 5/07/16.

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Authors: Alice Rose, University of Birmingham, BSc Hons
 Andy Soundy, University of Birmingham, PhD
 Sheeba Rosewilliam, University of Birmingham, MSc

Background

Shared decision-making in goal-setting

Shared decision-making (SDM) in healthcare is an approach where 'clinicians and patients work together to decide on the best course of action' [1] and is heavily emphasised in recent government papers [1-4]. Involving patients in decisions about their care is important to improve quality of healthcare to one that is safe, effective, person-centred, timely, efficient and equitable [5]. However the latest National Inpatient Survey [6] highlights that patients are still not involved as much as they wish to be in decisions about their care. Within rehabilitation the goal-setting process is suggested to be a key forum for SDM so that patients and professionals collaboratively set rehabilitation goals. Sharing decisions about their goals with patients can increase patient satisfaction [7-8], motivation [9-12] and functional outcomes [8; 13]. Yet, recent empirical evidence suggests that rehabilitation patients have little involvement in making decisions about their goals [14-18].

The research gap

Studies measuring the extent of SDM have traditionally focussed on clinical consultations [19] and have explored clinician perceived barriers to SDM [20]. Further research is required to consider the patient views within community rehabilitation settings.

Therefore, aims of this study are:

- (1) To measure the extent of shared decision making within goal-setting in an intermediate care set up
- (2) To determine if there is variation in perceived involvement between staff and patients in a goal-setting meeting
- (3) To explore the patient-reported barriers to participating in making decisions about their care and rehabilitation



Methodology

Mixed methods approach: explanatory sequential involving two phases: *Phase 1:* goal-setting meetings with patients were observed and SDM within these meetings were scored using the MAPPIN'SDM questionnaire by patients, staff and a trained observer.

Phase 2: semi-structured interviews were conducted with a sub-group of patients involved in phase one to identify patient reported barriers to SDM.

Setting: two community healthcare teams. One team provided in-patient rehabilitation to patients in a community rehabilitation centre and the other provided rehabilitation in the patient's own home.

Sample strategy: *Phase 1 inclusion criteria* – patients with frailty syndromes and rehabilitation staff. Frailty syndromes [21] included falls, immobility, delirium, incontinence and susceptibility to side effects of medication. Rehabilitation staff included physiotherapists, occupational therapists and rehabilitation workers. *Phase 2 inclusion criteria* - patients who had participated in phase one were considered for phase two if they had scored 0 or 1 (strongly disagree/ disagree) more than once, indicating lack of involvement in decisions about their goals on the MAPPIN'SDM questionnaire.

Ethical approval: granted by the North West NRES Committee (15/ NW/0688)

Data Analysis: Descriptive statistics for patient demographics and questionnaire data were calculated. Since the data was ordinal and patient data was skewed Welch tests were carried out on each sub question of the questionnaires to find out if there were overall differences in the responses between the three groups. If a significant difference was found, Games-Howell tests were undertaken to specifically show where these significant differences were found. Thematic analysis was carried out for the qualitative data from interviews.

Demographics	Frequency
Sample size	40
Gender: male (female)	9 (31)
Age group	
60 - 69	2
70 - 79	10
80 - 89	17
90 - 99	11
Ethnicity: White British	40
Frailty Syndrome	
Falls	22
Immobility	17
Incontinence	1
Side effect medications	0
Delirium	0
Falls history (past 12 months)	
0	9
1	14
2	10
3	4
4	1
5+	2

Question Key Message	Welch significance (p=0.003)
1a. Patient's problems discussed	0.002
1b. Patient understands problems	0.510
2a. Patient told their opinion important	0.004
2b. Patient happy their opinion important	0.000
3a. Rehab options discussed	0.000
3b. Patient understands rehab options	0.000
4a. Advantages/disadvantages rehab discussed	0.000
4b. Patient understands advan/disadvantages	0.000
5a. Patient's expectations/fears discussed	0.213
5b. Expectations/fears taken into account	0.499
6a. Staff check they understand patient	0.745
6b. Staff understand patient	0.010
7a. Language used made sense to patient	0.030
7b. Patient has opportunity to ask questions	0.000
8a. Goals decided in meeting	0.000
8b. Patient is clear on their goals	0.001
9a. Discussion of action plan	0.065
9b. Patient understands action plan	0.599

Welch test for significant difference between groups

Preliminary Findings

Phase 1: The Welch tests found significant differences between patient, observer and staff answers for MAPPIN questions 1a, 2b-4b and 7b-8b. Patient and staff only significantly disagreed on one out of eighteen questions i.e. *about whether the patient's problems were discussed in the GSM*. The patient and observer disagreed on six out of the eighteen questions.

Phase 2: Pre-defined themes were used from a systematic review of patient-reported barriers and facilitators to SDM [22].

References

- [1] The King's Fund. Making shared decision-making a reality: no decision about me without me. London: The King's Fund; 2011
- [2] Department of Health. Equity and excellence: liberating the NHS. London: Department of Health; 2010
- [3] NHS England. Five year forward view. London: NHS England; 2014
- [4] The Health Foundation. Helping measure person-centred care. London: The Health Foundation; 2014
- [5] The Health Foundation. Evidence: Helping people share decision making. London: The Health Foundation, 2012
- [6] Care Quality Commission. Inpatient survey 2015. London: Care Quality Commission; 2015
- [7] Byrnes, M, Beilby, J, Ray, P, McLennan, R, Ker, J, Schug, S. Patient-focused goal planning process and outcome after spinal cord injury rehabilitation: quantitative and qualitative audit. Clin Rehabil. 2012;26(12):1141-9.
- [8] Turner-Stokes, L, Rose, H, Ashford, S, Singer, B. Patient engagement and satisfaction with goal planning: Impact on outcome from rehabilitation. Int J Ther Rehabil. 2015;22(5):210-6.
- [9] Van De Weyer, R, C. Ballinger, C, Playford, E, D. Goal setting in neurological rehabilitation: staff perspectives. Disabil Rehabil. 2010;32(17):1419-27.
- [10] Leach, E, Cornwell, P, Fleming, J, Haines, T. Patient centered goal-setting in a subacute rehabilitation setting. Disabil Rehabil. 2010;32(2):159-72
- [11] Scobbie, L, McLean, D, Dixon, D, Duncan, E, Wyke, S. Implementing a framework for goal setting in community based stroke rehabilitation: a process evaluation. BMC Health Serv Res. 2013; 13(1):1-13
- [12] Brown, M, Leacock, W, McPherson, K, M. Dean, S, G. Reed, K, Weatherall, M, Taylor, W, J. Survival, momentum, and things that make me "me": patients' perceptions of goal setting after stroke. Disabil Rehabil. 2014;36(12):1020-6
- [13] Dalton, C, Farrell, R, De Souza, A, Wujanto, E, McKenna-Slade, A, Thompson, S, Liu, C, Greenwood, R. Patient inclusion in goal setting during early inpatient rehabilitation after acquired brain injury. Clin Rehabil. 2012;26(2):165-73.
- [14] Maitra, K, K. Erway, F. Perception of client-centered practice in occupational therapists and their clients. Am J Occup Ther. 2006;60(3):298-310.
- [15] Young, C, A. Manmathan, G, P. Ward, J, C. Perceptions of goal setting in a neurological rehabilitation unit: a qualitative study of patients, carers and staff. J Rehabil Med. 2008;40(3):190-4.
- [16] Leacock, W, M, M. Dean, S, G. Siegert, R, J. McPherson, K, M. Navigating patient-centered goal setting in inpatient stroke rehabilitation: How clinicians control the process to meet perceived professional responsibilities. Patient Educ Couns. 2011;85(2):206-13
- [17] Schoeb, V, Staffoni, L, Parry, R, Pilnick, A. "What do you expect from physiotherapy?": a detailed analysis of goal setting in physiotherapy. Disabil Rehabil. 2014;36(20):1679-86.
- [18] Rosewilliam, S, Sintler, C, Pandyan, A, D, Skellon, J, Roskell, C, A. Is the practice of goal-setting for patients in acute stroke care patient-centred and what factors influence this? A qualitative study. Clin Rehabil. 2015;7:1-12
- [19] Couët, N, Desroches, S, Robitaille, H, et al. Assessments of the extent to which health-care providers involve patients in decision making: a systematic review of studies using the OPTION instrument.
- [20] Gravel, K, Légaré, F, Graham, I. Barriers and facilitators to implementing shared decision-making in clinical practice: a systematic review of health professionals' perceptions. Implement Sci. 2006;1(16):1-15
- [21] British Geriatrics Society. (2014) Fit for Frailty. London: BGS.
- [22] Joseph-Williams, N, Elwyn, G, Edwards, A. Knowledge is not power for patients: a systematic review and thematic synthesis of patient-reported barriers and facilitators to shared decision making. Patient Educ Couns. 2014;94(3):291-309