

# UNIVERSITY OF BIRMINGHAM

University of Birmingham  
Research at Birmingham

## Traumatic injury and exposure to mitochondrial-derived damage associated molecular patterns suppresses neutrophil extracellular trap formation

Hazeldine, Jon; Harrison, Paul; Dinsdale, Robert; Lord, Janet

DOI:

[10.3389/fimmu.2019.00685](https://doi.org/10.3389/fimmu.2019.00685)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Hazeldine, J, Harrison, P, Dinsdale, R & Lord, J 2019, 'Traumatic injury and exposure to mitochondrial-derived damage associated molecular patterns suppresses neutrophil extracellular trap formation', *Frontiers in immunology*. <https://doi.org/10.3389/fimmu.2019.00685>

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**

Checked for eligibility: 04/04/2019

**General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

**Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

1           **Traumatic Injury and Exposure to Mitochondrial-derived**  
2           **Damage Associated Molecular Patterns Suppresses Neutrophil**  
3           **Extracellular Trap Formation.**

4 Jon Hazeldine<sup>1,2,\*</sup>, Robert J. Dinsdale<sup>1,3</sup>, Paul Harrison<sup>1,3</sup> and Janet M. Lord<sup>1,2,3</sup>.

5  
6 <sup>1</sup>Institute of Inflammation and Ageing, University of Birmingham, Birmingham, United  
7 Kingdom.

8 <sup>2</sup>National Institute for Health Research Surgical Reconstruction and Microbiology Research  
9 Centre, Queen Elizabeth Hospital Birmingham, Birmingham, United Kingdom.

10 <sup>3</sup>Scar Free Foundation Birmingham Centre for Burns Research, Queen Elizabeth Hospital  
11 Birmingham, Birmingham, United Kingdom.

12  
13 **Correspondence:**

14 Dr Jon Hazeldine

15 [j.hazeldine@bham.ac.uk](mailto:j.hazeldine@bham.ac.uk)

16  
17 **Short running title:** Neutrophil dysfunction post-trauma

18  
19 **Keywords:** Neutrophils, Neutrophil extracellular traps, Trauma, Mitochondrial-derived  
20 DAMPs, Immune suppression.

26 **Abstract**

27 Major traumatic injury induces significant remodelling of the circulating neutrophil pool and  
28 loss of bactericidal function. Although a well-described phenomenon, research to date has  
29 only analysed blood samples acquired post-hospital admission, and the mechanisms that  
30 initiate compromised neutrophil function post-injury are therefore poorly understood. Here,  
31 we analysed pre-hospital blood samples acquired from 62 adult trauma patients (mean age 44  
32 years, range 19-95 years) within 1-hour of injury (mean time to sample 39 minutes, range  
33 13-59 minutes). We found an immediate impairment in neutrophil extracellular trap (NET)  
34 generation in response to phorbol 12-myristate 13-acetate (PMA) stimulation, which  
35 persisted into the acute post-injury phase (4-72 hours). Reduced NET generation was  
36 accompanied by reduced reactive oxygen species production, impaired activation of mitogen-  
37 activated protein kinases and a reduction in neutrophil glucose uptake and metabolism to  
38 lactate. Pre-treating neutrophils from healthy subjects with mitochondrial-derived damage-  
39 associated molecular patterns (mtDAMPs), whose circulating levels were significantly  
40 increased in our trauma patients, reduced NET generation. This mtDAMP-induced  
41 impairment in NET formation was associated with an N-formyl peptide mediated activation  
42 of AMP-activated protein kinase (AMPK), a negative regulator of aerobic glycolysis and  
43 NET formation. Indeed, activation of AMPK via treatment with the AMP-mimetic AICAR  
44 significantly reduced neutrophil lactate production in response to PMA stimulation, a  
45 phenomenon that we also observed for neutrophils pre-treated with mtDAMPs. Furthermore,  
46 the impairment in NET generation induced by mtDAMPs was partially ameliorated by pre-  
47 treating neutrophils with the AMPK inhibitor compound C. Taken together, our data  
48 demonstrate an immediate trauma-induced impairment in neutrophil anti-microbial function  
49 and identify mtDAMP release as a potential initiator of acute post-injury neutrophil  
50 dysfunction.

51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75

## 76 Introduction

77

78 Major injury induces significant phenotypic and functional remodelling of the peripheral  
79 neutrophil pool, attributable in part to the emergence into circulation of immature  
80 granulocytes (IGs) and highly mature neutrophil subsets (*Hampson et al. 2017; Hazeldine et*  
81 *al. 2014; Hazeldine et al. 2017; Liefeld et al. 2016; Pillay et al. 2012*). Alongside changes  
82 in the surface expression of adhesion molecules and chemokine receptors (*Johansson et al.*  
83 *2011; Visser et al. 2011; Visser et al. 2012*), immediate and prolonged impairments in  
84 phagocytosis (*Hampson et al. 2017; Liao et al. 2013*), reactive oxygen species (ROS)  
85 production (*Hampson et al. 2017; Hazeldine et al. 2017*) and chemotaxis (*Li et al. 2015;*  
86 *Tarlowe et al. 2003*) have been reported for neutrophils isolated from critically-injured  
87 patients. Whilst understanding of trauma-induced changes in neutrophil intracellular  
88 bactericidal function is well developed, few studies have investigated the impact of injury on  
89 the extracellular defensive mechanisms of neutrophils and how soon after injury, any  
90 compromise occurs.

91

92 Comprised of a DNA backbone decorated with granular and cytosol-derived peptides and  
93 enzymes, neutrophil extracellular traps (NETs) are an extracellular anti-microbial defence  
94 mechanism deployed to prevent pathogen dissemination (*Brinkmann and Zychlinsky 2012*).  
95 Studies that have examined NET generation post-injury have reported an increase in *ex vivo*  
96 NET formation by resting neutrophils (*Itagaki et al. 2015; Li et al. 2015*), a hyperactivity that  
97 may reflect their *in vivo* exposure to high mobility group box-1 (HMGB-1) and interleukin  
98 (IL)-33, two NET-inducing alarmins whose circulating concentrations are significantly  
99 increased post-injury (*Cohen et al. 2009; Tadie et al. 2013; Xu et al. 2017*). However, in  
100 terms of stimulus-induced NET generation, comparable (*Itagaki et al. 2015*) or reduced  
101 (*Hampson et al. 2017; Li et al. 2015*) NET production in response to stimulation with  
102 phorbol 12-myristate 13-acetate (PMA) has been reported post- trauma. Of these studies, only  
103 one performed quantitative analysis (*Hampson et al. 2017*), and neither study that reported a  
104 post-injury reduction in NET formation investigated the mechanism(s) responsible (*Hampson*  
105 *et al. 2017; Li et al. 2015*).

106

107 Activation of three non-redundant molecular processes underpin PMA-induced NET  
108 generation. Produced by the multi-subunit enzyme nicotinamide adenine dinucleotide  
109 phosphate (NADPH) oxidase, ROS generation is essential for the process of chromatin  
110 decondensation that precedes NET release (*Fuchs et al. 2007; Kirchner et al. 2012*).  
111 Occurring prior to (*Hakkim et al. 2011*) or following (*Keshari et al. 2013*) ROS production,  
112 activation of the mitogen activated protein kinases p38 and extracellular signal regulated  
113 kinase 1/2 (ERK 1/2), triggers NET formation by inhibiting caspase activation and increasing  
114 expression of the pro-survival protein Mcl-1, thus promoting NET production over the  
115 induction of apoptosis (*Hakkim et al. 2011; Keshari et al. 2013*). Finally, glycolysis is a  
116 fundamental metabolic requirement for PMA-induced NET formation, with the uptake and  
117 breakdown of extracellular glucose a necessity for the process of DNA expulsion (*Rodriguez-*  
118 *Espinosa et al. 2015*).

119

120 Trauma-associated tissue damage results in the release into the circulation of damage-  
121 associated molecular patterns (DAMPs), a collection of cytosolic, mitochondrial and nuclear-  
122 derived proteins and DNA (*Cohen et al. 2009; Kaczmarek et al. 2018; Zhang et al. 2010*).  
123 Whilst renowned for their role in immune activation (*Hazeldine et al. 2015; Zhang et al.*  
124 *2010*), data are emerging that suggests mitochondrial-derived DAMPs (mtDAMPs), which  
125 include N-formylated peptides and mitochondrial DNA (mtDNA), possess immune tolerising

126 properties. For instance, it has been shown *in vitro* that monocytes pre-exposed to mtDNA  
127 (*Fernandez-Ruiz et al. 2014*) and neutrophils pre-treated with whole mtDAMP preparations  
128 (*Zhang et al. 2010*) exhibit impaired cytokine production and calcium mobilisation  
129 respectively upon secondary stimulation. Furthermore, a significant reduction in stimulus-  
130 induced ROS production and transmigration was reported for neutrophils pre-exposed to  
131 bacterial-derived or synthetic N-formylated peptides (*Gabl et al. 2018; Kaczmarek et al.*  
132 *2018; Li et al. 2015*), both of which signal through the same formyl peptide receptor (FPR) as  
133 mitochondrial-derived formyl peptides. Based on these observations, the concept of  
134 mtDAMP-induced tolerance has been coined and proposed to be a potential mechanistic  
135 explanation for the state of peripheral neutrophil dysfunction that develops in the aftermath of  
136 major trauma (*Kaczmarek et al. 2018; Li et al. 2015*).

137

138 Here, in a prospective observational study of trauma patients, we have performed for the first  
139 time a quantitative assessment of NET production during the pre-hospital, ultra-early ( $\leq 60$   
140 minutes) and acute (4-72 hours) post-injury phases, and assessed the impact that major injury  
141 has on the molecular processes and signalling pathways that underpin PMA-induced NET  
142 generation. Furthermore, based on the emerging concept of mtDAMP-induced tolerance, we  
143 have investigated whether pre-exposing neutrophils isolated from healthy subjects to  
144 mtDAMPs *in vitro* results in altered NET generation upon secondary stimulation with PMA  
145 and the mechanisms involved.

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176 **Materials and methods**

177

178 **Study design and setting**

179

180 This manuscript presents data acquired from subjects enrolled into the Brain Biomarkers after  
181 Trauma Study, an ongoing prospective longitudinal observational study of adult trauma  
182 patients conducted at a single Major Trauma Centre site in the UK (University Hospitals  
183 Birmingham NHS Foundation Trust, Birmingham). Ethical approval for the study was  
184 granted by the North Wales Research Ethics Committee - West (REC reference:  
185 13/WA/0399, Protocol Number: RG\_13-164).

186

187 Patient enrolment began in the pre-hospital setting, where on a 24/7 basis between March  
188 2016 and October 2018, emergency care teams acquired blood samples from adult trauma  
189 patients ( $\geq 18$  years) with a suspected injury severity score (ISS)  $\geq 8$  within 1-hour of injury  
190 (defined as the time of phone call to emergency services). In the pre-hospital setting, blood  
191 samples were not taken from patients who were deemed unlikely to survive transportation to  
192 hospital. Post admission, patients were excluded if they were aged  $< 18$  years, if pre-hospital  
193 blood samples had been acquired  $> 1$  hour post-injury and if clinical assessments confirmed  
194 either an ISS  $< 8$  or a previous diagnosis of neuro-degenerative disease. No patients received  
195 blood products in the pre-hospital setting.

196

197 **Capacity and consent**

198

199 Due to the nature of injuries sustained, patients were unlikely to provide informed consent for  
200 their participation at the time of study enrolment. Consequently, patient recruitment was  
201 performed under the guidance of the Mental Health Capacity Act 2005 for research in  
202 emergency situations and the Declaration of Helsinki. For patients who lacked capacity, an  
203 agreement for study participation was sought from a legal consultee (family member or  
204 clinician not directly involved in the study), with written consent obtained from the patient  
205 once they regained capacity. In instances where the patient did not regain capacity, data were  
206 retained in accordance with the agreement of the legal consultee.

207

208 **Blood sampling**

209

210 In the pre-hospital environment, peripheral venous blood samples were acquired during the  
211 intravenous cannulation of patients or by venepuncture. Once taken, blood tubes were stored  
212 at room temperature (RT) until arrival at hospital, where analysis began within 1-hour by a  
213 single laboratory researcher on a 24/7 basis. Additional blood samples were acquired 4-12  
214 and 48-72 hours post-injury. At all three time points, blood samples were collected into BD  
215 Vacutainers<sup>®</sup> (BD Biosciences, Oxford, UK) containing ethylenediaminetetraacetic acid, z-  
216 serum clotting activator or 1/10 volume of 3.2% trisodium citrate. Full blood counts were  
217 performed using a Sysmex XN-1000 haematology analyser (Sysmex UK, Milton Keynes,  
218 UK) that measures a white cell differential and IGs, which are defined as promyelocytes,  
219 myelocytes and metamyelocytes. The analyser uses fluorescence dyes that label intracellular  
220 DNA and RNA, with the intensity of the fluorescence signal directly proportional to the  
221 nucleic acid content of the cell. Due to their higher RNA content, IGs are discriminated from  
222 mature neutrophils via their stronger fluorescence signal. Daily internal quality control  
223 measurements (XN check, Sysmex UK) and monthly external quality control samples  
224 (UKNEQAS, Watford, UK) ensured instrument performance.

225

226 Sixty-seven adults (mean age 31 years, range 18-80) served as a cohort of healthy controls  
227 (HCs). HCs were volunteers who were not taking any regular medication for a diagnosed  
228 illness and did not have an acute episode of infection prior to the time of sampling. The  
229 recruitment of HCs was carried out in accordance with the ethical approval granted by the  
230 University of Birmingham Research Ethics Committee (Ref: ERN\_12-1184) with written  
231 informed consent from all subjects. All subjects gave written informed consent in accordance  
232 with the Declaration of Helsinki.

233

### 234 **Preparation of mtDAMPs and mtDNA**

235

236 MtDNA and mtDAMPs were prepared from mitochondria isolated from the K562 tumour  
237 cell line (ATCC<sup>®</sup>, Teddington, Middlesex, UK) as described previously (*Hazeldine et al.*  
238 *2015*). MtDNA concentration and protein content within mtDAMPs were determined by  
239 spectrophotometry (Nanodrop 2000; Thermo Fisher Scientific, Paisley, UK) and preparations  
240 stored at -80°C prior to use.

241

### 242 **Neutrophil isolation and treatment**

243

244 Neutrophils were isolated by Percoll density gradient centrifugation (Scientific Lab Supplies,  
245 Nottingham, UK) with cell purity, which was routinely  $\geq 99\%$ , determined using a Sysmex  
246 XN-1000 haematology analyser. Neutrophils were re-suspended at concentrations of  
247  $1-10 \times 10^6$  /ml in phenol red free or phenol red containing RPMI-1640 media supplemented  
248 with 2 mM L-glutamine, 100 U/ml penicillin and 100  $\mu\text{g/ml}$  streptomycin (GPS; Sigma-  
249 Aldrich, Dorset, UK), phenol red free or phenol red containing RPMI-1640 media  
250 supplemented with GPS and 10% heat-inactivated fetal calf serum (HI-FCS; hereafter  
251 referred to as complete medium (CM); Sigma-Aldrich), glucose free RPMI-1640 media  
252 supplemented with GPS (Gibco, Fisher Scientific UK Ltd, Loughborough, UK), Hank's  
253 balanced salt solution (HBSS) supplemented with calcium and magnesium (hereafter referred  
254 to as HBSS<sup>Ca<sup>2+</sup></sup>; Gibco, Life Technologies, Cheshire, UK) or HEPES buffer containing 1 mM  
255 Ca<sup>2+</sup>.

256

257 For mtDAMPs and mtDNA experiments, neutrophils were pre-treated for 15 minutes  
258 (37°C/5% CO<sub>2</sub>) with 40 or 100  $\mu\text{g/ml}$  mtDAMPs or mtDNA prior to secondary stimulation.  
259 Prior to inclusion in transmigration, lactate and ROS assays, neutrophils were pelleted,  
260 supernatants removed and cells resuspended in specified media. To inhibit FPR-1 signalling,  
261 neutrophils were treated for 60 minutes (37°C/5% CO<sub>2</sub>) with 2.5  $\mu\text{M}$  cyclosporin H (CsH;  
262 Abcam, Cambridge, UK) or vehicle control, prior to mtDAMP stimulation. For compound C  
263 experiments, neutrophils were treated for 60 minutes with 200  $\mu\text{M}$  compound C (Sigma-  
264 Aldrich) or vehicle control prior to mtDAMP and PMA treatment. To inhibit calcium-  
265 calmodulin-dependent protein kinase kinases (CaMKKs), neutrophils were incubated for 60  
266 minutes (37°C/5% CO<sub>2</sub>) with 2.5  $\mu\text{M}$  STO-609 (Sigma-Aldrich) or vehicle control prior to  
267 mtDAMP treatment. To induce AMP-activated protein kinase (AMPK) signalling,  
268 neutrophils were treated for 60 minutes with 1 mM AICAR (Sigma-Aldrich) prior to PMA  
269 stimulation.

270

### 271 **Ex vivo NET formation**

272

273 Neutrophils ( $2 \times 10^5$  in phenol red free or phenol red containing RPMI + GPS or glucose free  
274 RPMI-1640 media supplemented with GPS) were stimulated with 25 nM PMA (Sigma-  
275 Aldrich) for 3 hours at 37°C/5% CO<sub>2</sub>. Post-stimulation, supernatants were collected and

276 centrifuged at 2,200 x g for 10 minutes at 4°C, after which the DNA content of cell-free  
277 supernatants was analysed. Briefly, 100 µl aliquots of cell-free supernatant were incubated  
278 with 1 µM SYTOX Green dye (Life Technologies) for 10 minutes at RT. Fluorescence was  
279 measured using a BioTek Synergy 2 fluorometric plate reader (NorthStar Scientific Ltd,  
280 Sandy, UK) with excitation and emission set at 485 nm and 528 nm respectively. In our  
281 trauma-based studies, DNA quantification was performed using a λ-DNA standard curve  
282 (Fisher Scientific) with PMA-induced NET generation presented as DNA concentration after  
283 subtracting the readings obtained from untreated controls. For mtDAMP experiments,  
284 background fluorescence values acquired from SYTOX Green staining of mtDAMPs in the  
285 absence of neutrophils were subtracted from test readings, with NET production expressed as  
286 a fold increase above untreated controls.

287

### 288 **Visualisation of NETs by fluorescence microscopy**

289

290 2 x 10<sup>5</sup> neutrophils in phenol red free or phenol red containing RPMI + GPS or glucose free  
291 RPMI-1640 media supplemented with GPS were seeded onto glass coverslips and incubated  
292 for 30 minutes at 37°C/5% CO<sub>2</sub> to allow for cell adherence. Following a 3-hour stimulation  
293 with 25 nM PMA (37°C, 5% CO<sub>2</sub>), samples were fixed for 30 minutes with 4%  
294 paraformaldehyde (37°C, 5% CO<sub>2</sub>), washed three times in phosphate buffered saline (PBS)  
295 and permeabilised with 0.1% Triton X-100 (Sigma-Aldrich). DNA was then stained with  
296 1 µM SYTOX Green dye for 5 minutes, after which slides were washed once in PBS,  
297 mounted in fluoromount medium and visualized using a LEICA DMI 6000 B microscope  
298 (LEICA, Milton Keynes, UK) at x20 or x40 objective.

299

### 300 **ROS production**

301

302 For *ex vivo* analysis of neutrophils isolated from trauma patients, ROS generation was  
303 assessed by lucigenin-amplified chemiluminescence. The effect of mtDAMP pre-treatment  
304 on ROS production was examined using luminol-amplified chemiluminescence. In both  
305 instances, 100 µl aliquots of neutrophils (1x10<sup>6</sup>/ml in HBSS<sup>+/+</sup>) were dispensed into wells of  
306 a 96-well white-bottomed flat plate (BD Biosciences), pre-coated with PBS/2% BSA, that  
307 contained 25 µl of luminol (pH 7.3; final concentration 100 µM; Sigma-Aldrich) or lucigenin  
308 (final concentration 200 µM; Sigma-Aldrich) and 50 µl HBSS<sup>+/+</sup>. Neutrophils were then  
309 stimulated with 25 nM PMA or vehicle control, after which ROS generation was assessed at  
310 1-minute intervals for 180 minutes using a Berthold Centro LB 960 luminometer (Berthold  
311 Technologies, Hertfordshire, UK). Experiments were performed in quadruplicate, with ROS  
312 production measured as relative light units and calculated as area under the curve (AUC).

313

### 314 **Measurement of lactate concentration in cell-free culture supernatants**

315

316 Neutrophils (2x10<sup>6</sup> in phenol red free RPMI + GPS) were stimulated for 1, 2 or 3 hours  
317 (37°C, 5% CO<sub>2</sub>) with 25 nM PMA or vehicle control. At each time-point, cell-free  
318 supernatants were harvested (800 x g, 5 mins, 4°C) and samples stored at -80°C prior to  
319 analysis. Lactate concentration in 25 µl aliquots of supernatant was determined using a  
320 commercially available lactate assay kit according to manufacturer's instructions (Sigma-  
321 Aldrich).

322

323

324

325



326 **Glucose uptake assay**

327

328 Following a 15-minute rest period at 37°C/5% CO<sub>2</sub>, neutrophils (1x10<sup>6</sup> in RPMI-1640 media  
329 without glucose) were stimulated for 60 minutes (37°C/5% CO<sub>2</sub>) with 25 nM PMA or  
330 vehicle. With 10 minutes of the stimulation period remaining, the fluorescent glucose  
331 analogue 2-N-7-nitrobenzen-2-oxa-1,3-diazol-4-yl amino-2-deoxyglucose (2-NBDG; Thermo  
332 Fisher) at a final concentration of 100 µM was added. Post-incubation, samples were washed  
333 and cells re-suspended in glucose free RPMI in preparation for flow cytometric analysis,  
334 which was performed on a CyAn<sub>ADP</sub><sup>TM</sup> bench top cytometer (Dako, Cambridgeshire, UK).  
335 10,000 neutrophils were collected and FL1 mean fluorescence intensity values recorded.

336

337 **Neutrophil transmigration**

338

339 Neutrophils (1x10<sup>7</sup>/ml) in HEPES buffer containing 1 mM Ca<sup>2+</sup> were incubated for 30  
340 minutes in a 37°C water bath with 3 µg/ml calcein-acetoxmethyl ester (calcein-AM, Fisher  
341 Scientific), after which cells were pelleted, supernatants removed and neutrophils re-  
342 suspended at 1x10<sup>7</sup>/ml in phenol red free CM. A total of 1x10<sup>6</sup> neutrophils were dispensed  
343 into the upper chambers of polycarbonate membrane cell culture inserts with 3 µM pores  
344 (Corning, New York, USA) that had been pre-loaded into wells of a 24-well flat bottomed  
345 plate (BD Biosciences) containing pre-warmed phenol red free CM and 1 nM LTB<sub>4</sub> (R and D  
346 Systems, Abingdon, UK). Following a 90-minute incubation at 37°C, cell culture inserts were  
347 removed and plates read immediately for calcein fluorescence using a BioTek Synergy 2  
348 fluorometric plate reader with excitation and emission set at 485 nm and 528 nm respectively.  
349 Fluorescence intensities were converted into neutrophil numbers via the use of a standard  
350 curve that was generated from calcein-AM loaded neutrophils that had been incubated  
351 alongside the test samples in the conditions described above. The number of neutrophils  
352 measured in media in which no chemokine was added was subtracted from the numbers  
353 calculated for wells that contained 1 ng/ml LTB<sub>4</sub> in order to determine specific chemokine-  
354 mediated migration.

355

356 **Assessment of neutrophil phenotype**

357

358 Freshly isolated neutrophils (1x10<sup>5</sup> in CM) were stimulated with 100 µg/ml mtDAMPs or  
359 vehicle control for 15 minutes at 37°C in a humidified 5% CO<sub>2</sub> atmosphere. Post-treatment,  
360 samples were stained on ice for 20 minutes with the following mouse anti-human monoclonal  
361 antibodies or their concentration-matched isotype controls: 2 µg/ml fluorescein  
362 isothiocyanate (FITC)-labelled CD62L (clone DREG56; eBioscience, Hatfield, UK); 1 µg/ml  
363 CXCR1-FITC (clone eBIO8F1-1-4; eBioscience); 0.5 µg/ml R-phycoerythrin (PE)-labelled  
364 CXCR2-PE (clone eBio5E8-C7-F10; eBioscience) or 2.5 µg/ml allophycocyanin (APC)-  
365 labelled CD11b (clone ICRF44, BioLegend, London, UK). Post incubation, cells were  
366 pelleted (250 x g, 5 minutes, 4°C), supernatants discarded and neutrophils washed once in  
367 PBS/1%BSA. Following resuspension in PBS, samples were transferred to polypropylene  
368 FACS tubes for flow cytometric analysis, which was performed on an AccuriC6<sup>TM</sup> bench top  
369 cytometer (BD Biosciences). 10,000 neutrophils, gated according to their forward scatter  
370 (FS)/sideward scatter (SS) properties, were acquired for analysis, where receptor expression  
371 was measured as median fluorescence intensity (MedFI).

372

373

374

375

376 **Cell signalling measurements**

377

378 To determine signalling through AMPK and MAPK pathways, cell lysates prepared from  
379  $2 \times 10^6$  resting neutrophils,  $1 \times 10^6$  neutrophils stimulated with either 25 nM PMA or 100  $\mu\text{g/ml}$   
380 mtDAMPs for 2-90 minutes ( $37^\circ\text{C}/5\% \text{CO}_2$ ), or  $2 \times 10^6$  neutrophils stimulated with 100  $\mu\text{g/ml}$   
381 mtDAMPs for 5 minutes following 1-hour pre-treatment with 2.5  $\mu\text{M}$  STO-609 or 2.5  $\mu\text{M}$   
382 CsH were separated on 10 or 12% SDS-polyacrylamide gels. Following protein transfer to  
383 polyvinylidene difluoride membranes (Bio-Rad, Hertfordshire, UK), blots were probed  
384 overnight at  $4^\circ\text{C}$  with rabbit anti-human antibodies (Cell Signalling Technology,  
385 Massachusetts, USA) directed against phosphorylated AMPK (pAMPK), phosphorylated  
386 ERK1/2 (pERK1/2), phosphorylated P38 (pP38), lactate dehydrogenase A (LDHA) or  
387 pyruvate kinase (PKM2). Post incubation, membranes were washed in tris-buffered saline  
388 containing 0.001% tween (TBST) and incubated for 1 hour at RT with a goat anti-rabbit  
389 secondary antibody conjugated to horse radish peroxidase (HRP; diluted 1:4000 in TBST; GE  
390 Healthcare, Buckinghamshire, UK). HRP activity was detected using enhanced  
391 chemiluminescence (Bio-Rad). To confirm equal loading of proteins, blots were probed with  
392 antibodies against total ERK 1/2, total P38 (1:1000; Cell Signalling Technology) or  $\beta$ -actin  
393 (1:5000, GeneTex, California, USA). Densitometry analysis was performed using Image J  
394 software (National Institutes of Health, Bethesda, MD, USA).

395

396 **Enzyme-linked immunosorbent assays (ELISAs)**

397

398 Serum was prepared from blood collected into BD vacutainers containing z-serum clotting  
399 activator. Following a 30-minute incubation at RT, blood samples were centrifuged at  $1,620 \times$   
400  $g$  for 10 minutes at  $4^\circ\text{C}$ , after which serum was aliquoted and stored at  $-80^\circ\text{C}$  until analysed.  
401 ELISAs to measure serum concentrations of HMGB-1 (IBL International, Hamburg,  
402 Germany), mitochondrial encoded NADH dehydrogenase 6 (ND6; MyBioSource, San Diego,  
403 California, USA) and IL-33 (R and D Systems) were performed in accordance with  
404 manufacturer's instructions.

405

406 **Statistical analyses**

407

408 Statistical analyses were performed using GraphPad Prism<sup>®</sup> software (GraphPad Software  
409 Ltd, California, USA). Data distribution was examined using the Kolmogorov-Smirnov or  
410 Shapiro-Wilk normality test. For data that followed a normal distribution, paired student T-  
411 tests, a repeated measures ANOVA with Bonferroni multiple comparison post hoc test or a  
412 one way ANOVA with Dunnett's multiple comparison post hoc test were performed. For  
413 non-normally distributed data, a Wilcoxon matched-pairs signed rank test, a Friedman test  
414 with Dunn's multiple comparison post hoc test or a Kruskal-Wallis with Dunn's multiple  
415 comparison post hoc test was performed. For box and whisker plots, whiskers represent  
416 minimum and maximum values. Statistical significance was accepted at  $p \leq 0.05$ .

417

418

419

420

421

422

423

424

425

## 426 **Results**

427

### 428 **Patient enrolment and demographics**

429

430 1,070 adult trauma patients were screened for study inclusion, with 87 subjects enrolled into  
431 the study (Supplementary Figure 1). Of these, 62 patients with a mean age of 44 years (range  
432 19-95 years) and mean injury severity score of 26 (range 9-57) had their immune function  
433 analysed (Table 1). The mean time of pre-hospital blood sampling was 39 minutes post-injury  
434 (range 13-59 minutes).

435

### 436 **PMA-induced NET production is impaired post-trauma**

437

438 Compared to neutrophils isolated from HCs, neutrophils acquired from trauma patients within  
439 1-hour of injury exhibited significantly enhanced basal NET generation (Figure 1A), a  
440 hyperactivity that was accompanied by significantly elevated serum concentrations of  
441 HMGB-1 (Figure 1B) and IL-33 (Figure 1C). By the 4-12 and 48-72 hour post-injury time  
442 points, a significant reduction in basal NET production was observed (Figure 1A). In  
443 response to stimulation with PMA, patient neutrophils released significantly less DNA at all  
444 three sampling time-points when compared to HCs (Figure 1D). Fluorescence microscopy  
445 confirmed the impairment in NET generation (Figure 1E).

446

### 447 **ROS production in response to PMA stimulation is reduced in the acute post-injury 448 phase**

449

450 ROS generation is a non-redundant event in NET formation (*Fuchs et al. 2007*). Having  
451 observed trauma-induced alterations in both basal and stimulated NET formation, we  
452 examined the effect of injury on ROS production. In the absence of stimulation, patient  
453 neutrophils isolated 48-72 hours post-injury exhibited significantly enhanced ROS production  
454 when compared to the response of neutrophils from HCs (Figure 2A). No difference in basal  
455 ROS generation was seen between HCs and patient neutrophils acquired  $\leq$ 1-hour or 4-12  
456 hours post-injury (Figure 2A). In response to PMA stimulation, there was a significant  
457 reduction in ROS production, relative to HCs, for neutrophils isolated from patients only at  
458 the 48-72 hour post-injury time point (Figure 2B).

459

### 460 **Traumatic injury results in an immediate and sustained elevation in the frequency and 461 absolute number of circulating IGs**

462

463 Compared to their mature counterparts, immature neutrophils exhibit impaired *ex vivo* NET  
464 production and reduced ROS production upon stimulation with inflammatory agonists  
465 (*Hampson et al. 2017; Martinelli et al. 2004*). Relative to the values recorded for HCs,  
466 trauma patients presented, at all sampling time points, with a significantly elevated frequency  
467 (Figure 2C) and absolute number (Figure 2D) of circulating IGs.

468

### 469 **Traumatic injury is associated with impaired MAPK signalling**

470

471 MAPK signalling is a prerequisite for PMA-induced NET production (*Keshari et al. 2013*).  
472 Due to the significant lymphocytosis that occurs within minutes of traumatic injury  
473 (*Hazeldine et al. 2017*), and the small blood volume collected from patients at the scene of  
474 injury, we were unable to isolate a sufficient number of neutrophils from pre-hospital blood  
475 samples to examine MAPK signalling. However, we found neutrophils isolated from patients

476 4-12 and 48-72 hours post-injury exhibited significantly increased phosphorylation of P38  
477 MAPK (Figure 3A) but not ERK1/2 (Figure 4A) in the absence of exogenous stimulation.

478

479 In response to treatment with PMA, neutrophils obtained from HCs exhibited a significant  
480 increase in P38 phosphorylation (Figure 3B-D). In contrast, no significant PMA-induced  
481 increase in P38 phosphorylation was observed for neutrophils isolated from trauma patients  
482 4-12 or 48-72 hours post-injury (Figures 3B-D). Compared to untreated cells, neutrophils  
483 isolated from HCs and trauma patients at the 4-12 and 48-72 hour post-injury time points  
484 displayed a significant increase in ERK1/2 phosphorylation following 5, 10 and 15 minutes  
485 of PMA stimulation (Figure 4B-D). However, across these three stimulation time points, the  
486 degree of ERK1/2 phosphorylation was significantly greater in neutrophils isolated from HCs  
487 (Figure 4B-D)

488

### 489 **Altered neutrophil glucose uptake and metabolism post-trauma**

490

491 Confirming the results of a recent study that demonstrated a necessity for exogenous glucose  
492 in PMA-induced NET production (*Rodriguez-Espinosa et al. 2015*), we found neutrophils  
493 cultured in glucose free media released significantly less DNA upon PMA stimulation than  
494 neutrophils stimulated in glucose containing media (Supplementary Figure 2). Based on our  
495 observation of a trauma-induced impairment in *ex vivo* NET generation following PMA  
496 treatment, we investigated the effect of injury on neutrophil glucose uptake. Using the  
497 fluorescent glucose analogue 2-NBDG, enhanced basal glucose uptake was recorded for  
498 neutrophils isolated from trauma patients within 1-hour of injury (Figure 5A), but in response  
499 to PMA stimulation, a significant trauma-induced impairment in neutrophil glucose uptake  
500 was seen at all sampling time points (Figure 5B).

501

502 We next examined whether injury impacted upon glucose metabolism, a non-redundant step  
503 in NET formation triggered by PMA stimulation (*Rodriguez-Espinosa et al. 2015*). Using  
504 lactate production as a marker of neutrophil glycolytic activity, we measured lactate  
505 concentrations in supernatants collected from resting and PMA-stimulated neutrophils  
506 following a 3-hour *in vitro* culture. Compared to HCs, neutrophils isolated from trauma  
507 patients at the 48-72 hour post-injury time-point exhibited enhanced basal (Figure 5C) but  
508 impaired PMA-induced lactate production (Figure 5D). The increase in basal lactate  
509 generation was accompanied by a significant up-regulation in the expression of the glycolytic  
510 enzymes pyruvate kinase and lactate dehydrogenase A (Figure 5E-F).

511

### 512 **Neutrophils pre-treated with mtDAMPs exhibit impaired NET production but** 513 **enhanced ROS generation upon secondary stimulation**

514

515 Compared to the levels measured in samples from HCs, serum concentrations of the  
516 mitochondrial-derived protein ND6 were significantly increased in patients at all post-injury  
517 time points, confirming the release of mtDAMPs after trauma (Figure 6A). Demonstrating the  
518 immune stimulatory properties of mtDAMPs, we measured significantly reduced CD62L,  
519 CXCR1 and CXCR2 expression as well as increased CD11b density on the surface of  
520 mtDAMP treated neutrophils (Supplementary Table 1). These changes in neutrophil surface  
521 phenotype were accompanied by activation of ERK 1/2 MAPK signalling (Supplementary  
522 Figure 3A). The emerging concept of mtDAMP-induced tolerance of neutrophil function is  
523 based in part on experimental data that has shown prior activation of neutrophils with  
524 bacterial-derived N-formylated peptides results in impaired migration upon secondary  
525 stimulation (*Li et al. 2015*). Confirming these findings, we found that neutrophils pre-treated

526 with 40 or 100 µg/ml preparations of whole mtDAMPs exhibited significantly reduced  
527 transmigration towards the chemokine LTB<sub>4</sub> (Supplementary Figure 3B). In contrast, no  
528 impairment in migration was witnessed for neutrophils pre-treated with 100 µg/ml of purified  
529 mtDNA (Supplementary Figure 3C).

530

531 To determine whether prior mtDAMP treatment influenced PMA-induced NET production,  
532 fluorometric analysis was performed on cell-free supernatants collected from cultures of  
533 PMA stimulated neutrophils that had been pre-treated with mtDAMPs or vehicle control.  
534 Analysis revealed neutrophils pre-exposed to 40 or 100 µg/ml mtDAMPs released  
535 significantly less DNA following a 3-hour stimulation with PMA than vehicle-treated  
536 controls (Figure 6B). Fluorescence microscopy confirmed this mtDAMP-induced inhibition  
537 of PMA-induced NET generation (Figure 6C). Interestingly, focussing upon neutrophils pre-  
538 treated with 100 µg/ml mtDAMPs, images revealed that despite a significant reduction in  
539 NET formation, the cells had lost their multi-lobed nuclear morphology following stimulation  
540 with PMA, presenting with decondensed nuclear material (Figure 6C). In contrast to whole  
541 mtDAMP preparations, neutrophils pre-treated with 40 µg/ml of purified mtDNA prior to  
542 PMA stimulation showed no impairment in NET production (data not shown). Interestingly,  
543 ROS production, which is a prerequisite for NET formation, was significantly higher upon  
544 secondary PMA stimulation for neutrophils pre-treated with 100 µg/ml mtDAMPs  
545 (Figure 6D).

546

#### 547 **AMPK is activated by mtDAMP treatment and inhibition of AMPK partially** 548 **ameliorates the mtDAMP-induced reduction in NET formation**

549

550 In stimulated T cells, elevated intracellular calcium levels activate AMPK, a recently  
551 described negative regulator of PMA-induced NET formation (*Tamas et al. 2006; Wang et*  
552 *al. 2015*). As raised intracellular calcium levels are a feature of mtDAMP treated neutrophils  
553 (*Zhang et al. 2010*), we determined the activation status of AMPK in neutrophils following  
554 mtDAMP stimulation. To do this, cell lysates, prepared from neutrophils stimulated for 2, 5,  
555 10 and 15 minutes with 100 µg/ml mtDAMPs, were probed with a phospho-specific antibody  
556 directed against Thr172, a residue within the activation loop of AMPK. As shown in Figure  
557 7A, mtDAMP treatment resulted in an immediate and persistent phosphorylation of residue  
558 Thr172. Treating neutrophils with the FPR-1 antagonist CsH prior to mtDAMP stimulation  
559 resulted in a significant reduction in AMPK phosphorylation, suggesting that N-formyl  
560 peptides drive mtDAMP-induced activation of AMPK (Figure 7B). In antigen challenged T  
561 cells, phosphorylation of AMPK requires the activation of calcium-calmodulin-dependent  
562 protein kinase kinases (CaMKKs), a class of serine/threonine protein kinases activated by  
563 increases in intracellular calcium (*Tamas et al. 2006*). To investigate whether CaMKKs were  
564 involved in mtDAMP-induced phosphorylation of AMPK in neutrophils, we treated  
565 neutrophils with the CaMKK selective inhibitor STO-609 prior to mtDAMP stimulation.  
566 Compared to vehicle control, a significant impairment in mtDAMP-induced activation of  
567 AMPK was detected in neutrophils pre-treated with STO-609 (Figure 7C).

568

569 AMPK has recently been shown to be a negative regulator of PMA-induced NET formation  
570 (*Jiang et al. 2014; Wang et al. 2015*). To investigate whether AMPK signalling was involved  
571 in mtDAMP-mediated suppression of NET formation, we treated neutrophils with compound  
572 C, an inhibitor of AMPK, prior to mtDAMP exposure. Compared to vehicle control,  
573 significantly greater NET production in response to PMA stimulation was recorded for  
574 neutrophils pre-treated with compound C (Figure 7D).

575 **MtDAMP pre-treatment results in impaired lactate generation by neutrophils upon**  
576 **secondary stimulation with PMA**

577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624

Confirming results of previous studies that had shown AMPK to be a negative regulator of aerobic glycolysis (*Faubert et al. 2013*), we measured significantly lower concentrations of lactate in supernatants collected from PMA stimulated neutrophils that had been pre-treated with the AMP mimetic AICAR when compared to vehicle control (Figure 8A). Given that aerobic glycolysis is a key metabolic event in PMA-induced NET formation (*Rodriguez-Espinosa et al. 2015*) and our observation of reduced NET generation following PMA stimulation for neutrophils pre-treated with mtDAMPs (Figure 6B-C), we investigated whether AMPK activation triggered by mtDAMP exposure was associated with an impairment in neutrophil glycolysis. Following 1, 2 or 3 hour stimulation with PMA, significantly lower lactate concentrations were measured in supernatants collected from neutrophils pre-treated with 100 µg/ml mtDAMPs (Figure 8B).

## 625 Discussion

626

627 Here, via the analysis of blood samples acquired from trauma patients within 1-hour of  
628 injury, we have shown for the first time that major trauma results in an immediate impairment  
629 in neutrophil anti-microbial defences, specifically the formation of NETs. This defect  
630 persisted into the acute post-injury phase (4-72 hours) and was accompanied by impaired  
631 ROS production, MAPK activation and a reduction in both glucose uptake and breakdown.  
632 Furthermore, we showed that the trauma-induced reduction in NET formation could be  
633 replicated *in vitro* by treating neutrophils isolated from healthy donors with mtDAMPs prior  
634 to secondary stimulation. Thus, our data provides support for the emerging concept of  
635 mtDAMP-induced tolerance, where the post-injury release of mtDAMPs into the circulation  
636 has been proposed to contribute to the neutrophil dysfunction that develops in the aftermath  
637 of traumatic injury (Kaczmarek *et al.* 2018; Li *et al.* 2015).

638

639 In the absence of secondary stimulation, neutrophils obtained from patients within minutes,  
640 but not hours, of injury released significantly more DNA into culture supernatants than  
641 neutrophils isolated from healthy controls. This immediate trauma-induced increase in basal  
642 NET generation is in agreement with the findings of our previous trauma-based study, where  
643 in pre-hospital plasma samples, we reported the presence of citrullinated histone H3, a  
644 protein that decorates the nuclear DNA backbone of NETs (Hazeldine *et al.* 2017; Liu *et al.*  
645 2012). Thus, taken together, our results suggest that neutrophils are immediately exposed to  
646 NET-inducing stimuli post-injury. Supporting this proposal, data presented here and in our  
647 previous manuscript have shown serum concentrations of known NET inducers, which  
648 include TNF- $\alpha$ , IL-8, IL-33 and HMGB-1 are all significantly elevated within minutes of  
649 injury (Hazeldine *et al.* 2017; Keshari *et al.* 2012; Liu *et al.* 2012; Tadie *et al.* 2013; Yazdani  
650 *et al.* 2017). Of these agonists, the immediate release of HMGB-1 may be particularly  
651 pertinent given that within 60 minutes of *in vitro* co-culture, neutrophils stimulated with this  
652 nuclear-derived DAMP have been shown to generate NETs (Huang *et al.* 2015). Moreover,  
653 as HMGB-1 induced NET formation is independent of ROS generation by NADPH oxidase  
654 (Tadie *et al.* 2013), immediate exposure to this DAMP could explain our observation of  
655 enhanced *ex vivo* NET formation by neutrophils isolated from pre-hospital blood samples that  
656 exhibited no difference in basal ROS production when compared to neutrophils from HCs. In  
657 terms of stimulus-induced NET formation, this is the first study to show quantitatively that  
658 trauma results in reduced NET generation to PMA stimulation. We confirmed this  
659 impairment, which was evident at all three sampling time points, by fluorescent microscopy,  
660 with our images akin to those presented in a previous study that reported a qualitative post-  
661 injury reduction in PMA-induced NET production in a much smaller cohort of trauma  
662 patients (Li *et al.* 2015).

663

664 The processes that mediate PMA-induced NET generation are well defined, with prominent  
665 roles assigned to ROS generation, MAPK activation, glucose uptake and glycolysis (Fuchs *et al.*  
666 2007; Hakkim *et al.* 2011; Keshari *et al.* 2013; Rodriguez-Espinosa *et al.* 2015). Across  
667 our sampling time points, we observed defects in each of these processes, suggesting that  
668 multiple impairments rather than a single aberration underlie the post-injury reduction in  
669 NET formation. Whilst our data demonstrating a trauma-induced impairment in neutrophil  
670 ROS production has been described previously (Hampson *et al.* 2017; Hazeldine *et al.* 2017),  
671 we are the first to report a post-trauma reduction in MAPK activation, glucose uptake and  
672 metabolism by neutrophils in response to *ex vivo* stimulation. Aside from NET production,  
673 other anti-microbial mechanisms of neutrophils utilise glucose. For instance, chemotaxis  
674 requires the uptake of exogenous glucose (Weisdorf *et al.* 1982), whilst breakdown of

675 endogenous glucose is important for stimulus-induced ROS production and phagocytosis  
676 (*Rodriguez-Espinosa et al. 2015; Weisdorf et al. 1982*). Interestingly, as reported here for  
677 NET generation, these three defence strategies have all been shown in *ex vivo* assays to be  
678 significantly impaired following trauma (*Hampson et al. 2017; Li et al. 2015; Liao et al.*  
679 *2013; Tarlowe et al. 2003*). Thus, the injury-induced reduction we have demonstrated in  
680 glucose uptake and metabolism may be a mechanism underlying many facets of post-trauma  
681 neutrophil dysfunction. Aside from the changes we found in neutrophil signalling and  
682 metabolism, trauma patients presented at all three time points with elevated circulating levels  
683 of IGs. Shown *in vitro* to exhibit impaired NET production upon secondary stimulation  
684 (*Martinelli et al. 2004*), the immediate and persistent presence of immature cells offers  
685 another potential mechanistic explanation for the trauma-induced reduction in NET  
686 formation.

687  
688 Suggesting that traumatic injury modulates cell metabolism, we demonstrated a post-trauma  
689 elevation in lactate production by resting neutrophils, an observation that is in agreement with  
690 the findings of a previous study. In a cohort of polytrauma patients, Oehler et al reported a  
691 higher glycolytic activity, relative to HCs, for neutrophils isolated from subjects between 48  
692 and 120 hours post-injury, an enhancement they attributed to increased expression and  
693 activity of pyruvate kinase, a glycolytic enzyme that catalyses the conversion of  
694 phosphoenolpyruvate to pyruvate (*Oehler et al. 2000*). Here, we confirmed that traumatic  
695 injury induces increased expression of pyruvate kinase and showed that this is accompanied  
696 by increased expression of lactate dehydrogenase A. Whilst up-regulation of lactate  
697 dehydrogenase A, which converts pyruvate to lactate, has been described in transcriptomic  
698 analysis of whole blood leukocytes isolated from critically-ill patients (*Nalos et al. 2016*),  
699 this is the first study to demonstrate increased protein expression of lactate dehydrogenase in  
700 neutrophils post-injury.

701  
702 Accompanying the impairment we observed in *ex vivo* NET formation was a significant  
703 trauma-induced elevation in the circulating levels of the mitochondrial-derived N-formylated  
704 peptide ND6. In a recent study, suppressed chemotactic responses towards CXCL1 and  
705 LTB<sub>4</sub> were reported for neutrophils pre-treated with synthetic ND6 (*Kaczmarek et al. 2018*),  
706 a finding that mirrored results of previous studies where prior exposure to bacterial-derived  
707 N-formylated peptides or ND6 respectively was shown to reduce neutrophil migration and  
708 ROS production upon secondary stimulation (*Gabl et al. 2018; Li et al. 2015*). Adding to this  
709 growing body of literature that suggests a tolerising effect for mitochondrial-derived peptides  
710 on neutrophil function, we demonstrated that neutrophils pre-treated with whole mtDAMP  
711 preparations, but not purified mtDNA, exhibited significantly reduced NET production  
712 following PMA stimulation.

713  
714 A striking observation of our NET based assays was that despite an absence of NET  
715 production, mtDAMP pre-treated neutrophils lost their distinctive multi-lobed nuclear  
716 morphology upon PMA challenge. Interestingly, chromatin decondensation in the absence of  
717 DNA release was recently reported for PMA stimulated neutrophils pre-treated with the  
718 glycolysis inhibitor 2-deoxy-glucose (*Rodriguez-Espinosa et al. 2015*). Using lactate as a  
719 readout of glycolytic activity, we measured significantly reduced lactate concentrations in the  
720 supernatants of mtDAMP pre-treated neutrophils stimulated with PMA, demonstrating that  
721 exposure to mtDAMPs influences the metabolism of immune cells.

722  
723 We found that exposing neutrophils to mtDAMPs resulted in activation of the  
724 serine/threonine protein kinase AMPK. In T cells, increases in intracellular calcium levels



725 promote AMPK phosphorylation through activation of CaMKKs (*Tamas et al. 2006*). Our  
726 data demonstrating a significant reduction in mtDAMP-induced phosphorylation of AMPK in  
727 neutrophils pre-treated with the selective CAMMK inhibitor STO-506 indicates this  
728 signalling pathway is also activated in stimulated neutrophils. As the only component of  
729 mtDAMPs that promotes calcium mobilisation in neutrophils (*Zhang et al. 2010*), signals  
730 derived from N-formyl peptides are likely to have driven the mtDAMP-induced  
731 phosphorylation of AMPK. Supporting this idea, we observed significantly reduced  
732 mtDAMP-induced AMPK phosphorylation in neutrophils pre-treated with CsH, an FPR-1  
733 antagonist that prevents calcium mobilisation upon mtDAMP stimulation (*Zhang et al. 2010*).

734

735 Demonstrating that AMPK is a negative regulator of NET formation, significantly enhanced  
736 and impaired NET formation has been reported for PMA stimulated neutrophils pre-treated  
737 with AMPK inhibitors and activators respectively (*Jiang et al. 2014; Wang et al. 2015*).  
738 Supporting these data, we showed significantly greater PMA-induced NET production by  
739 neutrophils treated with the AMPK inhibitor compound C prior to mtDAMP exposure. How  
740 activation of AMPK inhibits NET formation is currently unknown. Based on published  
741 literature, we propose two mechanisms, both of which revolve around the ability of AMPK to  
742 inhibit the serine/threonine protein kinase mammalian target of rapamycin (mTOR) (*Faubert  
743 et al. 2013; MacIver et al. 2011*). Firstly, inhibition of mTOR has been shown to significantly  
744 reduce Glut1 transporter activity (*Wieman et al. 2007*). Given the importance of extracellular-  
745 derived glucose in NET formation (*Rodriguez-Espinosa et al. 2015*), reduced glucose uptake,  
746 secondary to impaired mTOR and Glut1 activity, could contribute to the reduction in NET  
747 production and lactate generation that we observed for neutrophils pre-treated with  
748 mtDAMPs. Furthermore, this mechanism could contribute to the mtDAMP-induced  
749 impairment we reported in neutrophil transmigration since chemotaxis also utilises  
750 extracellular sources of glucose (*Weisdorf et al. 1982*). Alongside Glut1, mTOR signalling  
751 has been implicated in regulating the expression/activity of hypoxia-inducible factor-1 $\alpha$   
752 (HIF-1 $\alpha$ ) (*Faubert et al. 2013*). A transcription factor involved in promoting aerobic  
753 glycolysis, activation of HIF-1 $\alpha$  precedes both NET formation (*McInturff et al. 2012*) and  
754 myeloid cell migration (*Cramer et al. 2003*). Thus, reduced activity/expression of HIF-1 $\alpha$ ,  
755 secondary to AMPK-mediated inhibition of mTOR, could be an additional/alternative  
756 explanation for the mtDAMP-induced reduction in lactate generation, NET generation and  
757 neutrophil chemotaxis. Importantly, both these proposed mechanisms could occur in  
758 neutrophils without affecting their capacity for ROS production, which we found was  
759 significantly increased following mtDAMP treatment. Indeed, as the energy required for ROS  
760 generation is derived from endogenous sources of glucose (*Rodriguez-Espinosa et al. 2015*),  
761 this anti-microbial function could occur in the background of an AMPK driven reduction in  
762 extracellular glucose uptake. Our finding of increased ROS production to PMA stimulation  
763 for neutrophils pre-treated with mtDAMPs contradicts the post-injury impairment we  
764 reported in ROS generation for trauma patients that presented with significantly elevated  
765 serum concentrations of mtDAMPs. We believe this discrepancy may be attributable to  
766 trauma-induced changes in the composition of the circulating neutrophil pool. For example,  
767 traumatic injury results in the emergence into circulation of IGs and CD16<sup>BRIGHT</sup> CD62L<sup>DIM</sup>  
768 neutrophils, both of which exhibit impaired stimulus-induced ROS generation (*Drifte et al.  
769 2013; Sauce et al. 2017*). In contrast, healthy subjects, who served as the cohort for our  
770 mtDAMP pre-treatment experiments, possess a homogenous pool of fully-functional mature  
771 neutrophils that would exhibit a greater capacity to respond to stimulation.

772

773 Although we have shown that neutrophils pre-treated with compound C exhibit increased  
774 NET generation upon PMA stimulation, inhibition of AMPK only partially ameliorated the

775 reduction in NET formation that occurred with mtDAMP treatment. Other factors aside from  
776 AMPK activation must therefore be involved in mediating the mtDAMP-induced suppression  
777 of NET production. Recently, through a proposed mechanism of action that involved the  
778 prevention of membrane rupture, lactoferrin, an iron-binding glycoprotein stored within the  
779 secondary granules of neutrophils, was found to suppress NET release triggered by PMA  
780 stimulation (*Okubo et al. 2016*). Indicative of a defect in the latter stages of NET production,  
781 we showed chromatin decondensation in the absence of DNA release was a feature of  
782 mtDAMP-treated neutrophils. Thus, based on this observation and the fact that exposure to  
783 mtDAMPs promotes neutrophil degranulation (*Hauser et al. 2010; Hazeldine et al. 2015;*  
784 *Zhang et al. 2010*), we suggest that a mtDAMP-induced release of lactoferrin could represent  
785 an additional mechanistic explanation for the impairment in NET formation that occurs  
786 following mtDAMP exposure.

787

788 This study has some limitations. Conducted at a single major trauma centre, the results of our  
789 prospective observational study are based on the analysis of a small number of patient  
790 samples, meaning our findings require validation in larger independent cohorts. This point is  
791 particularly pertinent to our analysis of pre-hospital blood samples, where inter-individual  
792 variability in immune cell number and volume of blood collected at the scene of injury meant  
793 we were unable to perform all assays on each patient at this time point. As a heterogeneous  
794 collection of proteins, lipids and DNA, no study to date has quantified the exact concentration  
795 of mtDAMPs released into the circulation post-injury. Thus, our *in vitro* treatment of  
796 neutrophils with 40 or 100 µg/ml mtDAMPs may not be physiologically relevant. However,  
797 these doses match those used in previous *in vitro* based studies that have examined the effect  
798 of mtDAMP exposure on neutrophil anti-microbial function (*Hauser et al. 2010; Hazeldine*  
799 *et al. 2015; Zhang et al. 2010*). Similarly, our decision to use the phorbol ester PMA as an  
800 agonist may be considered a study limitation. However, as a potent stimulus, it allowed us to  
801 study maximal neutrophil responses. Moreover, as the agonist of choice for all previous  
802 trauma-based studies that had examined stimulus-induced NET production post-injury  
803 (*Hampson et al. 2017; Itagaki et al. 2015; Li et al. 2015*), our use of PMA enabled us to  
804 compare our observations to those in the published literature.

805

806 In summary, this is the first study to describe a quantitative post-trauma reduction in NET  
807 formation as an immediate on-scene phenomenon, which is accompanied by aberrant  
808 intracellular signalling and cell metabolism. In addition, we have shown that the post-injury  
809 reduction in NET generation can be recreated *in vitro* by treating neutrophils isolated from  
810 healthy subjects with mtDAMPs prior to PMA stimulation. Thus, our data support the  
811 suggestion that the release of mtDAMPs from damaged tissue is a contributory factor in the  
812 reduction in neutrophil function that occurs post-injury (*Hampson et al. 2017; Hazeldine et*  
813 *al. 2017; Li et al. 2015; Liao et al. 2013; Tarlowe et al. 2003*).

814

815

816

817

818

819

820

821

822

823

824

825 **Acknowledgements**

826

827 The research was funded by the NIHR Surgical Reconstruction and Microbiology Research  
828 Centre (SRMRC). The authors would also like to acknowledge the Queen Elizabeth Hospital  
829 Birmingham Charity for funding the purchase of the Sysmex XN-1000 haematology analyser.  
830 The views expressed are those of the author(s) and not necessarily those of the NHS, the  
831 NIHR or the Department of Health. The authors thank all the research, nursing and  
832 administrative staff at the NIHR-SRMRC for their assistance in data collection and  
833 management. We also thank the emergency healthcare professionals in the West Midlands  
834 Ambulance Service NHS Foundation Trust, the Midlands Air Ambulance Charity and the  
835 University Hospitals Birmingham NHS Foundation Trust for their participation in the  
836 identification, enrolment and follow up of patients in this study.

837

838 **Author contributions statement**

839

840 JH designed the study, performed experimental work, analysed data and wrote the  
841 manuscript. RD performed experimental work and analysed data. PH critically appraised the  
842 manuscript and JL conceptualised the study and contributed to writing the manuscript.

843

844 **Conflict of interest statement**

845

846 The authors declare the submitted manuscript was not carried out in the presence of any  
847 personal, professional or financial relationships that could potentially be construed as a  
848 conflict of interest.

849

850

851

852

853

854

855

856

857

858

859

860

861

862

863

864

865

866

867

868

869

870

871

872

873

874

875 **References**

- 876 Brinkmann, V., and Zychlinsky, A. (2012). Neutrophil extracellular traps: is immunity the  
877 second function of chromatin? *J.Cell Biol.* 198, 773-783. DOI:10.1083/jcb.201203170.
- 878 Cohen, M.J., Brohi, K., Calfee, C.S., Rahn, P., Chesebro, B.B., Christiaans, S.C. et al. (2009).  
879 Early release of high mobility group box nuclear protein 1 after severe trauma in humans:  
880 role of injury severity and tissue hypoperfusion. *Crit Care* 13, R174. DOI:10.1186/cc8152.
- 881 Cramer, T., Yamanishi, Y., Clausen, B.E., Forster, I., Pawlinski, R., Mackman, N. et al.  
882 (2003). HIF-1alpha is essential for myeloid cell-mediated inflammation. *Cell* 112, 645-657.
- 883 Drifte, G., Dunn-Siegrist, I., Tissieres, P., and Pugin, J. (2013) Innate immune functions of  
884 immature neutrophils in patients with severe sepsis and severe systemic inflammatory  
885 response syndrome. *Crit. Care Med.* 41, 820-832. DOI: 10.1097/CCM.0b013e318274647d.
- 886 Faubert, B., Boily, G., Izreig, S., Griss, T., Samborska, B., Dong, Z. et al. (2013). AMPK is a  
887 negative regulator of the Warburg effect and suppresses tumor growth in vivo. *Cell Metab* 17,  
888 113-124. DOI:10.1016/j.cmet.2012.12.001.
- 889 Fernandez-Ruiz, I., Arnalich, F., Cubillos-Zapata, C., Hernandez-Jimenez, E., Moreno-  
890 Gonzalez, R., Toledano, V. et al. (2014). Mitochondrial DAMPs induce endotoxin tolerance  
891 in human monocytes: an observation in patients with myocardial infarction. *PLoS.One.* 9,  
892 e95073. DOI: 10.1371/journal.pone.0095073.
- 893 Fuchs, T.A., Abed, U., Goosmann, C., Hurwitz, R., Schulze, I., Wahn, V. et al. (2007). Novel  
894 cell death program leads to neutrophil extracellular traps. *J.Cell Biol.* 176, 231-241. DOI:  
895 10.1083/jcb.200606027.
- 896 Gabl, M., Sundqvist, M., Holdfeldt, A., Lind, S., Martensson, J., Christenson, K. et al (2018).  
897 Mitocryptides from Human Mitochondrial DNA-Encoded Proteins Activate Neutrophil  
898 Formyl Peptide Receptors: Receptor Preference and Signaling Properties. *J.Immunol.* 200,  
899 3269-3282. DOI:10.4049/jimmunol.1701719.
- 900 Hakkim, A., Fuchs, T.A., Martinez, N.E., Hess, S., Prinz, H., Zychlinsky, A. et al (2011).  
901 Activation of the Raf-MEK-ERK pathway is required for neutrophil extracellular trap  
902 formation. *Nat.Chem.Biol.* 7, 75-77. DOI:10.1038/nchembio.496.
- 903 Hampson, P., Dinsdale, R.J., Wearn, C.M., Bamford, A.L., Bishop, J.R.B., Hazeldine, J. et al  
904 (2017). Neutrophil Dysfunction, Immature Granulocytes, and Cell-free DNA are Early  
905 Biomarkers of Sepsis in Burn-injured Patients: A Prospective Observational Cohort Study.  
906 *Ann.Surg.* 265, 1241-1249. DOI: 10.1097/SLA.0000000000001807.
- 907 Hauser, C.J., Sursal, T., Rodriguez, E.K., Appleton, P.T., Zhang, Q. and Itagaki, K. (2010).  
908 Mitochondrial damage associated molecular patterns from femoral reamings activate  
909 neutrophils through formyl peptide receptors and P44/42 MAP kinase. *J.Orthop.Trauma* 24,  
910 534-538. DOI: 10.1097/BOT.0b013e3181ec4991.
- 911 Hazeldine, J., Hampson, P. and Lord, J.M. (2014). The impact of trauma on neutrophil  
912 function. *Injury* 45, 1824-1833. DOI: 10.1016/j.injury.2014.06.021.
- 913 Hazeldine, J., Hampson, P., Opoku, F.A., Foster, M. and Lord, J.M. (2015). N-Formyl  
914 peptides drive mitochondrial damage associated molecular pattern induced neutrophil

915 activation through ERK1/2 and P38 MAP kinase signalling pathways. *Injury* 46, 975-984.  
916 doi: 10.1016/j.injury.2015.03.028.

917 Hazeldine, J., Naumann, D.N., Toman, E., Davies, D., Bishop, J.R.B., Su,Z. et al (2017).  
918 Prehospital immune responses and development of multiple organ dysfunction syndrome  
919 following traumatic injury: A prospective cohort study. *PLoS.Med.* 14 (7), e1002338. DOI:  
920 10.1371/journal.pmed.1002338.

921 Huang, H., Tohme, S., Al-Khafaji, A.B., Tai, S., Loughran, P., Chen, L. et al (2015).  
922 Damage-associated molecular pattern-activated neutrophil extracellular trap exacerbates  
923 sterile inflammatory liver injury. *Hepatology* 62, 600-614. DOI: 10.1002/hep.27841.

924 Itagaki, K., Kaczmarek, E., Lee, Y.T., Tang, I.T., Isal, B., Adibnia, Y. et al (2015).  
925 Mitochondrial DNA released by trauma induces neutrophil extracellular traps. *PLoS.One.* 10  
926 (3), e0120549. DOI: 10.1371/journal.pone.0120549.

927 Jiang, S., Park, DW., Tadie, JM., Gregoire, M., Deshane, J., Pittet, JF. et al (2014). Human  
928 resistin promotes neutrophil proinflammatory activation and neutrophil extracellular trap  
929 formation and increases severity of acute lung injury. *J. Immunol.* 192(10):4795-4803. DOI:  
930 10.4049/jimmunol.1302764.

931 Johansson, J., Sjogren, F., Bodelsson, M. and Sjoberg, F. (2011). Dynamics of leukocyte  
932 receptors after severe burns: an exploratory study. *Burns* 37, 227-233. DOI:  
933 10.1016/j.burns.2010.08.015.

934 Kaczmarek, E., Hauser, C.J., Kwon, W.Y., Rica, I., Chen, L., Sandler, N. et al. (2018). A  
935 subset of five human mitochondrial formyl peptides mimics bacterial peptides and  
936 functionally deactivates human neutrophils. *J.Trauma Acute.Care Surg.* 85(5): 936-943. DOI:  
937 10.1097/TA.0000000000001971.

938 Keshari, R.S., Jyoti, A., Dubey, M., Kothari, N., Kohli, M., Bogra, J. et al (2012). Cytokines  
939 induced neutrophil extracellular traps formation: implication for the inflammatory disease  
940 condition. *PLoS.One.* 7, e48111. DOI: 10.1371/journal.pone.0048111.

941 Keshari, R.S., Verma, A., Barthwal, M.K. and Dikshit, M. (2013). Reactive oxygen species-  
942 induced activation of ERK and p38 MAPK mediates PMA-induced NETs release from  
943 human neutrophils. *J.Cell Biochem.* 114, 532-540. DOI: 10.1002/jcb.24391.

944 Kirchner, T., Moller, S., Klinger, M., Solbach, W., Laskay, T. and Behnen,M. (2012). The  
945 impact of various reactive oxygen species on the formation of neutrophil extracellular traps.  
946 *Mediators.Inflamm.* 2012, 849136. DOI: 10.1155/2012/849136.

947 Lelifeld, P.H., Wessels, C.M., Leenen, L.P., Koenderman, L. and Pillay,J. (2016). The role  
948 of neutrophils in immune dysfunction during severe inflammation. *Crit Care* 20, 73. DOI:  
949 10.1186/s13054-016-1250-4.

950 Li, H., Itagaki, K., Sandler, N., Gallo, D., Galenkamp, A., Kaczmarek,E. et al (2015).  
951 Mitochondrial damage-associated molecular patterns from fractures suppress pulmonary  
952 immune responses via formyl peptide receptors 1 and 2. *J.Trauma Acute.Care Surg.* 78, 272-  
953 279. DOI: 10.1097/TA.0000000000000509.

954 Liao, Y., Liu, P., Guo, F., Zhang, Z.Y. and Zhang, Z. (2013). Oxidative burst of circulating  
955 neutrophils following traumatic brain injury in human. *PLoS.One.* 8, e68963. DOI:  
956 10.1371/journal.pone.0068963.

957 Liu, C.L., Tangsombatvisit, S., Rosenberg, J.M., Mandelbaum, G., Gillespie, E.C.,  
958 Gozani, O.P. et al (2012). Specific post-translational histone modifications of neutrophil  
959 extracellular traps as immunogens and potential targets of lupus autoantibodies. *Arthritis*  
960 *Res. Ther.* 14, R25. DOI: 10.1186/ar3707.

961 MacIver, N.J., Blagih, J., Saucillo, D.C., Tonelli, L., Griss, T., Rathmell, J.C. et al (2011).  
962 The liver kinase B1 is a central regulator of T cell development, activation, and metabolism.  
963 *J.Immunol.* 187, 4187-4198. DOI: 10.4049/jimmunol.1100367.

964 Martinelli, S., Urosevic, M., Daryadel, A., Oberholzer, P.A., Baumann, C., Fey, M.F. et al  
965 (2004). Induction of genes mediating interferon-dependent extracellular trap formation during  
966 neutrophil differentiation. *J.Biol.Chem.* 279, 44123-44132. DOI:10.1074/jbc.M405883200.

967 McInturff, A.M., Cody, M.J., Elliott, E.A., Glenn, J.W., Rowley, J.W., Rondina, M.T. et al  
968 (2012). Mammalian target of rapamycin regulates neutrophil extracellular trap formation via  
969 induction of hypoxia-inducible factor 1 alpha. *Blood* 120, 3118-3125. DOI: 10.1182/blood-  
970 2012-01-405993.

971 Nalos, M., Parnell, G., Robergs, R., Booth, D., McLean, A.S. and Tang, B.M. (2016).  
972 Transcriptional reprogramming of metabolic pathways in critically ill patients. *Intensive Care*  
973 *Med.Exp.* 4, 21. DOI: 10.1186/s40635-016-0094-1.

974 Oehler, R., Weingartmann, G., Manhart, N., Salzer, U., Meissner, M., Schlegel, W. et al  
975 (2000). Polytrauma induces increased expression of pyruvate kinase in neutrophils. *Blood* 95,  
976 1086-1092.

977 Okubo, K., Kamiya, M., Urano, Y., Nishi, H., Herter, J.M. and Mayadas, T. et al (2016).  
978 Lactoferrin suppresses neutrophil extracellular traps release in inflammation. *EBioMedicine.*  
979 10, 204-215. DOI: 10.1016/j.ebiom.2016.07.012.

980 Pillay, J., Kamp, V.M., van, H.E., Visser, T., Tak, T., Lammers, J.W. et al (2012). A subset  
981 of neutrophils in human systemic inflammation inhibits T cell responses through Mac-1.  
982 *J.Clin.Invest* 122, 327-336. DOI: 10.1172/JCI57990.

983 Rodriguez-Espinosa, O., Rojas-Espinosa, O., Moreno-Altamirano, M.M., Lopez-Villegas,  
984 E.O. and Sanchez-Garcia, F.J. (2015). Metabolic requirements for neutrophil extracellular  
985 traps formation. *Immunology* 145, 213-224. DOI: 10.1111/imm.12437.

986 Sauce, D., Dong, Y., Campillo-Gimenez, L., Casulli, S., Bayard, C., Autran, B. et al (2017).  
987 Reduced oxidative burst by primed neutrophils in the elderly individuals is associated with  
988 increased levels of the CD16BRIGHT/CD62LDIM immunosuppressive subset. *J. Gerontol.*  
989 *A. Biol. Sci. Med. Sci.* 72, 163-172. DOI: 10.1093/gerona/glw062.

990 Tadie, J.M., Bae, H.B., Jiang, S., Park, D.W., Bell, C.P., Yang, H. et al (2013). HMGB1  
991 promotes neutrophil extracellular trap formation through interactions with Toll-like receptor  
992 4. *Am.J.Physiol Lung Cell Mol.Physiol* 304, L342-L349. DOI: 10.1152/ajplung.00151.2012.

993 Tamas, P., Hawley, S.A., Clarke, R.G., Mustard, K.J., Green, K., Hardie, D.G. et al. (2006).  
994 Regulation of the energy sensor AMP-activated protein kinase by antigen receptor and Ca<sup>2+</sup>  
995 in T lymphocytes. *J.Exp.Med.* 203, 1665-1670. DOI:10.1084/jem.20052469.

996 Tarlowe, M.H., Kannan, K.B., Itagaki, K., Adams, J.M., Livingston, D.H. and Hauser, C.J.  
997 (2003). Inflammatory chemoreceptor cross-talk suppresses leukotriene B4 receptor 1-  
998 mediated neutrophil calcium mobilization and chemotaxis after trauma. *J.Immunol.* 171,  
999 2066-2073.

1000 Visser, T., Hietbrink, F., Groeneveld, K.M., Koenderman, L. and Leenen, L.P. (2011).  
1001 Isolated blunt chest injury leads to transient activation of circulating neutrophils.  
1002 *Eur.J.Trauma Emerg.Surg.* 37, 177-184. DOI: 10.1007/s00068-010-0041-x.

1003 Visser, T., Pillay, J., Pickkers, P., Leenen, L.P. and Koenderman, L. (2012). Homology in  
1004 systemic neutrophil response induced by human experimental endotoxemia and by trauma.  
1005 *Shock* 37, 145-151. DOI: 10.1097/SHK.0b013e31823f14a4.

1006 Wang, H., Li, T., Chen, S., Gu, Y. and Ye, S. (2015). Neutrophil Extracellular Trap  
1007 Mitochondrial DNA and Its Autoantibody in Systemic Lupus Erythematosus and a Proof-of-  
1008 Concept Trial of Metformin. *Arthritis Rheumatol.* 67, 3190-3200. DOI: 10.1002/art.39296.

1009 Weisdorf, D.J., Craddock, P.R. and Jacob, H.S. (1982). Glycogenolysis versus glucose  
1010 transport in human granulocytes: differential activation in phagocytosis and chemotaxis.  
1011 *Blood* 60, 888-893.

1012 Wieman, H.L., Wofford, J.A. and Rathmell, J.C. (2007). Cytokine stimulation promotes  
1013 glucose uptake via phosphatidylinositol-3 kinase/Akt regulation of Glut1 activity and  
1014 trafficking. *Mol.Biol.Cell* 18, 1437-1446. DOI:10.1091/mbc.e06-07-0593.

1015 Xu, J., Guardado, J., Hoffman, R., Xu, H., Namas, R., Vodovotz, Y. et al (2017). IL33-  
1016 mediated ILC2 activation and neutrophil IL5 production in the lung response after severe  
1017 trauma: A reverse translation study from a human cohort to a mouse trauma model.  
1018 *PLoS.Med.* 14, e1002365. DOI: 10.1371/journal.pmed.1002365.

1019 Yazdani, H.O., Chen, H.W., Tohme, S., Tai, S., van der Windt, D.J., Loughran, P. et al  
1020 (2017). IL-33 exacerbates liver sterile inflammation by amplifying neutrophil extracellular  
1021 trap formation. *J.Hepatol.* DOI: 10.1016/j.jhep.2017.09.010.

1022 Zhang, Q., Raouf, M., Chen, Y., Sumi, Y., Sursal, T., Junger, W. et al (2010). Circulating  
1023 mitochondrial DAMPs cause inflammatory responses to injury. *Nature* 464, 104-107. DOI:  
1024 10.1038/nature08780.

1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035

1036 **Figure legends**

1037

1038 **Figure 1. Effect of traumatic injury on PMA-induced NET formation.** (A) Basal NET  
1039 generation by resting neutrophils isolated from healthy controls (HC) and trauma patients as  
1040 assessed by DNA concentration in cell free supernatants following a 3-hour *in vitro* culture.  
1041 (B) HMGB-1 and (C) IL-33 concentrations in serum samples from HC and trauma patients.  
1042 IL-33 levels were undetectable (N.D) in serum samples from HC. (D-E) Following a 3-hour  
1043 *in vitro* stimulation with PMA, NET production by neutrophils from HC and trauma patients  
1044 was compared by measuring DNA concentration in cell free supernatants (D) and  
1045 fluorescence microscopy (E). For supernatant analysis, number of samples are shown below  
1046 each time-point. For microscope images, HC (n=12), ≤1H (n=6), 4-12H (n=6) and 48-72H  
1047 (n=8). \* p<0.01, \*\* p<0.001, \*\*\* p<0.0001 vs. HC.

1048

1049 **Figure 2. Neutrophil ROS production and immature granulocyte count post-injury.**  
1050 Comparison of basal (A) and PMA-induced (B) ROS production by neutrophils isolated from  
1051 healthy controls (HC) and trauma patients. Data are presented as area under the curve (AUC)  
1052 (A) or fold increase above vehicle treated controls (B). (C-D) Frequency (C) and absolute  
1053 number (D) of immature granulocytes (IG) in peripheral blood samples from healthy controls  
1054 (HC) and trauma patients. Number of samples analysed are shown below each time-point.  
1055 \* p<0.01, \*\* p<0.001, \*\*\* p<0.0001 vs. HC.

1056

1057 **Figure 3. Traumatic injury results in impaired PMA-induced activation of p38 MAPK.**  
1058 Phosphorylation status of P38 in resting and PMA-stimulated neutrophils isolated from  
1059 healthy controls (HC) and trauma patients 4-12 and 48-72 hours post-injury. Data are  
1060 presented as representative Western blots (B) and densitometry analysis of p38  
1061 phosphorylation in resting (A) or PMA-stimulated neutrophils at the 4-12H (C) and 48-72H  
1062 (D) post-injury time points. HC (n=7), 4-12h (n=9) and 48-72h (n=4). For (A) \* p<0.01,  
1063 \*\* p<0.001 vs. HC. For (C) and (D) \*\* p<0.001 vs. Time 0, # p<0.01, ### p<0.0001 vs. HC  
1064 sample at matched time point.

1065

1066 **Figure 4. Traumatic injury results in impaired PMA-induced activation of ERK**  
1067 **MAPK.** Phosphorylation status of ERK 1/2 in resting and PMA-stimulated neutrophils  
1068 isolated from healthy controls (HC) and trauma patients 4-12 and 48-72 hours post-injury.  
1069 Data are presented as representative Western blots (B) and densitometry analysis of ERK  
1070 phosphorylation in resting (A) or PMA-stimulated neutrophils at the 4-12h (C) and 48-72h  
1071 (D) post-injury time points. HC (n=4), 4-12h (n=8) and 48-72h (n=6). For (C) and (D)  
1072 \* p<0.01, \*\* p<0.001, \*\*\* p<0.0001 vs. Time 0, ## p<0.001, ### p<0.0001 vs. HC  
1073 sample at matched time point.

1074

1075 **Figure 5. Effect of traumatic injury on neutrophil glucose uptake and metabolism.**  
1076 Glucose uptake by basal (A) or PMA-stimulated (B) neutrophils isolated from healthy  
1077 controls (HC) and trauma patients. MFI, Mean fluorescence intensity. (C-D) Comparison of  
1078 lactate concentration in cell-free supernatants collected from resting (C) or PMA stimulated  
1079 (D) neutrophils isolated from HC and trauma patients following a 3-hour *in vitro* culture.  
1080 Number of samples analysed are indicated below each time-point. \* p<0.01, \*\* p<0.001,  
1081 \*\*\* p<0.0001 vs. HC. (E-F) Expression of the glycolytic enzymes pyruvate kinase (PKM2)  
1082 and lactate dehydrogenase A (LDHA) in resting neutrophils isolated from HC (n=5) and  
1083 trauma patients 48-72 hours post-injury (n=14). Data are presented as representative Western  
1084 blots (E) and densitometry analysis of collated data for LDHA (F, top panel) and PKM2  
1085 (F, bottom panel). \* p<0.01 vs. HC.



1086 **Figure 6. Effect of mtDAMP pre-treatment on neutrophil NET generation.** (A) Serum  
1087 concentrations of the mitochondrial-derived N-formylated peptide ND6 in peripheral blood  
1088 samples acquired from healthy controls (HC) and trauma patients. Number of samples  
1089 analysed are indicated below each time-point. \*\*\* p<0.0001 vs. HC. (B) NET production by  
1090 PMA stimulated neutrophils pre-treated with 40 or 100 µg/ml mtDAMPs was assessed by  
1091 measuring DNA content of cell-free supernatants (n=5). \* p<0.01 vs. Vehicle. (C) MtDAMP-  
1092 induced inhibition of NET generation was confirmed by fluorescence microscopy (n=5; top  
1093 panel x20 magnification; bottom panel x40 magnification). (D) ROS generation by  
1094 neutrophils pre-treated with 100 µg/ml mtDAMPs or vehicle control was measured in  
1095 response to stimulation with 25 nM PMA using luminol-based chemiluminescence (n=10).  
1096 Data are presented as area under the curve (AUC) and are mean±SEM. \*\*\* p<0.0001 vs. PMA.  
1097

1098 **Figure 7. Treatment of neutrophils with mtDAMPs results in phosphorylation of**  
1099 **AMPK.** (A) Whole cell lysates prepared from purified neutrophils stimulated for 2-15  
1100 minutes with 100 µg/ml mtDAMPs were screened for phosphorylated AMPK. Western blot  
1101 in top panel is representative of 4 independent experiments. For densitometry analysis  
1102 \*\*\* p<0.0001 vs. 0 minutes. (B) AMPK phosphorylation in neutrophils treated for 1-hour with  
1103 the FPR-1 antagonist Cyclosporin H (CsH) or (C) or the CaMKK inhibitor STO-609 prior to  
1104 a 5-minute stimulation with 100 µg/ml mtDAMPs. Blots are representative of 5 (B) and 10  
1105 (C) independent experiments, with densitometric data depicted in the accompanying  
1106 histogram. \*\* p<0.001, \*\*\* p<0.0001 vs. vehicle. (D) Comparison of PMA-induced NET  
1107 formation by mtDAMP stimulated neutrophils pre-treated with the AMPK inhibitor  
1108 compound C or vehicle control (n=10). \*\* p<0.01 vs. PMA treated.  
1109

1110 **Figure 8. MtDAMP pre-treatment results in impaired neutrophil glycolysis.**  
1111 (A) Neutrophils pre-treated for 1-hour with 1 mM AICAR or vehicle control were stimulated  
1112 for 3 hours with 25 nM PMA, after which lactate concentrations were measured in cell-free  
1113 supernatants. Data are mean±SEM of 12 independent experiments. \*\*\* p<0.0001.  
1114 (B) Comparison of lactate concentrations in supernatants collected from neutrophils pre-  
1115 treated with 100 µg/ml mtDAMPs or vehicle-control and subsequently stimulated with 25  
1116 nM PMA for 1, 2 and 3 hours. Data are mean±SEM of 10 independent experiments.  
1117 \*\* p<0.001, \*\*\* p<0.0001 vs. vehicle.  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135

1136 **Table 1. Cohort demographics.**

1137

1138

	<b>Patients (n=62)</b>
Age, years	44 (19-95)
Male, n (%)	56 (90)
Time to pre-hospital sample, minutes post-injury	39 (13-59)
ISS	26 (9-57)
NISS	38 (9-75)
<b>AIS</b>	
Head, n (%)	24 (48)
Face, n (%)	17 (34)
Chest, n (%)	29 (58)
Abdomen, n (%)	14 (28)
Spine, n (%)	19 (38)
Pelvis, n (%)	8 (16)
Limbs, n (%)	28 (56)
Other, n (%)	6 (12)
<b>Mechanism of injury</b>	
Fall, n (%)	10 (16)
A/P, n (%)	13 (21)
Blunt, n (%)	2 (3)
RTC, n (%)	37 (60)
<b>Outcomes</b>	
ICU-free days	22 (0-30)
Hospital-free days	13 (0-29)
Mortality, n (%)	8 (13)

1139

1140 Data are expressed as mean (range) unless indicated otherwise.


1141 The number of data points for each clinical variable are: ISS and NISS, n=48; AIS scores,  
1142 n=50; ICU and hospital free days, n=53.

1143 ICU-free days and hospital-free days were calculated by 30 minus the number of days the  
1144 patient stayed in hospital).

1145 A/P, Assault/penetrating; AIS, Abbreviated injury scale; ISS, Injury severity score; ICU,  
1146 Intensive care unit; NISS, New injury severity score; RTC, Road traffic collision.

1147

1148 **Figure 1**  
 1149  
 1150




1151  
 1152  
 1153  
 1154  
 1155  
 1156  
 1157  
 1158  
 1159  
 1160  
 1161  
 1162  
 1163  
 1164  
 1165  
 1166  
 1167  
 1168

1169 **Figure 2**

1170

1171



1172

1173

1174

1175

1176

1177

1178


1179

1180


1181

1182


**A**




**B**



**C**



**D**



1184

1185

1186

1187

1188

1189

1190


1191

1192


1193

1194


**A**




**B**



**C**



**D**



1196

1197

1198

1199

1200

1201

1202


1203

1204

1205


1206

1207



1209  
 1210  
 1211  
 1212  
 1213  
 1214  
 1215  
 1216  
 1217  
 1218  
 1219  
 1220  
 1221

1222 **Figure 6**



1223

1224

1225

1226

1227

1228

1229

1230

1231


1232

1233

1234



1235 **Figure 7**



1236

1237

1238

1239

1240

1241

1242

1243


1244

1245

1246

1247

**A**



**B**

