

University of Montana

## ScholarWorks at University of Montana

---

Numerical Terradynamic Simulation Group  
Publications

Numerical Terradynamic Simulation Group

---

2010

### Global Landscape Freeze-Thaw Classification using Spaceborne Microwave Remote Sensing

Youngwook Kim

John S. Kimball

*University of Montana - Missoula*

Kyle C. McDonald

*City College of New York*

Joseph M. Glassy

Follow this and additional works at: [https://scholarworks.umt.edu/ntsg\\_pubs](https://scholarworks.umt.edu/ntsg_pubs)

**Let us know how access to this document benefits you.**

---

#### Recommended Citation

Kim, Youngwook; Kimball, John S.; McDonald, Kyle C.; and Glassy, Joseph M., "Global Landscape Freeze-Thaw Classification using Spaceborne Microwave Remote Sensing" (2010). *Numerical Terradynamic Simulation Group Publications*. 363.

[https://scholarworks.umt.edu/ntsg\\_pubs/363](https://scholarworks.umt.edu/ntsg_pubs/363)

This Poster is brought to you for free and open access by the Numerical Terradynamic Simulation Group at ScholarWorks at University of Montana. It has been accepted for inclusion in Numerical Terradynamic Simulation Group Publications by an authorized administrator of ScholarWorks at University of Montana. For more information, please contact [scholarworks@mso.umt.edu](mailto:scholarworks@mso.umt.edu).

Youngwook Kim<sup>1,2,\*</sup>, J. S. Kimball<sup>1,2</sup>, K. C. McDonald<sup>3</sup>, and J. Glassy<sup>4</sup>

(1) Flathead Lake Biological Station, The University of Montana 32125 Biostation Lane, Polson, MT, 59860-9659 \*Corresponding author: [youngwook.kim@ntsg.umt.edu](mailto:youngwook.kim@ntsg.umt.edu)

(2) Numerical Terradynamic Simulation Group, The University of Montana, Missoula, MT, 59812

(3) Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena CA, 91109

(4) Lupine Logic, Inc., Missoula, MT, 59812

## Introduction:

The freeze-thaw (F/T) status of the landscape is closely linked to surface energy budget and hydrological activity, the seasonal dynamics of vegetation growing seasons, terrestrial carbon budgets and land-atmosphere trace gas exchange. Satellite microwave radars and radiometers are well suited for global F/T monitoring due to insensitivity to signal degradation by atmospheric contamination and solar illumination effects, are uniquely capable of detecting the distinct change in landscape dielectric properties between predominantly frozen and thawed states, and provide a surrogate measure of a range of biophysical processes associated with the F/T signal, especially at high latitudes.


## Data and Methods:

	Passive		Active
Sensor	SSM/I	AMSR-E	SeaWinds
Platform	DMSP	EOS Aqua	QuikSCAT
Frequency	37GHz	36GHz	13.4GHz (Ku-band)
Polarization	V-pol	V-pol	V-pol/H-pol
Resolution	25x25km	25x25km	15x25km
Overpass	6am/pm	1:30am/pm	Daily average
Periods	1988-2007	2003-2007	2000-2007

## Surface air temperature data for FT\_ESDR calibration and verification

- (1) NASA DAO (2000-2006) 6-hour reanalysis (1° x 1.25°)
- (2) NCEP/NCAR (NNR, 1988-2007) 6-hour reanalysis (1.875° x 2°)
- (3) NCDC (1988-2007): daily summary of the day from WMO weather stations

## Ancillary data for quality control (QC) assessment




## F/T Algorithms: <sup>1</sup>Seasonal Threshold Approach (STA)


$$\Delta(\sigma \text{ or } T_b) = \frac{\sigma(t) - \sigma_{fr}}{\sigma_{th} - \sigma_{fr}} \quad \sigma_{fr} = \text{frozen reference state (mean } T_b \text{ or } \sigma \text{ in Jan)} \quad \Delta(\sigma \text{ or } T_b) > T \quad \text{Thawed}$$

$$\Delta(\sigma \text{ or } T_b) = \frac{\sigma(t) - \sigma_{th}}{\sigma_{fr} - \sigma_{th}} \quad \sigma_{th} = \text{non-frozen reference state (mean } T_b \text{ or } \sigma \text{ in Jul)} \quad \Delta(\sigma \text{ or } T_b) \leq T \quad \text{Frozen}$$

where  $T_b$  = brightness temperature and  $\sigma$  = backscatter



Example pixel-wise dynamic  $T_b$  [K] change thresholds [T] derived using SSM/I P37V (PM overpass) and NNR  $T_{min}$  (at left) and SSM/I A37V (AM overpass) and NNR  $T_{min}$  (at right) in 2004




## Global FT\_ESDR domain:

-defined by 7-year DAO reanalysis climatology using a <sup>2</sup>cold temperature constraint index [CCI, days yr<sup>-1</sup>]; -encompasses all vegetated regions where low temperatures are a major constraint to ecosystem processes.

$$iT_{Min} = \begin{cases} 0, & \text{if } T_{min} \leq T_{Minmin} \\ \frac{T_{min} - T_{Minmin}}{T_{Maxmax} - T_{Minmin}}, & \text{if } T_{Minmin} < T_{min} < T_{Maxmax} \\ 1, & \text{if } T_{min} \geq T_{Maxmax} \end{cases}$$

$$CCI = 365 - iT_{Min}$$

$T_{Maxmax}$  and  $T_{Minmin}$  vary by major land cover class using a global Biome Properties Lookup Table (BPLUT)



<sup>1</sup>Source: McDonald and Kimball, *EHS* 2004.


<sup>2</sup>Source: Jolly et al., *Global Change Biol* 2005.

## Global F/T classification criteria:


- The STA based SSM/I and AMSR-E AM/PM F/T classifications are produced as discrete frozen (0) or non-frozen (1) daily values;
- The combined SSM/I and AMSR-E F/T classifications are determined as frozen (0) or non-frozen (1) where the same F/T classification value occurs for both AM and PM retrievals, as daily output; as transitional (2) where AM frozen and PM non-frozen values occur, and inverse transitional (3) where AM non-frozen and PM frozen results occur;
- The SeaWinds daily F/T classification (frozen or non-frozen) is derived using the BYU daily global sigma-0 browse product from QuikSCAT L1B data.

## FT global accuracy assessment using NCDC stations:


Global mean annual FT classification accuracy (%) between 1988 and 2007




Seasonal pattern of daily mean global FT classification accuracy (%) in 2004




Spatial pattern of mean annual FT accuracy (%) relative to daily air temperatures from 3,701 NCDC stations in 2004



## Selected global daily FT patterns for 2004:




## 2004 annual non-frozen period:




## 20-year QC Map:

FT\_ESDR quality (QC; 1988-2007) assessment was developed to identify regions of relative high to low quality of F/T classification results in relation to general climate and landscape features; the QC map ranges from low (estimated accuracy < 70%) to best (estimated accuracy > 90%) quality categories; areas in white are outside the FT\_ESDR domain



## 20-year SSM/I global non-frozen period variation:



## Conclusions:

- The various microwave sensors produce similar FT spatial & temporal patterns, with 72-93% mean annual classification accuracy relative to NCDC stations, while global SSM/I F/T time series and corresponding annual frozen/non-frozen periods were quantified over a 20 year record;
- Approx. 66 million km<sup>2</sup> of the global land area are constrained by seasonally frozen temperatures; the average (1988-2007) seasonal progression of SSM/I derived frozen area ranges from 0.48 (±0.03) million km<sup>2</sup> (August) to 34.6 (±0.9) million km<sup>2</sup> (January);
- Daily F/T transitional areas occupy from 3.2 ±0.27 million km<sup>2</sup> (August) to 13.2 ±0.59 million km<sup>2</sup> (November);
- These results are being used to construct a consistent, systematic long-term (>20 yr) global daily record of F/T dynamics with well defined accuracy;
- The FT\_ESDR will be available online at (<http://freezethaw.ntsg.umt.edu/>) and archived to the NSIDC DAAC.

## Acknowledgements

Portions of this work were conducted at the University of Montana and Jet Propulsion Laboratory, California Institute of Technology under contract to NASA. This work was supported under the NASA Making Earth Science Data Records for Use in Research Environments (MEaSUREs) program (NNH062DA001N-MEaSUREs)