

Article

Antioxidant Activity and Phytochemical Characterization of *Senecio clivicolus* Wedd.

Immacolata Faraone ^{1,*} , Dilip K. Rai ² , Lucia Chiummianto ¹ , Eloy Fernandez ³,
Alka Choudhary ², Flavio Prinzo ¹ and Luigi Milella ^{1,*} 

¹ Department of Science, University of Basilicata, 85100 Potenza, Italy; lucia.chiummianto@unibas.it (L.C.); prinzo92@live.it (F.P.)

² Teagasc Food Research Centre, Ashtown, Dublin D15KN3K, Ireland; Dilip.Rai@teagasc.ie (D.K.R.); alkachoudhary12@gmail.com (A.C.)

³ Faculty of Tropical AgriSciences, Czech University of Life Sciences Prague, Kamýcká 129, 16500 Prague 6, Suchdol, Czech Republic; eloy@ftz.czu.cz

* Correspondence: immafaraone88@gmail.com (I.F.); luigi.milella@unibas.it (L.M.);
Tel.: +39-0971-20-5525 (L.M.); Fax: +39-0971-20-5503 (L.M.)

Academic Editor: Francesco Epifano

Received: 7 September 2018; Accepted: 28 September 2018; Published: 29 September 2018



Abstract: Antioxidant phytochemicals play a key role in oxidative stress control and in the prevention of related disorders, such as premature aging, degenerative diseases, diabetes, and cancer. The aim of this study was to investigate the potential antioxidant activity and the phytochemical profile of *Senecio clivicolus* Wedd., a perennial shrub, belonging to the Asteraceae family. Despite the wide interest of this family, this specie has not been investigated yet. *S. clivicolus* aerial parts were extracted with 96% ethanol. Then, the ethanol extract was fractionated by liquid/liquid extraction using an increasing solvents polarity. Total polyphenol and terpenoid contents were measured. Moreover, the antioxidant activity was evaluated by six different complementary in vitro assays. The Relative Antioxidant Capacity Index (RACI) was used to compare data obtained by different tests. The sample showing the highest RACI was subjected to characterization and quantitation of its phenolic composition using LC-MS/MS analysis. The ethyl acetate fraction, investigated by LC-MS/MS analysis, showed 30 compounds, most of them are chlorogenic acid and flavonoid derivatives. To the best of our knowledge, this is the first report about the evaluation of antioxidant activity and phytochemical profile of *S. clivicolus*, underlying the importance of this species as a source of health-promoting phytochemicals.

Keywords: Asteraceae; *Senecio clivicolus*; DPPH; beta-carotene bleaching; RACI; phenolic characterization; UHPLC-MS/MS; polyphenols; flavonoids; health-promoting compounds

1. Introduction

Plants are an important source of bioactive compounds with antioxidant capacity and during the last years, the interest for the use of natural products has significantly increased. The protective effects of plant secondary metabolites can be attributed to direct scavenging activities against reactive oxygen species (ROS), as well as to the induction of intracellular antioxidant effect [1,2]. In fact, different epidemiological studies have shown as the decrease of premature death and mortality from cancer or other chronic diseases are associated with antioxidant-rich diet including fruit, vegetables, and other botanicals [3,4]. Reactive oxygen and nitrogen species (RNS) are physiologically produced during metabolic processes and especially during electron transport chain reactions [4], low concentration of these species is essential for several biochemical processes. An important endogenous antioxidant system compensates the production of ROS, RNS, and other free radicals, but an overall increase

in cellular levels of these unstable molecules above the cells defenses results in oxidative stress that can ultimately cause cell death [5]. Oxidative stress is responsible for toxicity and damage of cell components, including nucleic acids, proteins, and lipids [3,6] and it has been recognized responsible for the onset of several diseases, including liver diseases, diabetes, neurodegenerative, and cardiovascular diseases. Thus, dietary antioxidants have been proposed as therapeutic agents to counteract oxidative stress-based diseases [4].

Senecio is one of the largest genus of the Asteraceae family, with a wide distribution in the world, consisting more than 1500 species of herbs, shrubs, vines, and trees. Many of these species grow in Bolivia, a central country of South America, with an extremely rich biodiversity of endemic species [7]. The traditional use of medicinal plants by various indigenous populations in Bolivia has been documented in the literature [8–13]. In particular, traditional medicine used extracts of leaves and roots of several *Senecio* species as a remedy for fever, cough, stomach pain, gastric ulcer, for the treatment of diabetes, skin wounds, and as a vasodilator, antiemetic, and anti-inflammatory agents [14].

Senecio clivicolus Wedd. grows in the mountain region of western Bolivia, where it is known as “*chini waycha*” or “*waycha negra*”. The natives used this perennial shrub as a remedy for stomach pain and diarrhea [14,15]. Moreover, this plant extract was used against fungal infections in the skin [16].

Pyrrolizidine alkaloids, eremophilanes, terpenes, and furanoeremophilanes were reported in previous phytochemical studies on *S. clivicolus* and other species belonging to *Senecio* genus [17–20]. These metabolites have several important properties as insect antifeedant, antifungal, cytotoxic, antioxidant, anti-inflammatory, and antimicrobial agents. Some furanoeremophilanes, such as cacalone compound, have a radical scavenging and antioxidant activity [14,21].

Following our investigation of medicinal plants belonging to the Asteraceae family [22,23], and due to the ethno medical use of this specie, the present study was focused on the investigation of total polyphenolic content (TPC), the potential antioxidant activity, the structural characterization and quantification of secondary metabolites present *S. clivicolus* aerial parts.

Extracts were analyzed, using six different complementary assays, for their radical-scavenging activity against 2,2-diphenyl-1-picrylhydrazyl (DPPH), 2,2'-azino-bis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS), superoxide anion (O_2^-) and nitric oxide (NO) radicals, for their ferric reducing antioxidant power (FRAP) and the capacity to inhibit lipid peroxidation by β -carotene bleaching test (BCB). Then, the phytochemical profile was performed by LC-ESI-MS/MS analysis and the identification and quantification of polyphenols were achieved using commercially available standards [24].

To the best of our knowledge, this is the first report about the evaluation of antioxidant activity and phytochemical profile of *S. clivicolus*, underling the importance of this specie as a source of health-promoting phytochemicals.

2. Results and Discussion

2.1. The Extraction Yield and the Influence of Solvents on Total Polyphenolic and Terpenoid Content

The dried aerial parts of *S. clivicolus* were extracted by dynamic maceration using 96% ethanol [25]. The extraction yield showed a value of 27.06%, higher than other reports. In fact, it has been reported that the extraction yield of the aerial parts of other species of *Senecio* with 95% ethanol (yield of 5.60% in *S. biafrae* [26] and 12.57% in *S. aegyptius* [27], methanol (yield of 13.85% in *S. gibbosus* [28]) or water (yield of 13.60% in *S. candicans* [29]) was considerably lower than values obtained in this study.

Then, liquid/liquid partition of a part of the crude ethanol extract (20.00 g) was performed for three times using increasing polarity solvents (*n*-hexane, chloroform, ethyl acetate and *n*-butanol); thus, a separation of the compounds was obtained based on the affinity with the solvent used [30,31]. The extraction yield of dry *n*-hexane (ScH), chloroform (ScC), ethyl acetate (ScEA), *n*-butanol (ScB) and water (ScW) fractions is reported in Figure 1. The chloroform fraction showed the highest extraction

yield ($63.39 \pm 5.23\%$); instead, the water and ethyl acetate fractions demonstrated the lower extraction yields, 7.60 ± 0.42 and $6.87 \pm 0.39\%$, respectively.




Figure 1. Extraction yields of *S. clivicolus* EtOH extract partitioned fractions. Results were expressed as mean \pm standard deviation of the triplicate experiments. Samples are crude ethanol extract (Sc EtOH), *n*-hexane fraction (ScH), chloroform fraction (ScC), ethyl acetate fraction (ScEA), *n*-butanol fraction (ScB), and water fraction (ScW).

Total phenolic content (TPC) of Sc EtOH and its fractions was carried out based on the reaction of the samples with Folin-Ciocalteu reagent, which results in a blue colored solution whose intensity was directly proportional to the amount of phenolic compounds present. Gallic acid was used as standard and results were expressed as milligram of gallic acid equivalents per gram of dry extract (mg GAE/g) and shown in Figure 2. TPC ranged from 24.32 ± 0.43 to 170.11 ± 1.49 mg GAE/g in ScW and ScEA fractions, respectively. Out of all samples, ScEA had the highest value, followed by ScB and Sc EtOH (82.50 ± 1.94 and 48.31 ± 1.32 mg GAE/g, respectively).




Figure 2. Total polyphenolic content (TPC) and total terpenoids content (TTeC). Results were expressed as mean \pm standard deviation in mg of gallic acid equivalents per gram of dried sample (mg GAE/g), and in mg of linalool equivalents per gram of dried sample (mg LE/g). In each tests, the values with the same letter are not significant different at the $p < 0.05$ level, according to one-way analysis of variance (ANOVA). Samples are crude ethanol extract (Sc EtOH), *n*-hexane fraction (ScH), chloroform fraction (ScC), ethyl acetate fraction (ScEA), *n*-butanol fraction (ScB), and water fraction (ScW).

Moreover, total terpenoids content (TTeC) was evaluated for their characteristic reddish brown color at 538 nm. The linalool, a monoterpene, was used as standard reagent and results (Figure 2) were expressed as mg of linalool equivalents per gram of dry sample (mg LE/g). TTeC ranged from 470.77 ± 3.61 to 2074.24 ± 12.76 mg LE/g in ScEA and ScC fractions, respectively. ScC, followed by ScB, ScW, and Sc EtOH, exhibited higher values.

Results demonstrated as the yield of phenolic and terpenoid clearly differs on the basis of solvent polarity. Fractions obtained by using more polar solvents reported the highest total polyphenols, indicating that the majority of polyphenolic compounds in the aerial parts of *S. clivicolus* could be of polar nature, data congruent with what reported previously in literature [32].

2.2. Antioxidant Activity

Low concentrations of biological radicals are important in the human body; in fact, they are involved in biological functions as vascular homeostasis, neurotransmission, antimicrobial, antioxidant and antitumor activities. In contrast, the overproduction of biological radicals is associated with several diseases [1]. Phenolic compounds are gaining attention due to their significant antioxidant activities. Different in vitro assays were reported for the measurement of antioxidant activity on foods and plants and it has been demonstrated as more than one method is necessary to elucidate the antioxidant capacity of samples because these assays differ in the principles and experimental conditions. In this study, the antioxidant activity of the Sc EtOH extract and its fractions (ScH, ScC, ScEA, ScB, and ScW) were tested using six different complementary in vitro antioxidant assays.

In particular, the radical scavenging activity was evaluated by using synthetic cationic and neutral (ABTS and DPPH) and physiological (superoxide anion and nitric oxide) radicals. The ScEA showed the highest radical scavenging-activity in all investigated tests (Table 1). In particular, the ScEA reported 409.53 ± 9.53 mg TE/g and 317.53 ± 5.81 mg TE/g values in ABTS and DPPH assays respectively, followed by ScB and Sc EtOH. Instead, in both assays, the lowest activity was found in ScH and ScC fractions.

Table 1. Results of ABTS, DPPH and super oxide (SO) scavenging activity, ferric reducing antioxidant power (FRAP) and β -carotene bleaching (BCB) of *S. clivicolus* samples.

Samples	ABTS (mgTE/g)	DPPH (mgTE/g)	SO (IC ₂₅ mg/mL)	FRAP (mgTE/g)	BCB %AA
Sc EtOH	137.87 ± 1.45^d	63.42 ± 0.78^b	0.37 ± 0.02^d	93.08 ± 1.12^d	53.11 ± 0.45^{de}
ScH	28.94 ± 2.50^a	nc	$0.16 \pm 0.01^{b,c}$	12.98 ± 1.04^a	4.75 ± 0.23^a
ScC	55.93 ± 2.24^b	12.70 ± 0.94^a	0.14 ± 0.01^b	23.90 ± 1.32^b	44.58 ± 0.96^c
ScEA	409.53 ± 9.53^f	317.53 ± 5.81^d	0.08 ± 0.00^a	507.66 ± 5.26^f	55.82 ± 2.22^e
ScB	208.37 ± 3.21^e	119.54 ± 6.71^c	0.20 ± 0.01^c	184.18 ± 4.59^e	51.70 ± 1.97^d
ScW	107.01 ± 0.94^c	55.58 ± 1.07^b	0.64 ± 0.04^e	53.41 ± 2.55^c	23.28 ± 0.70^b

Samples are crude ethanol extract (Sc EtOH), *n*-hexane fraction (ScH), chloroform fraction (ScC), ethyl acetate fraction (ScEA), *n*-butanol fraction (ScB) and water fraction (ScW). Data are expressed as means \pm standard deviation from three experiments; mg GAE/g = mg of gallic acid equivalents per gram of dried sample; mg TE/g = mg of Trolox equivalents per gram of dried sample; IC₂₅ mg/mL = concentration of the sample required to inhibit the activity of the radical by 25%; %AA = percentage of antioxidant activity at initial sample concentration of 1 mg/mL; different superscripts in the same row indicate significant difference ($p < 0.05$); nc = not calculable.

The ability of samples to scavenge superoxide anion and nitric oxide was expressed as IC₂₅, as it was not possible to reach 50% inhibition at tested concentrations, and the results were compared with ascorbic acid. The *S. clivicolus* extract and fractions caused a dose-dependent inhibition in the superoxide anion assay (SO) and ScEA fraction showed the highest inhibition, that means the lowest IC₂₅ value (IC₂₅ of 0.08 ± 0.00 mg/mL), followed by ScC, ScH, and ScB (Table 1). These samples showed a higher activity than that measured for ascorbic acid (IC₂₅ of 0.26 ± 0.02 mg/mL).

The scavenging ability against the biological nitric oxide (NO) was not detectable in any fraction, but ScEA showed a dose-dependent inhibition ability (IC₂₅ of 1.11 ± 0.05 mg/mL); its value was four times lower than ascorbic acid (IC₂₅ 4.78 ± 0.09 mg/mL), demonstrating its effectiveness.

The reducing ability of samples in a redox reaction was assessed using the FRAP test. ScEA fraction also in this case showed the highest activity (507.66 ± 5.26 mg TE/g) followed by ScB fraction (184.18 ± 4.59 mg TE/g). ScC and ScH were again the less active (Table 1).

To get a broader overview of the antioxidant potential of the plant complex of *S. clivicolus* samples, the β -carotene bleaching test (BCB) was conducted. The phenolic compounds compete with β -carotene for binding to the radical derived from linoleic acid and prevent the destruction of the conjugated system of the molecule responsible for coloring, for which, the higher is the antioxidant activity of the extract, the greater is the concentration of β -carotene in solution. Results were expressed as percentage of antioxidant activity (%AA) at the initial sample concentration of 1 mg/mL (Table 1). The analysis evidenced that the most active sample in BCB test was the ScEA, followed by Sc EtOH and ScB (55.82 ± 2.22 , 53.11 ± 0.45 and 51.70 ± 1.97 %AA, respectively). While, there was a low activity for the apolar ScH fraction.

Moreover, correlations were calculated based on the average of the results by the Pearson test. This test calculates the linear correlation coefficient (r), a dimensionless number between -1 and 1 , inclusive, reflecting the extent of a linear relationship between two datasets. The more the value of the coefficient comes close to the extremes, the more the correlation is present, in a positive or negative manner. The correlation value of 0 indicates no linear correlation. High correlation was found between the total polyphenol content and radical-scavenging activity ($r_{\text{TPC}/\text{ABTS}} = 0.98$; $r_{\text{TPC}/\text{DPPH}} = 0.98$; $r_{\text{TPC}/\text{NO}} = 0.92$; $r_{\text{TPC}/\text{SO}} = 0.77$) and reducing power ($r_{\text{TPC}/\text{FRAP}} = 0.99$) demonstrating as the polyphenols are the compounds mostly involved in the antioxidant activity [33]. The highest correlation was observed between the reducing power of the samples and the radical-scavenging activity by ABTS and DPPH methods ($r_{\text{FRAP}/\text{ABTS}} = 0.99$ and $r_{\text{FRAP}/\text{DPPH}} = 1.00$). Polyphenols are less involved in the inhibition of lipid peroxidation ($r_{\text{TPC}/\text{BCB}} = 0.61$). The antioxidant activity determined by the BCB method, a lipophilic system constituted by the emulsion β -carotene-linoleic acid, generally has a low correlation with the phenolic compounds and the other methods tested because of the effectiveness of lipophilic compounds [34]. Instead, terpenoids are clearly less involved in tested activities (Table 2).

Table 2. Pearson correlation coefficients calculated among tested *Senecio clivicolus* extract and fractions.

	TPC	TTeC	ABTS	DPPH	SO	NO	FRAP	BCB
TPC	1.00							
TTeC	-0.69	1.00						
ABTS	0.98	-0.64	1.00					
DPPH	0.98	-0.68	0.99	1.00				
SO	0.77	-0.56	0.65	0.70	1.00			
NO	0.92	-0.75	0.89	0.93	0.84	1.00		
FRAP	0.99	-0.70	0.99	1.00	0.75	0.94	1.00	
BCB	0.61	0.01	0.65	0.59	0.29	0.41	0.58	1.00

Total phenolic content (TPC); total terpenoids content (TTeC); ABTS assay; DPPH assay; Super oxide anion scavenging activity (SO); nitric oxide radical scavenging activity (NO); Ferric reducing antioxidant power assay (FRAP); β -carotene bleaching assay (BCB).

The results of the antioxidant activity obtained by the above-described chemical methods have been integrated by calculating the Relative Antioxidant Capacity Index (RACI). The RACI index allows the comparison of phytocomplex antioxidant capacity derived from different chemical methods (ABTS, DPPH, SO, NO, FRAP, and BCB) even the TPC. It has recently been shown, that the results obtained with the method of Folin-Ciocalteu reagent can be interpreted to determine the total content of polyphenols, and as an alternative way to measure the total reducing capacity of the samples since the Folin reagent reacts with any reducing substance present in solution [22]. This index is dimensionless and measures the distance between the average of the results obtained and the raw data expressed in standard deviation units. RACI values obtained agrees with the results obtained so far (Figure 3) evidencing as ScEA fraction presented the highest value (1.73), followed by the ScB

(0.16). The ScH fraction presented the lowest index (-0.73) and, therefore, a relative lack of antioxidant activity. In particular, the RACI values may be related to the high total polyphenol content of the ethyl acetate fraction.




Figure 3. Relative Antioxidant Capacity Index (RACI) of *Senecio clivicolus* samples. Samples are crude ethanol extract (Sc EtOH), *n*-hexane fraction (ScH), chloroform fraction (ScC), ethyl acetate fraction (ScEA), *n*-butanol fraction (ScB), and water fraction (ScW).

To date, no study reported the antioxidant activity of *S. clivicolus* extracts, but Conforti et al. [28] evaluated the antioxidant activity of *Senecio gibbosus* aerial parts. This study reported as the methanol extract and ethyl acetate fraction had potent antioxidant property on DPPH radical as well as on lipid peroxidation. In particular, the paper reported an IC_{50} of 0.01 mg/mL for the ethyl acetate fraction of *S. gibbosus* on DPPH and showed higher inhibition than that ScEA (IC_{50} of 0.10 ± 0.00 mg/mL) obtained in present study. It is possible to explain these results with the different methods used to obtain the ethyl acetate fraction. In fact, Conforti et al. extracted the plant materials with methanol; then, the methanolic extract was acidified with 2.50% H_2SO_4 and partitioned with *n*-hexane, dichloromethane, and ethyl acetate. Moreover, in this paper the *n*-hexane fraction tested by DPPH assay was inactive, such as ScH.

Similarly, the in vitro antioxidant activity of *Senecio inaequidens* and *Senecio vulgaris* by the DPPH assay showed that the ethyl acetate fraction gave the 61.60% and 44.57% of inhibition, respectively, at a concentration of 0.31 mg/mL [35]. In present study, ScEA showed a stronger inhibition compared to both species tested by Conforti et al. ($90.55 \pm 0.61\%$ of inhibition at 0.31 mg/mL).

2.3. Identification and Quantification of Polyphenols by Mass Spectrometry

The compounds previously identified in the aerial parts of *S. clivicolus* belong mainly to the class of alkaloids, eremophilanes and furanoeremophilanes [17,25]. To evaluate the compounds responsible of the measured antioxidant activity, ScEA fraction partitioned from ethanol extract of *S. clivicolus* aerial parts was subject to mass spectrometry analysis in negative ionization [36]. The LC-MS profile of ScEA is shown in Figure 4.




Figure 4. Ethyl acetate fraction *Senecio clivicolus* base peak intensity chromatogram (BPI).

Several compounds (>26) were detected and tentative identification of most of them could be reached through accurate masses and fragmentation pattern and aided by the existing literature. As many as 19 of the compounds were of chlorogenic acid derivatives. Only eight phenolic compounds could be identified in the ethyl acetate fraction by comparing their retention times with those of the available commercial standards. In particular, these compounds were phenolic derivatives of benzoic acid (4-hydroxybenzoic acid (**1**)), phenylpropanoids, and in particular cinnamic acid derivatives (chlorogenic acid methyl ester (**3**), 3,5-di-*O*-caffeoyl quinic acid (**7**), 3,4-di-*O*-caffeoyl quinic acid (**8**), caffeic acid (**10**) and chlorogenic acid (**15**)), and flavonols (rutin (**4**) and quercetin-3-*O*-glucoside (**5**)).

The most abundant was the 3,5-di-*O*-caffeoyl quinic acid (45.44 ± 0.91 mg/g DW), followed by 3,4-di-*O*-caffeoyl quinic acid (19.27 ± 0.68 mg/g DW). These phenolic compounds are known for their antioxidant properties [37–42].

As mentioned earlier, a number of the tentatively identified compounds were chlorogenic acid derivatives; in particular, five of the compounds with m/z 529.13 tentatively identified as feruloyl-caffeoylquinic acid isomers (**9**, **10**, **12**, **13**, and **20**). Upon fragmentation by CID, these compounds produced the ions at m/z 367, 353, 293, 193, 191, 179, 173, 161, 134, and 111 [43,44]. The signals at 9.23 (**13**) and 11.87 (**21**) min m/z 793.40 indicating the presence of an extra hexose sugar unit with respect to m/z 529.13 and were tentatively identified as chlorogenic acid methylester hexoside derivative; they produced ions at m/z 529, 191, 179, 173, and 161 [45]. The ESI-MS signals at m/z 601.22 (**15**, **16**), 617.23 (**15**, **16**), 779.23 (**17**), 763.23 (**18**, **19**), 819.26 (**22**), 807.30 (**20**, **23**), and 735.32 (**24**) were tentatively identified as chlorogenic acid derivatives; they produced the ions at m/z 353, 191, 179, 161 and 135 [46]. The ESI-MS signal at m/z 479.07 (**2**) was tentatively identified as quercetagenin-*O*-glucoside because it produced the ion m/z 317 of quercetagenin, and the typical ions m/z 166 and 139 [47]. The ESI-MS signals at 7.17 (**4**) and 9.61 (**14**) min m/z 493.09 were tentatively identified as mearnsenin-*O*-hexoside isomers and they gave dominant product ions m/z 478, 331 and 315 [48]. The ESI-MS signal at m/z 477.10 (**6**) was tentatively identified as isorhamnetin glycoside and it gave dominant product ions m/z 462, 315 and 299 [49]. Instead, the ESI-MS signal at m/z 519.10 (**11**)

was tentatively identified as isorhamnetin-acetyl-glucoside because it produced the ion m/z 315 due to the loss of acetyl-glucose residue [50] (Table 3).

Previous reports evidenced as identified compounds are able to exert a strong antioxidant activity. In particular, Hung T.M. et al. (2006) reported 3,5-di-*O*-caffeoyl quinic acid and 3,4-di-*O*-caffeoyl quinic acid as potent scavengers of the neutral radical DPPH. These compounds demonstrated to be more potent than BHT used as a positive control [39].

Also, flavonoid derivative identified compounds in *S. clivicolus* EA fraction, showed to be effective when tested for their antioxidant activity. For example, rutin exhibited a strong DPPH radical scavenging activity and at 0.05 mg/mL showed 90.40% inhibition compared with BHT (58.80%) as reported from Yang et al. [40]. Moreover, rutin showed to be also effective in preventing lipid peroxidation. Razivi et al. [41] instead showed the cytotoxic, antimicrobial, and antioxidant activities of flavonol quercetin 3-*O*-glucoside.

The antioxidant activity analysis was effective in identifying the important polyphenolic contributors to the antioxidant capacity of *S. clivicolus* aerial parts, as assessed by TPC, TTeC, and *in vitro* antioxidant assays. The ethyl acetate fraction demonstrated the highest relative antioxidant capacity index and was subjected to mass spectrometry analysis. In the present study, the known antioxidant compounds such as phenolic derivatives of benzoic acid, cinnamic acid derivatives, and flavonoids were reported for the first time in *Senecio* genus. The most abundant of them were the known antioxidant 3,5-di-*O*-caffeoyl quinic acid and 3,4-di-*O*-caffeoyl quinic acid.

Table 3. Liquid chromatography-tandem mass spectrometry (LC-MS/MS) of ethyl acetate fraction of *Senecio clivicolus*. Identification of compounds based on *m/z*, and fragmentation pattern and retention time of standards. Quantities of the detected phenolic compounds were determined using commercial standards.

Peak No.	RT (min)	ESI (–) MS Observed	ESI (–) MS Calc.	Molecular Formula	MS/MS	Tentative Identity	mg/g DW	Reference
1	5.39	137.0242	137.0239	C ₇ H ₅ O ₃	93	4-Hydroxybenzoic acid	2.25 ± 1.31	[36]
2	6.64	479.0779	479.0826	C ₂₁ H ₁₉ O ₁₃	317, 166, 139	Quercetagenin- <i>O</i> -glucoside	nq	[47]
3	6.92	367.1029	367.1029	C ₁₇ H ₁₉ O ₉	161, 85	Chlorogenic acid methylester	2.57 ± 0.07	[36]
4	7.17	493.0982	493.0982	C ₂₂ H ₂₁ O ₁₃	478, 331, 315, 287, 271, 244, 166	Mearnsetin- <i>O</i> -glucoside isomer	nq	[48]
		609.1461	609.1456	C ₂₇ H ₂₉ O ₁₆	300, 285, 271, 255, 179, 151	Rutin	0.16 ± 0.02	[36]
5	7.25	463.0894	463.0877	C ₂₁ H ₁₉ O ₁₂	300, 271, 255, 179, 151	Quercetin-3- <i>O</i> -glucoside	0.84 ± 0.05	[36,51]
6	7.80	477.1067	477.1033	C ₂₂ H ₂₁ O ₁₂	462, 315, 299, 271, 254, 243, 227, 151	Isorhamnetin glycoside	nq	[49]
7	7.91	515.1174	515.1190	C ₂₅ H ₂₃ O ₁₂	179, 135	3,5-di- <i>O</i> -caffeoyl quinic acid	45.44 ± 0.91	[36]
8	8.13	515.1169	515.1190	C ₂₅ H ₂₃ O ₁₂	179, 135	3,4-di- <i>O</i> -caffeoyl quinic acid	19.27 ± 0.68	[36]
9	8.66	529.1370	529.1346	C ₂₆ H ₂₅ O ₁₂	367, 349, 191, 179, 173, 161, 135, 133, 101, 93	Feruloyl-caffeoylquinic acid isomer	nq	[43,44]
10	8.86	179.0353	179.0344	C ₉ H ₇ O ₄	135, 79	Caffeic acid	1.88 ± 0.13	[36]
		529.1370	529.1346	C ₂₆ H ₂₅ O ₁₂	367, 349, 191, 179, 173, 161, 135, 101	Feruloyl-caffeoylquinic acid isomer	nq	[43]
11	8.99	519.1039	519.1139	C ₂₄ H ₂₃ O ₁₃	504, 315, 299, 285, 271, 243, 191	Isorhamnetin-acetyl-glucoside	nq	[50]
12	9.12	529.1370	529.1346	C ₂₆ H ₂₅ O ₁₂	367, 349, 191, 179, 173, 161, 135, 101	Feruloyl-caffeoylquinic acid isomer	nq	[43]
13	9.23	529.1370	529.1346	C ₂₆ H ₂₅ O ₁₂	367, 349, 191, 179, 173, 161, 135, 101	Feruloyl-caffeoylquinic acid isomer	nq	[43]
		793.4029	793.4010	C ₄₁ H ₆₁ O ₁₅	529, 397, 353, 219, 191, 179, 173, 161, 101, 71	Chlorogenic acid methylester hexoside derivative	nq	[45]
14	9.61	493.0982	493.0982	C ₂₂ H ₂₁ O ₁₃	331, 316, 179, 161, 135, 133, 101	Mearnsetin- <i>O</i> -glucoside isomer	nq	[48]
15/16	9.72/10.21	353.0885	353.0873	C ₁₆ H ₁₇ O ₉	191, 179, 173, 93, 85	Chlorogenic acid	2.33 ± 0.45	[36]
		617.2367	617.2387	C ₃₅ H ₃₇ O ₁₀	353, 245, 191, 179, 173, 161, 135	Chlorogenic acid derivative	nq	[46]
		601.2267	601.2285	C ₃₁ H ₃₇ O ₁₂	439, 353, 263, 191, 179, 173, 161, 135, 85	Dicafeoyl-methoxyoxaloylquinic acids	nq	[46]
17	10.55	779.2360	779.2340	C ₄₃ H ₃₉ O ₁₄	515, 375, 353, 335, 191, 179, 173, 161, 155, 135, 111, 93	Chlorogenic acid derivative	nq	[46]
18/19	10.88/11.21	763.2333	763.2332	C ₅₀ H ₃₅ O ₈	515, 353, 191, 179, 173, 161, 135, 110	Dicafeoylquinic acid derivative	nq	[46]
20	11.50	529.1370	529.1346	C ₂₆ H ₂₅ O ₁₂	367, 353, 293, 193, 191, 179, 173, 161, 134, 111	Feruloyl-caffeoylquinic acid isomer	nq	[43]
		807.3002	807.3017	C ₄₆ H ₄₇ O ₁₃	353, 335, 191, 179, 173, 161, 155, 135	Chlorogenic acid derivative	nq	[46]

Table 3. Cont.

Peak No.	RT (min)	ESI (–) MS Observed	ESI (–) MS Calc.	Molecular Formula	MS/MS	Tentative Identity	mg/g DW	Reference
21	11.87	793.2888	793.2860	C ₄₅ H ₄₅ O ₁₃	353, 191, 179, 173, 161, 155, 135	Chlorogenic acid derivative	nq	[45]
22	12.25	819.2629	819.2618	C ₂₈ H ₅₁ O ₂₇	353, 335, 191, 179, 173, 161, 155, 135	Chlorogenic acid derivative	nq	[46]
23	12.79	807.3002	807.3017	C ₄₆ H ₄₇ O ₁₃	353, 335, 191, 179, 173, 161, 155, 135	Chlorogenic acid derivative	nq	[46]
24	12.99	735.3240	735.3228	C ₃₇ H ₅₁ O ₁₅	353, 335, 191, 179, 173, 161, 135	Chlorogenic acid derivative	nq	[46]
25	14.70	675.3726	675.3744	C ₃₇ H ₅₅ O ₁₁	415, 397, 277, 235, 161, 143, 125, 119, 113, 101, 89	unknown	nq	
26	16.25	480.3110	480.3087	C ₂₇ H ₄₄ O ₇	255, 242, 224, 168, 153, 79	unknown	nq	

nq: not quantified.

3. Materials and Methods

3.1. Chemicals, Reagents and Equipment

Solvents as *n*-butanol, chloroform, *n*-hexane, hydrochloric acid, ethanol, ethyl acetate, glacial acetic acid, methanol and phosphoric acid were purchased from Carlo Erba (Milan, Italy).

Acetonitrile and formic acid were purchased from Merck (Wicklow, Ireland). Folin-Ciocalteu reagent 2N, sodium carbonate, 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS), potassium persulfate, 2,2-diphenyl-1-picrylhydrazyl (DPPH), potassium phosphate monobasic, β -nicotinamide adenine dinucleotide reduced form (NADH), phenazine methosulfate (PMS), nitrotetrazolium blue chloride (NBT), sodium nitroprusside dehydrate (SNP), sulfanilamide, *N*-(1-Naphthyl)ethylenediamine dihydrochloride, sodium acetate trihydrate, 2,4,6-tripyridyl-*s*-triazine (TPTZ), iron (III) chloride ($\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$), β -carotene, linoleic acid, and Tween 20 were purchased from Sigma-Aldrich (Milan, Italy).

Standards as 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox), ascorbic acid, butylhydroxytoluen (BHT), gallic acid, and Leucine-Enkephalin were purchased from Sigma-Aldrich (Milan-Italy) and Merck (Wicklow-Ireland), respectively. Standard as 3,4-di-*O*-caffeoyl quinic acid, 3,5-di-*O*-caffeoyl quinic acid, 4-hydroxybenzoic acid, caffeic acid, chlorogenic acid, chlorogenic acid methyl ester, quercetin-3-*O*-glucoside, and rutin were purchased from Extrasynthese (Genay, France).

Water was deionized using a Milli-Q water purification system (Millipore, Bedford, MA, USA).

All spectrophotometric measurements were done in 96-well microplates or cuvettes on a UV-VIS spectrophotometer (SPECTROstar^{Nano} BMG Labtech, Ortenberg, Germany).

LC-MS/MS analysis were performed on a Q-ToF Premier mass spectrometer (Waters Corporation, Milford, MA, USA) coupled to an Alliance 2695 HPLC system (Waters Corporation, Milford, MA, USA). Mass spectrometry quantification of the polyphenols was performed using multiple reaction monitoring (MRM) experiments using Waters Acquity (Waters Corporation, Milford, MA, USA) ultra-high performance liquid chromatography coupled with tandem mass spectrometry (UHPLC-MS/MS).

3.2. Plant Material and Sample Preparation

The aerial parts of *S. clivicolus* (Asteraceae family) were collected in the 2014, near the Aymaya population/community (18.45° S to 66.46° W; 3750 msnm), Bustillo province, Potosí department, Bolivia (Provincia di Nor Yungas, GPS coordinates –16.190040, –67.884402) dried at room temperature and crushed. A voucher specimen has been stored at the University of La Paz (number SC108-2014). Dried plant material (110.00 g) have been extracted by maceration in a darkness shaker set at 25 °C using 96% ethanol for 24 h at a solid to solvent ratio of 1:10 (*w/v*) per extraction (3 times). The extracts were filtered through a Buchner funnel (0.45 μm) and combined; subsequently supernatants have been dried using a rotary evaporator with a water bath set at 37 °C.

Twenty grams of ethanol extract (Sc EtOH) were solubilized in water (400.00 mL) and subjected to liquid/liquid partitioning in triplicate using an equal volume of *n*-hexane, chloroform, ethyl acetate and *n*-butanol in order to separate the compounds on the basis of increasing solvent polarity. Then the *n*-hexane, chloroform, ethyl acetate, *n*-butanol, and water fractions (ScH, ScC, ScEA, ScB and ScW, respectively) were dried and stored in darkness at room temperature until further use.

Sc EtOH and all the above-mentioned fractions were analyzed for their polyphenolic and terpenoids contents and their antioxidant activities.

3.3. Total Phenolic Content (TPC)

Total phenolic content (TPC) of *S. clivicolus* samples was determined by Folin-Ciocalteu assay as reported by Todaro et al. [52] with slight modification. The results were expressed as milligrams of gallic acid equivalents per gram of dried extract (mg GAE/g).

3.4. Total Terpenoid Content (TTeC)

The total terpenoid content (TTeC) was determined using linalool as standard reagent [53]. The absorbance was measured at 538 nm and the results were expressed as milligrams of linalool equivalents per gram of dried sample (mg LE/g).

3.5. Radical-Scavenging Activity

The radical scavenging capacity of *S. clivicolus* was evaluated by four different complementary in vitro assays. In particular, the synthetic coloured ABTS⁺ and DPPH[•] radicals, and the biological super oxide anion (O₂⁻) and nitric oxide (NO) radicals. The electron transfer involves reduction of a colored oxidant and the capacity of the samples to scavenge the radicals was monitored by spectrophotometer and quantified in Trolox equivalents, used as standard [15].

3.5.1. ABTS Assay

The free radical-scavenging capacity of all *S. clivicolus* samples was studied using the 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt (ABTS) radical assay, by the procedure described by Armentano et al. [54] with slight modification. The reaction between ABTS salt and potassium persulfate generates the ABTS⁺ radical after 16 h of incubation at room temperature. The reaction for scavenging the cationic radicals from the samples was monitored at 734 nm and the results were expressed as milligrams of Trolox equivalents per gram of dried extract (mg TE/g) by the Trolox standard curve.

3.5.2. DPPH Assay

The radical-scavenging ability of samples was also evaluated by in vitro 2,2-diphenyl-1-picrylhydrazyl (DPPH) neutral radical and, also in this case, Trolox was used as standard. The reaction was monitored at 515 nm and the data were expressed as milligrams of Trolox equivalents per gram of dried extract (mg TE/g) [22].

3.5.3. Super Oxide Anion Scavenging Activity (SO)

Superoxide anion radicals (O₂⁻) were generated in vitro by the NADH/PMS system as described by Russo et al. [55]. The scavenging activity of samples on the inhibition of formazan formation was monitored at 560 nm by spectrophotometer and ascorbic acid was used as positive control. The results were expressed as the concentration inhibiting 25% of radical inhibition in mg/mL (IC₂₅).

3.5.4. Nitric Oxide Radical Scavenging Activity (NO)

The nitric oxide radical (NO) was spontaneously generated at physiological pH from nitroprusside. Then, the nitric oxide interacts with oxygen to give nitrite ions that can be determined spectrophotometrically by Griess reagent [55]. Results were expressed as IC₂₅ of radical inhibition in mg/mL and ascorbic acid was used as positive control.

3.6. Ferric Reducing Antioxidant Power Assay (FRAP)

FRAP reagent was prepared fresh before the experiment by mixing 300 mM sodium acetate buffer at pH 3.60, 20 mM FeCl₃·6H₂O in distilled water and 10 mM TPTZ in 40 mM HCl in a ratio of 10:1:1. The reaction was monitored at 593 nm and Trolox was used as reference antioxidant standard. FRAP values were expressed as milligrams of Trolox equivalents per gram of dried extract (mg TE/g) [56].

3.7. β-Carotene Bleaching Assay (BCB)

Lipid peroxidation inhibition was evaluated by the β-carotene bleaching method (BCB) [57]. The β-Carotene emulsion (950.00 μL) was mixed with 50.00 μL of sample at 1.00 mg/mL and

BHT was used as positive control, then the mixture (250.00 μL) was transferred into a 96-well plate. Outer wells were filled with 250.00 μL of water to provide a large thermal mass because the reaction was temperature-sensitive and close temperature control throughout the plate was essential in this assay [58]. The microplate was placed at 50 °C for 3 h and the absorbance was measured at 470 nm at 0', 30', 60', 90', 120', 150', and 180'. The results were expressed as percentage of β -carotene bleaching inhibition (% Antioxidant Activity, %AA).

3.8. Identification and Quantification by Liquid Chromatography Mass Spectrometry

LC-MS characterization was carried out on a Q-ToF Premier mass spectrometer (Waters Corporation, Milford, MA, USA) coupled to an Alliance 2695 HPLC system (Waters Corporation, Milford, MA, USA) as described previously [59,60]. Accurate mass measurements of the analytes were achieved through the use of an internal reference compound (leucine-enkephalin). The separation of the compounds was performed on an Atlantis T3 C18 column (Waters Corporation, Milford, USA, 100.00 mm \times 2.10 mm; 3.00 μm particle size) maintained at 40 °C and using 0.10% aqueous formic acid (solvent A) and 0.10% formic acid in acetonitrile (solvent B). A stepwise gradient from 10% to 90% solvent B was applied at a flow rate of 0.30 mL/min for 25 min. Electrospray mass spectra data were acquired on a negative ionization mode for a mass range m/z 100 to m/z 1000. Cone voltage and capillary voltage were set at 30 V and 3 kV, respectively. Collision induced fragmentation (CID) of the analytes was achieved using 12 eV to 30 eV energy with argon as the collision gas.

For the quantification of the polyphenols in the ethyl acetate fraction of *S. clivicolus*, Waters Acquity (Waters Corporation, Milford, MA, USA) ultra-high performance liquid chromatography coupled with tandem mass spectrometry (UHPLC-MS/MS) was used. The compounds were separated on a Waters Acquity HSS T3 C18 column (2.10 \times 100.00 mm; 1.80 μm particle size) using a binary solvent system consisting of water containing 0.10% formic acid (mobile phase A) and acetonitrile containing 0.10% formic acid (mobile phase B). The following gradient program was carried out: 0–2.50 min 2% B, 2.50–3.00 min 10% B, 3.00–7.50 min 15% B, 7.50–8.50 min 35% B, 8.50–9.50 min 98% B and 9.50–10.00 min 2% B at a flow rate of 0.50 mL/min. The injection volume for all the samples and the standards was 3.00 μL . All the standards in the concentration ranging from 0.01 to 50.00 $\mu\text{g}/\text{mL}$ were dissolved in 80% methanol. Multiple reaction monitoring (MRM) quantitative method was developed for each of the standard compound using the Waters Intellistart software, and the quantifications of the data were carried out using the Waters TargetLynx™ software. The ionization source conditions were as follows: capillary voltage 3 kV, cone voltage 35 V, source temperature 150 °C, desolvation temperature 350 °C, desolvation gas flow 200 L/h, cone gas flow 50 L/h, and collision gas flow 0.10 mL/min. The column temperature was maintained at 40 °C.

3.9. Statistical Analysis

All data were expressed as mean \pm standard deviation (SD) of three independent experiments performed in triplicate. To verify the correlation among used methods, the p value of 0.05 or less was considered significant and calculated by one-way analysis of variance (ANOVA). Pearson coefficient and all other statistical calculations were determined using GraphPad Prism 5 Software (San Diego, CA, USA).

4. Conclusions

In this report the ethanol extract of *S. clivicolus* aerial parts was subjected to liquid/liquid fractionation using solvent with increasing polarity. The antioxidant activity, determined by six complementary in vitro assays, showed different values among fractions. Relative Antioxidant Capacity Index (RACI), evidenced *S. clivicolus* ethyl acetate fraction as the most active. More in detail, fraction reported activity was: *n*-hexane < water < chloroform < *n*-butanol < ethyl acetate. Moreover, this study reports the profile of the phenolic compounds in *S. clivicolus* ethyl acetate fraction. The phytochemical investigation allowed the identification or tentative identification of 30 compounds

and confirms that the LC-MS-based profiling is a powerful technique for the phenolic characterization. This first report on *S. clivicolus* phenolics demonstrated as it could be considered a rich source of health promoting compounds, particularly chlorogenic acid, and flavonoid derivatives. Hung et al. (2006) demonstrated as this compounds are able to exert a protective role against oxidative stress. In fact chlorogenic acid, the most abundant isomer among caffeoylquinic acid isomers, is a biologically active polyphenol, which play several biological activities such as antioxidant [42].

In conclusion, based on the observed results in conjunction with the existing literature, it is anticipated that *S. clivicolus*, especially its polar extracts, might be used for mitigating human and animal diseases. The presence of potential nutraceuticals is suitable and promising for the development of safe food products, natural additives and cosmetics. Its compounds, previously tested for their biological activity, seems to explain the most of the measured antioxidant activity. This will further trigger extensive research for better understanding of the impact of *S. clivicolus* phenolic compounds on health.

Author Contributions: Conceptualization: I.F., L.C., and L.M.; data curation: I.F., L.C., and F.P.; funding acquisition: D.K.R., L.C., and L.M.; investigation, I.F., A.C., and F.P.; methodology: I.F., D.K.R., L.C., E.F., A.C., F.P., and L.M.; project administration, L.C. and L.M.; resources: D.K.R. and E.F.; validation: E.F. and A.C.; writing—original draft: I.F. and L.M.; writing—review and editing: I.F. and L.M.

Funding: This research was supported by the University of Basilicata, Project “Monitoraggio delle acque marine costiere e profonde in Basilicata” D.G.R. 1490 of 4/12/2014.

Acknowledgments: The authors are grateful to Victor CussyPoma, National University Siglo XX, Llalagua, Bolivia for his help in the identification of the species.

Conflicts of Interest: The authors declare that there is no conflict of interests regarding the publication of this paper.

References

1. Patel Rajesh, M.; Patel Natvar, J. In vitro antioxidant activity of coumarin compounds by DPPH, super oxide and nitric oxide free radical scavenging methods. *J. Adv. Pharm. Educ. Res.* **2011**, *1*, 52–68.
2. Milutinović, V.; Niketić, M.; Ušjak, L.; Nikolić, D.; Krunic, A.; Zidorn, C.; Petrović, S. Methanol extracts of 28 *Hieracium* species from the Balkan Peninsula—Comparative LC–MS analysis, chemosystematic evaluation of their flavonoid and phenolic acid profiles and antioxidant potentials. *Phytochem. Anal.* **2018**, *29*, 30–47. [[CrossRef](#)] [[PubMed](#)]
3. Pizarro, J.G.; Folch, J.; Vazquez De la Torre, A.; Verdager, E.; Junyent, F.; Jordán, J.; Pallas, M.; Camins, A. Oxidative stress-induced DNA damage and cell cycle regulation in B65 dopaminergic cell line. *Free Radic. Res.* **2009**, *43*, 985–994. [[CrossRef](#)] [[PubMed](#)]
4. Scalbert, A.; Manach, C.; Morand, C.; Rémésy, C.; Jiménez, L. Dietary polyphenols and the prevention of diseases. *Crit. Rev. Food Sci.* **2005**, *45*, 287–306. [[CrossRef](#)] [[PubMed](#)]
5. Sakagami, H.; Satoh, K.; Hatano, T.; Yoshida, T.; Okuda, T. Possible role of radical intensity and oxidation potential for gallic acid-induced apoptosis. *Anticancer Res.* **1997**, *17*, 377–380. [[PubMed](#)]
6. Forman, H.J.; Davies, K.J.; Ursini, F. How do nutritional antioxidants really work: Nucleophilic tone and para-hormesis versus free radical scavenging in vivo. *Free Radic. Biol. Med.* **2014**, *66*, 24–35. [[CrossRef](#)] [[PubMed](#)]
7. Araujo, N.; Mü Ller, R.; Nowicki, C.; Ibisch, P. *Análisis de Vacíos de Representatividad del Sistema Nacional de Áreas Protegidas*; FAN: Santa Cruz, CA, USA, 2005.
8. Deharo, E.; Bourdy, G.; Quenevo, C.; Munoz, V.; Ruiz, G.; Sauvain, M. A search for natural bioactive compounds in Bolivia through a multidisciplinary approach. Part V. Evaluation of the antimalarial activity of plants used by the Tacana Indians. *J. Ethnopharmacol.* **2001**, *77*, 91–98. [[CrossRef](#)]
9. Muñoz, V.; Sauvain, M.; Bourdy, G.; Callapa, J.; Bergeron, S.; Rojas, I.; Bravo, J.; Balderrama, L.; Ortiz, B.; Gimenez, A. A search for natural bioactive compounds in Bolivia through a multidisciplinary approach: Part I. Evaluation of the antimalarial activity of plants used by the Chacobo Indians. *J. Ethnopharmacol.* **2000**, *69*, 127–137. [[CrossRef](#)]
10. Bourdy, G.; Oporto, P.; Gimenez, A.; Deharo, E. A search for natural bioactive compounds in Bolivia through a multidisciplinary approach: Part VI. Evaluation of the antimalarial activity of plants used by Isoceno-Guarani Indians. *J. Ethnopharmacol.* **2004**, *93*, 269–277. [[CrossRef](#)] [[PubMed](#)]

11. Macía, M.J.; García, E.; Vidaurre, P.J. An ethnobotanical survey of medicinal plants commercialized in the markets of La Paz and El Alto, Bolivia. *J. Ethnopharmacol.* **2005**, *97*, 337–350. [[CrossRef](#)] [[PubMed](#)]
12. Thomas, E.; Semo, L.; Morales, M.; Noza, Z.; Nuñez, H.; Cayuba, A.; Noza, M.; Humaday, N.; Vaya, J.; Van Damme, P. Ethnomedicinal practices and medicinal plant knowledge of the Yuracarés and Trinitarios from indigenous territory and national park Isiboro-Sécure, Bolivian Amazon. *J. Ethnopharmacol.* **2011**, *133*, 153–163. [[CrossRef](#)] [[PubMed](#)]
13. Hajdu, Z.; Hohmann, J. An ethnopharmacological survey of the traditional medicine utilized in the community of Porvenir, Bajo Paraguá Indian Reservation, Bolivia. *J. Ethnopharmacol.* **2012**, *139*, 838–857. [[CrossRef](#)] [[PubMed](#)]
14. Floegel, A.; Kim, D.O.; Chung, S.J.; Koo, S.I.; Chun, O.K. Comparison of ABTS/DPPH assays to measure antioxidant capacity in popular antioxidant-rich US foods. *J. Food Comp. Anal.* **2011**, *24*, 1043–1048. [[CrossRef](#)]
15. Fournet, A.; Barrios, A.A.; Munoz, V. Leishmanicidal and trypanocidal activities of Bolivian medicinal plants. *J. Ethnopharmacol.* **1994**, *41*, 19–37. [[CrossRef](#)]
16. Bustamante, G.; Escalante, L.; Mejia, U.; Valdivia, M.; Soria, I. *Estudio Etnobotánico y Actividad Antimicrobiana de las Plantas Medicinales de los Valles Bajos de Cochabamba*; Universidad Mayor de San Simón: Cochabamba, Bolivia, 2001; Volume 78.
17. Pelsner, P.B.; Nordenstam, B.; Kadereit, J.W.; Watson, L.E. An ITS phylogeny of tribe *Senecioneae* (Asteraceae) and a new delimitation of *Senecio* L. *Taxon* **2007**, *56*, 1077–1104. [[CrossRef](#)]
18. Sánchez-Muñoz, B.A.; Aguilar, M.I.; King-Díaz, B.; Rivero, J.F.; Lotina-Hennsen, B. The sesquiterpenes β -caryophyllene and caryophyllene oxide isolated from *Senecio salignus* act as phyto-growth and photosynthesis inhibitors. *Molecules* **2012**, *17*, 1437–1447. [[CrossRef](#)] [[PubMed](#)]
19. Oladipupo, L.; Adebola, O. Chemical composition of the essential oils of the flowers, leaves and stems of two *Senecio polyanthemoides* Sch. Bip. samples from South Africa. *Molecules* **2009**, *14*, 2077–2086. [[CrossRef](#)] [[PubMed](#)]
20. Balzaretto, V.; Arancibia, A.; Marchiaro, A.; Arce, M.; Feijóo, M. Variation in the composition of the essential oil of *Senecio filaginoides* DC. *Molecules* **2000**, *5*, 459–461. [[CrossRef](#)]
21. Krasovskaya, N.; Kulesh, N.; Denisenko, V. Natural antioxidants. Furanoeremophilanes from *Cacalia* roots. *Chem. Nat. Compd.* **1989**, *25*, 545–548. [[CrossRef](#)]
22. Milella, L.; Bader, A.; De Tommasi, N.; Russo, D.; Braca, A. Antioxidant and free radical-scavenging activity of constituents from two *Scorzonera* species. *Food Chem.* **2014**, *160*, 298–304. [[CrossRef](#)] [[PubMed](#)]
23. Saltos, M.B.V.; Puente, B.F.N.; Faraone, I.; Milella, L.; De Tommasi, N.; Braca, A. Inhibitors of α -amylase and α -glucosidase from *Andromachia igniaria* Humb. & Bonpl. *Phytochem. Lett.* **2015**, *14*, 45–50. [[CrossRef](#)]
24. Fraige, K.; Dametto, A.C.; Zeraik, M.L.; de Freitas, L.; Saraiva, A.C.; Medeiros, A.I.; Castro-Gamboa, I.; Cavalheiro, A.J.; Silva, D.H.S.; Lopes, N.P. Dereplication by HPLC-DAD-ESI-MS/MS and screening for biological activities of *Byrsonima* species (Malpighiaceae). *Phytochem. Anal.* **2018**, *29*, 196–204. [[CrossRef](#)] [[PubMed](#)]
25. Dávila, M.; Sterner, O.; Hinojosa, N. Furanoeremophilanes from *Senecio clivicolus* Wedd. In *Revista Boliviana de Química*; Universidad Mayor de San Andrés: San Andrés, Bolivia, 2013; Volume 30, pp. 80–83, ISSN 0250-5460.
26. Lienou, L.L.; Telefo, P.B.; Bayala, B.; Yemele, D.M.; Lemfack, M.C.; Mouokeu, C.; Goka, S.C.; Tagnie, R.S.; Fewou, P. Effect of ethanolic extract of *Senecio biafrae* on puberty onset and fertility in immature female rat. *Cameroon J. Exp. Biol.* **2010**, *6*, 101–109. [[CrossRef](#)]
27. Hassan, W.; Gendy, A.; Al-youssef, H.; El-Shazely, A. Chemical constituents and biological activities of *Senecio aegyptius* var. *discoideus* Boiss. *J. Biosci. C* **2012**, *67*, 144–150. [[CrossRef](#)]
28. Conforti, F.; Marrelli, M.; Statti, G.; Menichini, F. Antioxidant and cytotoxic activities of methanolic extract and fractions from *Senecio gibbosus* subsp. *gibbosus* (GUSS) DC. *Nat. Prod. Res.* **2006**, *20*, 805–812. [[CrossRef](#)] [[PubMed](#)]
29. Hariprasath, L.; Raman, J.; Nanjian, R. Gastroprotective effect of *Senecio candicans* DC on experimental ulcer models. *J. Ethnopharmacol.* **2012**, *140*, 145–150. [[CrossRef](#)] [[PubMed](#)]
30. Saeed, N.; Khan, M.R.; Shabbir, M. Antioxidant activity, total phenolic and total flavonoid contents of whole plant extracts *Torilis leptophylla* L. *BMC Complement. Altern. Med.* **2012**, *12*, 221. [[CrossRef](#)] [[PubMed](#)]
31. Saada, M.; Falleh, H.; Catarino, M.D.; Cardoso, S.M.; Ksouri, R. Plant growth modulates metabolites and biological activities in *Retama raetam* (Forssk.) Webb. *Molecules* **2018**, *23*, 2177. [[CrossRef](#)] [[PubMed](#)]

32. Nakamura, M.; Ra, J.H.; Jee, Y.; Kim, J.S. Impact of different partitioned solvents on chemical composition and bioavailability of *Sasa quelpaertensis* Nakai leaf extract. *J. Food Drug Anal.* **2017**, *25*, 316–326. [[CrossRef](#)] [[PubMed](#)]
33. Sytar, O.; Boško, P.; Živčák, M.; Brestic, M.; Smetanska, I. Bioactive phytochemicals and antioxidant properties of the grains and sprouts of colored wheat genotypes. *Molecules* **2018**, *23*, 2282. [[CrossRef](#)] [[PubMed](#)]
34. Mariod, A.; Matthaeus, B. Fatty acids, tocopherols, sterols, phenolic profiles and oxidative stability of *Cucumis melo* var. *agrestis* oil. *J. Food Lip.* **2008**, *15*, 56–67. [[CrossRef](#)]
35. Conforti, F.; Loizzo, M.R.; Statti, G.A.; Houghton, P.J.; Menichini, F. Biological properties of different extracts of two *Senecio* species. *Int. J. Food Sci. Nutr.* **2006**, *57*, 1–8. [[CrossRef](#)] [[PubMed](#)]
36. Hossain, M.B.; Rai, D.K.; Brunton, N.P.; Martin-Diana, A.B.; Barry-Ryan, C. Characterization of phenolic composition in *Lamiaceae* spices by LC-ESI-MS/MS. *J. Agric. Food Chem.* **2010**, *58*, 10576–10581. [[CrossRef](#)] [[PubMed](#)]
37. Wang, H.; Nair, M.G.; Strasburg, G.M.; Booren, A.M.; Gray, J.I. Novel antioxidant compounds from *Tart Cherries* (*Prunus cerasus*). *J. Nat. Prod.* **1999**, *62*, 86–88. [[CrossRef](#)] [[PubMed](#)]
38. McDonald, S.; Prenzler, P.D.; Antolovich, M.; Robards, K. Phenolic content and antioxidant activity of olive extracts. *Food Chem.* **2001**, *73*, 73–84. [[CrossRef](#)]
39. Hung, T.M.; Na, M.; Thuong, P.T.; Su, N.D.; Sok, D.; Song, K.S.; Seong, Y.H.; Bae, K. Antioxidant activity of caffeoyl quinic acid derivatives from the roots of *Dipsacus asper* Wall. *J. Ethnopharmacol.* **2006**, *108*, 188–192. [[CrossRef](#)] [[PubMed](#)]
40. Yang, J.; Guo, J.; Yuan, J. In vitro antioxidant properties of rutin. *LWT-Food Sci. Technol.* **2008**, *41*, 1060–1066. [[CrossRef](#)]
41. Razavi, S.M.; Zahri, S.; Zarrini, G.; Nazemiyeh, H.; Mohammadi, S. Biological activity of quercetin-3-O-glucoside, a known plant flavonoid. *Russ. J. Bioorg. Chem.* **2009**, *35*, 376–378. [[CrossRef](#)]
42. Naveed, M.; Hejazi, V.; Abbas, M.; Kamboh, A.A.; Khan, G.J.; Shumzaid, M.; Ahmad, F.; Babazadeh, D.; FangFang, X.; Modarresi-Ghazani, F. Chlorogenic acid (CGA): A pharmacological review and call for further research. *Biomed. Pharmacother.* **2018**, *97*, 67–74. [[CrossRef](#)] [[PubMed](#)]
43. Han, J.; Ye, M.; Qiao, X.; Xu, M.; Wang, B.R.; Guo, D.A. Characterization of phenolic compounds in the Chinese herbal drug *Artemisia annua* by liquid chromatography coupled to electrospray ionization mass spectrometry. *J. Pharm. Biomed. Anal.* **2008**, *47*, 516–525. [[CrossRef](#)] [[PubMed](#)]
44. Clifford, M.N.; Johnston, K.L.; Knight, S.; Kuhnert, N. Hierarchical scheme for LC-MSⁿ identification of chlorogenic acids. *J. Agric. Food Chem.* **2003**, *51*, 2900–2911. [[CrossRef](#)] [[PubMed](#)]
45. Yépez, A.M.; de Ugaz, O.L.; Alvarez, C.M.; De Feo, V.; Aquino, R.; De Simone, F.; Pizza, C. Quinovic acid glycosides from *Uncaria guianensis*. *Phytochemistry* **1991**, *30*, 1635–1637. [[CrossRef](#)]
46. Jaiswal, R.; Kuhnert, N. Identification and characterization of two new derivatives of chlorogenic acids in *Arnica* (*Arnica montana* L.) flowers by high-performance liquid chromatography/tandem mass spectrometry. *J. Agric. Food Chem.* **2011**, *59*, 4033–4039. [[CrossRef](#)] [[PubMed](#)]
47. Ibdah, M.; Zhang, X.H.; Schmidt, J.; Vogt, T. A novel Mg²⁺-dependent O-methyltransferase in the phenylpropanoid metabolism of *Mesembryanthemum crystallinum*. *J. Biol. Chem.* **2003**, *278*, 43961–43972. [[CrossRef](#)] [[PubMed](#)]
48. Song, Y.; Desta, K.T.; Kim, G.S.; Lee, S.J.; Lee, W.S.; Kim, Y.H.; Jin, J.S.; Abd El-Aty, A.M.; Shin, H.C.; Shim, J.H.; et al. Polyphenolic profile and antioxidant effects of various parts of *Artemisia annua* L. *Biomed. Chromatogr.* **2016**, *30*, 588–595. [[CrossRef](#)] [[PubMed](#)]
49. Schieber, A.; Keller, P.; Streker, P.; Klaiber, I.; Carle, R. Detection of isorhamnetin glycosides in extracts of apples (*Malus domestica* cv. “Brettacher”) by HPLC-PDA and HPLC-APCI-MS/MS. *Phytochem. Anal.* **2002**, *13*, 87–94. [[CrossRef](#)] [[PubMed](#)]
50. Carazzone, C.; Mascherpa, D.; Gazzani, G.; Papetti, A. Identification of phenolic constituents in red chicory salads (*Cichorium intybus*) by high-performance liquid chromatography with diode array detection and electrospray ionisation tandem mass spectrometry. *Food Chem.* **2013**, *138*, 1062–1071. [[CrossRef](#)] [[PubMed](#)]
51. Krzyzanowska-Kowalczyk, J.; Pecio, Ł.; Mołdoch, J.; Ludwiczuk, A.; Kowalczyk, M. Novel phenolic constituents of *Pulmonaria officinalis* L. LC-MS/MS Comparison of Spring and Autumn metabolite profiles. *Molecules* **2018**, *23*, 2277. [[CrossRef](#)] [[PubMed](#)]

52. Todaro, L.; Russo, D.; Cetera, P.; Milella, L. Effects of thermo-vacuum treatment on secondary metabolite content and antioxidant activity of poplar (*Populus nigra* L.) wood extracts. *Ind. Crops Prod.* **2017**, *109*, 384–390. [[CrossRef](#)]
53. Ghorai, N.; Chakraborty, S.; Gucchait, S.; Saha, S.K.; Biswas, S. Estimation of total Terpenoids concentration in plant tissues using a monoterpene, linalool as standard reagent. *Protoc. Exch.* **2012**, *5*. [[CrossRef](#)]
54. Armentano, M.F.; Bisaccia, F.; Miglionico, R.; Russo, D.; Nolfi, N.; Carmosino, M.; Andrade, P.B.; Valentão, P.; Diop, M.S.; Milella, L. Antioxidant and proapoptotic activities of *Sclerocarya birrea* [(A. Rich.) Hochst.] methanolic root extract on the hepatocellular carcinoma cell line HepG2. *Biomed. Res. Int.* **2015**, *2015*. [[CrossRef](#)] [[PubMed](#)]
55. Russo, D.; Valentão, P.; Andrade, P.B.; Fernandez, E.C.; Milella, L. Evaluation of antioxidant, antidiabetic and anticholinesterase activities of *Smallanthus sonchifolius* landraces and correlation with their phytochemical profiles. *Int. J. Mol. Sci.* **2015**, *16*, 17696–17718. [[CrossRef](#)] [[PubMed](#)]
56. Dekdouk, N.; Malafronte, N.; Russo, D.; Faraone, I.; De Tommasi, N.; Ameddah, S.; Severino, L.; Milella, L. Phenolic compounds from *Olea europaea* L. possess antioxidant activity and inhibit carbohydrate metabolizing enzymes in vitro. *Evid. Based Complement. Alternat. Med.* **2015**, *2015*. [[CrossRef](#)] [[PubMed](#)]
57. Fidelis, Q.C.; Faraone, I.; Russo, D.; Aragão Catunda, F.E.; Vignola, L.; de Carvalho, M.G.; de Tommasi, N.; Milella, L. Chemical and biological insights of *Ouratea hexasperma* (A. St.-Hil.) Baill.: A source of bioactive compounds with multifunctional properties. *Nat. Prod. Res.* **2018**, *1–4*. [[CrossRef](#)] [[PubMed](#)]
58. MikaMi, I.; YaMaguChi, M.; Shinmoto, H.; Tsushida, T. Development and validation of a microplate-based β -carotene bleaching assay and comparison of antioxidant activity (AOA) in several crops measured by β -carotene bleaching, DPPH and ORAC assays. *Food Sci. Technol. Res.* **2009**, *15*, 171–178. [[CrossRef](#)]
59. Rico, D.; Diana, A.B.M.; Milton-Laskibar, I.; Fernández-Quintela, A.; Silván, J.M.; Rai, D.K.; Choudhary, A.; Peñas, E.; de Luis, D.A.; Martínez-Villaluenga, C. Characterization and in vitro evaluation of seaweed species as potential functional ingredients to ameliorate metabolic syndrome. *J. Funct. Foods* **2018**, *46*, 185–194. [[CrossRef](#)]
60. Choudhary, A.; Naughton, L.M.; Dobson, A.D.; Rai, D.K. HPLC-ESI-MS/MS characterisation of metabolites produced by *Pseudovibrio* sp. W64, a marine sponge-derived bacterium isolated from Irish waters. *Rapid Commun. Mass Spectrom.* **2018**, *32*, 1737–1745. [[CrossRef](#)] [[PubMed](#)]

Sample Availability: Samples of *Senecio clivicolus* are available from the authors and stored as reported above.



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).