

## Soil-derived greenhouse gas emissions as influenced by farming management

Andreas Gattinger<sup>a,b</sup>, C. Skinner<sup>a</sup>, H.-M. Krause<sup>a</sup>, M. Krauss<sup>a</sup> and Paul Mäder<sup>a</sup>

<sup>a</sup> *Research Institute of Organic Agriculture (FiBL)/Department of Soil Sciences, Frick, Switzerland*

<sup>b</sup> *Institute of Crop Science and Plant Breeding II, Organic Farming with focus on Sustainable Soil Use/Justus-Liebig University Giessen, Giessen, Germany*

Agricultural practices contribute considerably to emissions of greenhouse gases. So far, knowledge on the impact of organic compared to non-organic farming on soil-derived nitrous oxide (N<sub>2</sub>O) and methane (CH<sub>4</sub>) emissions is rather limited. Meta-studies studies show, that organically managed soils emit less N<sub>2</sub>O and take up more CH<sub>4</sub> than those under non-organic management. This in contrast of that what has been found in the laboratory with soil material from the DOK trial. When subjected to the same water-filled pore space (90%) and N fertilization level (40 kg N ha<sup>-1</sup>), BIOORG showed a higher N<sub>2</sub>O production potential than CONMIN. This can be related to the higher soil C content and higher microbial activity in BIOORG than in CONMIN soil. Production of N<sub>2</sub> was similar in BIOORG and CONMIN and significantly lower in NOFERT, most likely due to significantly decreased pH inhibiting N<sub>2</sub>O reduction. This caused the greatest N<sub>2</sub>O/(N<sub>2</sub>O + N<sub>2</sub>) ratios in NOFERT (0.88 ± 0.02) followed by BIOORG (0.79 ± 0.01) and CONMIN (0.68 ± 0.02) (p < 0.001).

Furthermore, we investigated N<sub>2</sub>O and CH<sub>4</sub> fluxes with manual chambers during 571 days in a grass-clover– silage maize – green manure cropping sequence in the field, making use of the contrasting farming systems of the DOK trial. We compared two organic farming systems – BIODYN and BIOORG with the two non-organic systems CONMIN and CONFYM – all reflecting Swiss farming practices—together with the unfertilised control NOFERT. We observed a 40.2% reduction of N<sub>2</sub>O emissions per hectare for organic compared to non-organic systems. In contrast to current knowledge, yield-scaled cumulated N<sub>2</sub>O emissions under silage maize were similar between organic and non-organic systems. Cumulated on area scale we recorded under silage maize a modest CH<sub>4</sub> uptake for BIODYN and CONMIN and high CH<sub>4</sub> emissions for CONFYM. We found that, in addition to N input, quality properties such as pH, soil organic carbon and microbial biomass significantly affected N<sub>2</sub>O emissions. This study showed that organic farming systems can be a viable measure contributing to greenhouse gas mitigation in the agricultural sector.