

Weeds community structure on the rice field (*Oryza sativa* L.) in bulusari village, Sayung district, Demak regency

by Amnan Haris

Submission date: 20-Aug-2019 08:19AM (UTC+0700)

Submission ID: 1161578282

File name: Haris_2019_J._Phys.__Conf._Ser._1217_012177.pdf (497.24K)

Word count: 2409

Character count: 12635

PAPER • OPEN ACCESS**Weeds community structure on the rice field (*Oryza sativa* L.) in bulusari village, Sayung district, Demak regency**

1

To cite this article: A Haris *et al* 2019 *J. Phys.: Conf. Ser.* **1217** 012177View the [article online](#) for updates and enhancements.**IOP | ebooks™**

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

Weeds community structure on the rice field (*Oryza sativa* L.) in bulusari village, Sayung district, Demak regency

A Haris¹, S Utami¹ and Murningsih¹

¹Biologi Department, Faculty of Science and Mathematics, Diponegoro University
Jl. Prof. Sudharto, SH., Tembalang, Semarang, Central Java, Indonesia, 50275
E-mail: amnan.haris43@gmail.com

Abstract. Rice weeds are disturbing plants that grow together with rice and otherwise well-controlled, it will reduce the quantity and quality of crops. The aim of this research is to analyze the community structure of weed rice in 8-week-old. Benefits of this research are expected to provide information about the types of rice weeds and their characteristics. Research was conducted in June 2018 in Bulusari Village, Sayung District, Demak Regency; and Ecology and Biosystematics Laboratory, Biology Departement, Diponegoro University. Method of this research is a random sampling method used a plot sized 1 m x 1 m as many as 18 plots. Results of the research were the highest Important Value Index is *Echinochloa colonum* (L) Link species and rice weed diversity index is low.

1. Introduction

Rice (*O. sativa* L.) is the main food crop in Asia, especially Indonesia, because the majority of its population consume rice as a staple food. Rice production must be increased to meet food demand due to population growth. One of the important things to increase rice production is minimizing crop losses caused by weeds. Yield loss caused by weeds were exceeds than yield loss caused by plant pests and diseases. In addition, competition with weeds also decreases the quality of rice grains [1].

Rice weeds are disturbing plants that grow with rice and otherwise well-controlled, it will reduce the quantity and quality of crops [2]. Rice weeds are associated and will fight each other for the nutrients needed by rice, especially if the amount of nutrients is very limited. Potential competition for rice weeds are increasingly supported by breeding both vegetatively and generatively [3].

According to research conducted by Fitri *et al* [4] in Nagari Singkarak, Solok Regency, West Sumatra, there were 10 types of rice weeds : *Eclipta prostrata* (L.), *Lindernia ciliata* (Colsm.) Pennell, *Ludwigia hyssopifolia* (G. Don) Exell, *Hedyotis diffusa* Willd, *Echinochloa colonum* (L.) Link, *Echinochloa crus-galli* (L.) Beauv, *Leptochloa chinensis* (L.) Nees, *Cyperus difformis* L., *Cyperus iria* L., *Fimbristylis miliacea* (L.) Vahl. The highest Important Value Index was (IVI) owned by *Fimbristylis miliacea* (L.) Vahl which was 108.50% and the lowest was *Hedyotis diffusa* Willd of 1.61%. This shows that *Fimbristylis miliacea* (L.) Vahl have an important role in maintaining the balance of weed communities in these fields. The diversity index is 1.6 which means the diversity of weeds in the rice field is moderate. Rice weeds always in the fields, one of which is the rice fields in Demak.



Demak is the third largest rice production center in Central Java with a production of 632.751 ton in 2015. The wide of rice field area is 94,877 hectares in 14 existing districts [5]. But rice cultivation activities in Demak still rely on rainwater. So, the farmer will only plant rice when the rainy season arrives [6].

The aims of this research is to determine the structure of rice weed communities in Bulusari Village, Sayung District, Demak Regency by calculating the Important Value Index (IVI) and Shannon-Wiener Diversity Index (H'). The expected benefits from this research are it can be used by farmers as a guide in weed management to increase rice productivity.

2. Material and Methods

2.1 Time and Place of Research

Reserach was conducted in June 2018 on 700 m² of inorganic rice field in Bulusari Village, Sayung District, Demak Regency. Identification of the rice weeds at the Ecology and Biosystematics Laboratory, Department of Biology, Faculty of Science and Mathematics, Diponegoro University.

2.2 Tools and materials

Tools used were camera, meters, ropes, pegs, scissors, plastic bags, stationery and rice weed identification book "Weed of Rice in Indonesia" (1987) by Mohamad Soerjani, Achmad Jahja G.H. Kostermans, Gembong Tjitrosoepomo. The materials used are labels, envelopes and rice weed samples.

2.3 Method

Rice weed sampling used a random method. Rice weeds were allowed to grow with rice plants until the age of 8 weeks. Size plot used for the research is 1 mx 1 m, 18 plots. Recorded on the types of rice weeds that grow and counted the number of individuals of each type of rice weed. The types of rice weeds were brought to the Ecology and Biosystematics laboratory to be identified. Identification of rice weed samples using the "Weed of Rice in Indonesia" book by Soerjani *et al.* [7].

Environmental factors were measured including light intensity with lux meter, pH and soil moisture with soil tester, air temperature with thermometer, humidity with hygrometer and altitude with altimeter.

2.4 Data analysis

The data obtained were analyzed by calculating type dominance, Important Value Index (IVI) and Shannon-Wiener Diversity Index

$$\begin{aligned} & \text{Dominance of Plant Types} \\ &= \frac{\text{Number of individuals in a family}}{\text{Number of individuals in all families}} \times 100\% \end{aligned}$$

Dominant families if they have a percentage of > 20% total individuals, codominant 10% - 20%, and not dominant <10% [8].

$$\begin{aligned} & \text{Important Value Index} \\ & \text{Species density [9],} \\ & \text{Absolute density (AD)} \\ &= \frac{\text{Number of Individuals of a Type}}{\text{Area of observation plot}} \end{aligned}$$

$$\begin{aligned} & \text{Relative Density (RD)} \\ &= \frac{\text{Absolute density of a type}}{\sum \text{Total absolute density of all types}} \times 100\% \end{aligned}$$

Frequency [10].

$$\text{Absolute Frequency (AF)} = \frac{\text{The number of plots occupied by a type}}{\text{The total number of observation plots}}$$

Relative Frequency (RF) =

$$= \frac{\text{Absolute frequency of a Type}}{\text{Absolute Frequency of All Types}} \times 100\%$$

$$\text{IVI} = \text{RD} + \text{RF}$$

Information :

IVI = Important Value Index

RD = Relative Density

RF = Relative Frequency

Shannon-Wiener Diversity Index [12]:

$$H' = - \sum (n_i / N) \ln (n_i / N)$$

Information :

H' = Shannon-Wiener diversity index

n_i = Number of i-type individuals

N = Total number of individuals of all types


The level of species diversity uses the Fachrul criteria [13] :

- H' value > 3 : species diversity is high.
- $1 \leq H' \leq 3$ value: species diversity is moderate.
- H' < 1 value : species diversity is low or small.

3. Results and discussion

3.1 Weed Composition

The results showed that at 8 weeks of age, Rice weeds were consisting of 6 families, 10 species, and 310 individuals. There are 5 types of grasses with 258 individuals, there are 2 types of sedges with 41 individuals, and 4 broad-leaves groups with 11 individuals.



Remarks: * Grasses, ** Sedges, *** Broad-leaves

Figure 1. Percentage of Rice Weed Types in Bulusari Village, Sayung Districts, Demak Regency

In Figure 1, The dominant plant types of the rice weed were *Echinochloa colonum* (L) Link. It caused by its ability to produce large amounts of seedlings and seeds. Tanasale [14] stated that weeds which can survive were weeds that can reproduce vegetatively and generatively whereas the non-dominant families were Portulacaceae and Solanaceae. It because Portulacaceae and Solanaceae need wider growth space, of course, it will be difficult because there is not available enough space. Sravani [15] describes *Portulaca oleracea* L. including broadleaf weeds which have many branches with a height of up to 1 m. The dominating nature of a particular species can cause adverse effects on the ecosystem occupied by the species. Dominance is closely related to invasion. Tjitrosoedirdjo [16], states invasion is a trait that describes the performance of a plant species that becomes dominant and threatens ecosystems, habitats and other species found in a location.

3.2. Important Value Index (IVI)

Based on Table 1, species with the highest Important Value Index (IVI) was *Echinochloa colonum* (L) Link and the lowest were *Leptochloa chinensis* (L.) Nees, *Cyperus sanguinolentus* Vahl, *Portulaca oleracea* L and *Physalis angulata* L. *Echinochloa colonum* (L) Link has the highest Important Value Index (IVI) because it has the most number and it found in almost all plots. It shows the important effect of the *Echinochloa colonum* (L) Link on these fields [17]. whereas *Leptochloa chinensis* (L.) Nees, *Cyperus sanguinolentus* Vahl, *Portulaca oleracea* L., and *Physalis angulata* L. have the lowest Important Value Index (IVI) because they least found among other types of weeds. Solfiyeni [18] stated that the low relative density value of a species indicates that the species has fewer individuals than other weed types, while the relatively low-frequency value of a species is an indication that this type have a narrow distribution.

Table 1. Structure of Rice Weed in Bulusari Village, Sayung Districts, Demak Regency

No.	Type	Famili	Nama spesies	IVI (%)
1	Grasses	Poaceae	<i>Digitaria ciliaris</i> (Retz.) Koel	12.08
2			<i>Echinochloa colonum</i> (L) Link	105.16
3			<i>Leptochloa chinensis</i> (L.) Nees	5.88
5	Sedge	Cyperaceae	<i>Cyperus sanguinolentus</i> Vahl	5.88

6	weed		<i>Cyperus cephalotes</i> Vahl	29.57
7	Broad-	Asteraceae	<i>Eclipta prostrata</i> (L.) L	12.08
8	leaved	Portulacaceae	<i>Portulaca oleracea</i> L.	5.88
9		Solanaceae	<i>Physalis angulata</i> L.	5.88
10		Euphorbiaceae	<i>Euphorbia hypericifolia</i> L.	18.60
Total Number of Individuals of all Types				310
Total number of all types (S)				10
Species Diversity Index (H')				0.69

3.3 Species Diversity Index

Weed species diversity index in 8 weeks old rice was 0.69. This value shows that weed species diversity in 8 weeks old rice was low. This value indicates that the rice field ecosystem has low weed diversity. This condition indicates that the ecosystem is less balanced. Suryatini [19] stated that the value of H' would increase if the number of species in the community increased and the distribution was evener. Species diversity has a close relationship with environmental conditions

3.4 Environmental factor

Based on the measurement of environmental factors conducted at the research location, the results can be seen in table 2 below

Table 2. Measurement of Environmental Factors on Rice Field in Bulusari Village, Sayung Districts, Demak Regency

No.	Environmental Factors	Average
1.	Light intensity	11,562 lux
2.	Air temperature	27 °f
3.	Humidity	86 %
4.	Soil pH	7.2
5.	Altitude	9 m fsl

Environmental factors are the factor which has influenced the diversity of the rice weeds from the outside. Based on Table 2, the light intensity at the research site was 11,562 lux. Utami [20] stated that high light intensity has an impact on growth and yield of good rice plants. Air temperature at the research site is 27°C. According to Balitan [21], the optimum temperature for rice plants to grow is 24-28 °C. Air humidity at the research site was 86%. Uluputty [22], stating that high air humidity is needed by weeds to germinate the soil pH at the location of this research is 7.2. Suryatini [19], stated that the soil pH that is best for growth and availability of nutrients is close to neutral (6.5-7.5). Soil pH determines whether or not nutrients can be absorbed easily by plants. The height of the research area is 9 meters above sea level. Suryatini [19], stated that the height of the place is not a limiting factor for the spread of weeds, because weeds can grow well to a height above 500 m above sea level.

4. Conclusion

From this present research, we conclude that *Echinochloa colonum* (L) Link was commonly found on the rice field in Bulusari Village, Sayung Districts, Demak Regency. It needs advance research to know

fluctuating community structure of the rice weeds in Field in Bulusari Village, Sayung Districts, Demak Regency

References

- [1] Antralina M, Istina I N, Yuwariah Y, Simarmata T 2015 *Procedia Food Science* **3** 323.
- [2] Sarifin M, Sujana I P, Pura N L S 2017 *Agrimeta* **7** 50-55.
- [3] Utami S dan Pudyaningrum, L R 2012 *J Bioma* **2** 91.
- [4] Fitri D S, Syam Z dan Solfiyeni 2014 *J Biologi Universitas Andalas* **3** 68.
- [5] Badan Pusat Statistik Provinsi Jawa Tengah. 2015 *Luas Panen, Produksi dan Produktivitas padi Sawah dan Padi Ladang Menurut Kabupaten dan Kota di Provinsi Jawa Tengah tahun 2015*.
- [6] Martitik D A 2018 *Keragaman Serangga OPT dan Musuh Alami di Lahan Kacang Hijau Sebelum dan Pasca Panen* Skripsi Departemen Biologi Undip
- [7] Soerjani M, Jahja A, Kostermans G H, Tjitrosoepomo G 1987 *Weed of Rice in Indonesia*.
- [8] Johnston and Gillman 1995 *Biodiversity and Conversation* **4** 339.
- [9] Irwanto 2007 *Analisis Vegetasi untuk Pengelolaan Kawasan Hutan Lindung Pulau Marsegu, Kabupaten Seram Barat, Provinsi Maluku* Sekolah Pasca Sarjana, Universitas Gadjah Mada, Yogyakarta.
- [10] Indriyanto 2006 *Ekologi Hutan* Jakarta: Bumi Aksara.
- [11] Brower J E, Zar J H 1997 *Field and Laboratory Methods for General Ecology* Iowa: Brown
- [12] Krebs C J 1989 *Ecological Methodology* Harper and Row Publisher New York
- [13] Fachrul M F 2007 *Metode Sampling Bioteknologi* Jakarta: Bumi Aksara.
- [14] Tanasale V L 2012 *Budidaya Pertanian* **8** 7.
- [15] Sravani V L, Abbas Z, Surya P 2017 *IAJPS* **4** 2845
- [16] Tjitrosoedirdjo S 2015 *Tumbuhan Invasif Pelaihan ke III Pengelolaan Gulma dan Tumbuhan Bogor: Invasif Seameo Biotrop*.
- [17] Amarullah E T, Trizelia, Yaharwandi, Hamid H 2017 *Biodiversitas* **18** 1218.
- [18] Solfiyeni, Chairul, Marpaung M 2016 *Proceeding Biology Education Conference* vol 13 p 743.
- [19] Suryatini Luh 2018 *J Sains dan Teknologi* **7** 77.
- [20] Utami D N 2017 *Pengaruh Intensitas Cahaya Terhadap Pertumbuhan dan Hasil Beberapa Varietas padi (Oryza sativa L.)* Electronic Thesis and Dissertation Unsyah Aceh.
- [21] Balitan 2008 *Teknologi Budidaya Padi* Balai Besar Pengkajian dan Pengembangan Teknologi Pertanian Solok.
- [22] Uluputty M R 2014 *J Agrolgia* **3** 37.

Weeds community structure on the rice field (*Oryza sativa* L.) in bulusari village, Sayung district, Demak regency

ORIGINALITY REPORT

19%

SIMILARITY INDEX

12%

INTERNET SOURCES

10%

PUBLICATIONS

18%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Universitas Diponegoro Student Paper	8%
2	repository.tudelft.nl Internet Source	2%
3	Submitted to Universiti Sains Malaysia Student Paper	2%
4	Submitted to King Mongkut's Institute of Technology Ladkrabang Student Paper	1%
5	Submitted to Universitas Brawijaya Student Paper	1%
6	ediss.uni-goettingen.de Internet Source	1%
7	Tahir Hussain Awan, Pompe C. Sta Cruz, Bhagirath Singh Chauhan. "Effect of pre-emergence herbicides and timing of soil saturation on the control of six major rice weeds and their phytotoxic effects on rice seedlings",	1%

Crop Protection, 2016

Publication

8	Submitted to University of Seoul Student Paper	1%
9	jurnal.untad.ac.id Internet Source	1%
10	china.iopscience.iop.org Internet Source	1%
11	Submitted to Universiti Putra Malaysia Student Paper	<1%
12	www.spotidoc.com Internet Source	<1%
13	Submitted to Sriwijaya University Student Paper	<1%
14	GUI H. LIU. "Landscape-scale variation in the seed banks of floodplain wetlands with contrasting hydrology in China", Freshwater Biology, 10/2006 Publication	<1%
15	Subudhi, H. N., S. P. Panda, P. K. Behera, and Chitra Patnaik. "A Check List of Weeds in Rice Fields of Coastal Orissa, India", Journal of Agricultural Science, 2015. Publication	<1%
16	Chauhan, Bhagirath S., Sharif Ahmed, and Tahir	

H. Awan. "Performance of sequential herbicides in dry-seeded rice in the Philippines", Crop Protection, 2015.

<1%

Publication

Exclude quotes On

Exclude matches < 7 words

Exclude bibliography On

Weeds community structure on the rice field (*Oryza sativa* L.) in bulusari village, Sayung district, Demak regency

GRADEMARK REPORT

FINAL GRADE

/500

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
