



Cauchy Problem for Fractional Ricatti Differential Equations Type with Alpha Order Caputo Fractional Derivatives

Ayad R. Khudair¹, S.A.M. Haddad², Sanaa L. khalaf³

^{1,2,3}Department of Mathematics, Faculty of Science, University of Basrah, Basrah, Iraq.

¹ayadayad1970@yahoo.com, ²shathahaddad@hotmail.com, ³sanaasanaa1978@yahoo.com

Abstract

In this paper, we investigate solution of the fractional Ricatti differential equations (FRDEs) with alpha order Caputo fractional derivatives. In fact, FRDEs are analogous of the Ricatti ordinary differential equations. The multi power series method is used to obtain a useful formula that is implemented to find an explicit solution of Cauchy problem for FRDEs without solving any integral. This formula is explicit and easy to compute by using Maple software to get explicit solution. Also, it is shown that the proposed formula can be used to solve the Cauchy problem for Ricatti ordinary differential equations.

Keywords: Ricatti differential equation; Caputo fractional derivatives; multi power series method; Cauchy problems.

1. Introduction

The history of fractional calculus is back to 1695 when L'Hospital asked Leibniz about $\frac{d^{1/2}u(x)}{dx^{1/2}}$.

Leibniz replied, "It will lead to a paradox." But he added prophetically, "From this apparent paradox, one day useful consequences will be drawn" [5]. In the past, only pure mathematician deal with fractional calculus because they are believed there is no applications to the fractional derivatives [14]. Based on the fact that a reasonable modeling of many physical phenomena having to depend on the time instant to gather with the prior time history, fractional calculus can be used successfully. Therefore, in recent year, fractional derivatives have been used in many phenomena in electromagnetic theory, fluid mechanics, viscoelasticity, circuit theory, control theory, biology, atmospheric physics, etc., [10] and [21]. However, there are two difficulties raised in the study of fractional derivatives, fractional derivatives cannot be expressed to a tangent direction as the classical first derivative. The second difficulty comes from complex integro-differential definitions which make the chain rule not valid for all type of fractional derivatives.

Fractional differential equations (FDEs) have been used to describe many real world problems such as damping laws, fluid mechanics, rheology, physics, mathematical biology, diffusion processes, electrochemistry, and so on. The solvability of a wide fractional differential equation types has been attracted many researchers [4], [6], [7], [11], [12], [16], [17], [19], [20], and [21]. Some theorems related

to existence and uniqueness solutions of initial value problem for FDEs can be found in the books [18] and [10].

Any differential equation that is quadratic in the unknown function is called the Riccati differential equation. This equation has many applications, especially in control theory. In recent year, the analytic or approximate solution of FRDEs has investigated by using many methods. For example, the decomposition method [15], the generalized Haar wavelet [13], Laplace-Adomian-Padé [8], the operator matrix of shifting Legendre polynomial [9], Legendre wavelet operational matrix [3], the generalized differential transform method [2]. Also, FRDEs may be considered as a good test example for investigating the accuracy and effectiveness of the numerical methods as shown in [1]. Motivated and inspired by the on-going research in this field, we will consider the following FRDEs type.

$$\begin{aligned} D^\alpha u(x) &= A(x) + B(x)u(x) + C(x)u^2(x), \quad x \geq 0, \quad m-1 < \alpha \leq m, \quad m \in \mathbb{N} \\ u(0) &= u_0, u'(0) = u_1, \dots, u^{(m-1)}(0) = u_m, \end{aligned} \quad (1)$$

Where $D^\alpha u(x)$ is Caputo derivative, [18] and [10], for any analytic function $u(x)$ which defined by

$$D^\alpha u(x) = \frac{1}{\Gamma(n-\alpha)} \int_0^x \frac{u^{(n)}(t)}{(x-t)^{\alpha-n+1}} dt, \quad n-1 < \alpha < n, \quad n \in \mathbb{Z}^+ \quad (2)$$

Clearly Caputo fractional derivative is linear operator i.e.

$$D^\alpha (a f(x) + b g(x)) = D^\alpha a f(x) + D^\alpha a g(x), \quad a, b \in \mathbb{R} \quad (3)$$

For all $\alpha > 0$ order Caputo derivative, one can have

$$D^\alpha C = 0, \quad \alpha > 0, \quad C, \text{ is a constant} \quad (4)$$

$$D^\alpha x^n = \begin{cases} 0, & \text{for } n \in \mathbb{N}_0 \text{ and } n < [\alpha] \\ \frac{\Gamma(n+1)}{\Gamma(n-\alpha+1)} x^{n-\alpha} & \text{for } n \in \mathbb{N}_0 \text{ and } n \geq [\alpha] \end{cases} \quad (5)$$

Where $[\alpha]$ is the smallest integer greater than or equal to α and $\mathbb{N}_0 = \{0\} \cup \mathbb{N}$. Note that, when $\alpha \in \mathbb{N}$, the Caputo differential derivative is coinciding with the classical differential derivative. For more details on fractional derivatives definitions and their properties, the reader may see the books [18], [10] and [21].

2. Main Result

In this section, we state and prove a new theorem to find a useful explicit formula to compute a solution of Cauchy problems for FRDEs without solving any integral. Now, we will state and prove the following lemma.

Lemma: Let $f(x)$ is a smooth functions for $x \geq 0$, and $u(x) = \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} U_{i,j} x^{i+\alpha j}$ then

$$(1) \quad D^\alpha u(x) = \sum_{i=m}^{\infty} U_{i,0} \frac{\Gamma(i+1)}{\Gamma(i-\alpha+1)} x^{i-\alpha} + \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} U_{i,j+1} \frac{\Gamma(i+\alpha j+\alpha+1)}{\Gamma(i+\alpha j+1)} x^{i+\alpha j}, \quad m-1 < \alpha < m$$

$$(2) \quad f(x)u(x) = \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \left(\sum_{s=0}^i \frac{f^{(s)}(0)}{s!} U_{i-s,j} \right) x^{i+\alpha j}$$

$$(3) \quad f(x)u^2(x) = \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \left(\sum_{s=0}^i \sum_{r=0}^{i-s} \sum_{k=0}^j \frac{f^{(s)}(0)}{s!} U_{r,j-k} U_{i-s-r,k} \right) x^{i+\alpha j}$$

Proof (1): Let us rewrite $u(x)$ as $u(x) = \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} U_{i,j} x^{i+\alpha j} = \sum_{i=0}^{\infty} U_{i,0} x^i + \sum_{i=0}^{\infty} \sum_{j=1}^{\infty} U_{i,j} x^{i+\alpha j}$.

Then we apply D^α to have

$$D^\alpha u(x) = \sum_{i=m}^{\infty} U_{i,0} \frac{\Gamma(i+1)}{\Gamma(i-\alpha+1)} x^{i-\alpha} + \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} U_{i,j+1} \frac{\Gamma(i+\alpha j+\alpha+1)}{\Gamma(i+\alpha j+1)} x^{i+\alpha j}, \quad m-1 < \alpha < m$$

Proof (2): By direct computation, one can have $f(x)u(x) = \left(\sum_{i=0}^{\infty} \frac{f^{(i)}(0)}{i!} x^i \right) \left(\sum_{i=0}^{\infty} \sum_{j=0}^{\infty} U_{i,j} x^{i+\alpha j} \right)$.

Assume $W_i = \sum_{j=0}^{\infty} U_{i,j} x^{\alpha j}$ we have,

$$\begin{aligned} f(x)u(x) &= \left(\sum_{i=0}^{\infty} \frac{f^{(i)}(0)}{i!} x^i \right) \left(\sum_{i=0}^{\infty} W_i x^i \right) \\ &= \sum_{i=0}^{\infty} \sum_{s=0}^i \frac{f^{(s)}(0)}{s!} W_{i-s} x^i \\ &= \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \left(\sum_{s=0}^i \frac{f^{(s)}(0)}{s!} U_{i-s,j} \right) x^{i+\alpha j} \end{aligned}$$

Proof (3): First, we compute $u^2(x)$

$$u^2(x) = \left(\sum_{i=0}^{\infty} \sum_{j=0}^{\infty} U_{i,j} x^{i+\alpha j} \right) \left(\sum_{i=0}^{\infty} \sum_{j=0}^{\infty} U_{i,j} x^{i+\alpha j} \right) = \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \left(\sum_{r=0}^i \sum_{s=0}^j U_{r,j-s} U_{i-r,s} \right) x^{i+\alpha j}$$

Now, by direct computation, one can have

$$f(x)u^2(x) = \left(\sum_{i=0}^{\infty} \frac{f^{(i)}(0)}{i!} x^i \right) \left(\sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \left(\sum_{r=0}^i \sum_{k=0}^j U_{r,j-k} U_{i-r,k} \right) x^{i+\alpha j} \right).$$

Assume $W_i = \sum_{j=0}^{\infty} \left(\sum_{r=0}^i \sum_{k=0}^j U_{r,j-k} U_{i-r,k} \right) x^{\alpha j}$ we have,

$$\begin{aligned} f(x)u^2(x) &= \left(\sum_{i=0}^{\infty} \frac{f^{(i)}(0)}{i!} x^i \right) \left(\sum_{i=0}^{\infty} W_i x^i \right) = \sum_{i=0}^{\infty} \sum_{s=0}^i \frac{f^{(s)}(0)}{s!} \sum_{j=0}^{\infty} \left(\sum_{r=0}^{i-s} \sum_{k=0}^j U_{r,j-k} U_{i-s-r,k} \right) x^{i+\alpha j} \\ &= \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \left(\sum_{s=0}^i \sum_{r=0}^{i-s} \sum_{k=0}^j \frac{f^{(s)}(0)}{s!} U_{r,j-k} U_{i-s-r,k} \right) x^{i+\alpha j} \end{aligned}$$

Theorem: If $A(x), B(x)$ and $C(x)$ are smooth functions for $x \geq 0$, then the Cauchy problems for FRDEs type

Eq.(1) has a solution $u(x) = \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} U_{i,j} x^{i+\alpha j}$ where, $U_{i,j}$ is the solution of the following forward linear

algebraic system

$$U_{i,0} = u^{(i)}(0) = u_i, \quad \forall i = 0, 1, 2, \dots, m-1, \quad (6)$$

$$U_{i,0} = 0, \quad i = m, m+1, m+2, \dots, \quad (7)$$

$$U_{i,1} = \frac{\Gamma(i+1)}{\Gamma(i+\alpha+1)} \left(\frac{A^{(i)}(0)}{i!} + \left(\sum_{s=0}^i \frac{B^{(s)}(0)}{s!} U_{i-s,0} \right) + \left(\sum_{s=0}^i \sum_{r=0}^{i-s} \frac{C^{(s)}(0)}{s!} U_{r,0} U_{i-s-r,0} \right) \right), \quad \forall i = 0, 1, 2, \dots, \quad (8)$$

$$U_{i,j+1} = \frac{\Gamma(i+\alpha j+1)}{\Gamma(i+\alpha j+\alpha+1)} \left(\left(\sum_{s=0}^i \frac{B^{(s)}(0)}{s!} U_{i-s,j} \right) + \left(\sum_{s=0}^i \sum_{r=0}^{i-s} \sum_{k=0}^j \frac{C^{(s)}(0)}{s!} U_{r,j-k} U_{i-s-r,k} \right) \right), \quad (9)$$

$$\forall i = 0, 1, 2, \dots, \quad j = 1, 2, 3, \dots$$

Proof: Let $u(x) = \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} U_{i,j} x^{i+\alpha j}$, be a solution to Eq. (1), where $U_{i,j}$, $i, j = 0, 1, 2, \dots$ are unknown

coefficients. To find these coefficients one can substitute $u(x)$ in Eq. (1) and by using the above Lemma, we get

$$\sum_{i=m}^{\infty} U_{i,0} \frac{\Gamma(i+1)}{\Gamma(i-\alpha+1)} x^{i-\alpha} + \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} U_{i,j+1} \frac{\Gamma(i+\alpha j+\alpha+1)}{\Gamma(i+\alpha j+1)} x^{i+\alpha j} = \sum_{i=0}^{\infty} \frac{A^{(i)}(0)}{i!} x^i$$

$$+ \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \left(\sum_{s=0}^i \frac{B^{(s)}(0)}{s!} U_{i-s,j} \right) x^{i+\alpha j} + \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \left(\sum_{s=0}^i \sum_{r=0}^{i-s} \sum_{k=0}^j \frac{C^{(s)}(0)}{s!} U_{r,j-k} U_{i-s-r,k} \right) x^{i+\alpha j}.$$

Assume $U_{i,0} = 0$, $i = m, m+1, m+2, \dots$, so one can have

$$\sum_{i=0}^{\infty} U_{i,1} \frac{\Gamma(i+\alpha+1)}{\Gamma(i+1)} x^i + \sum_{i=0}^{\infty} \sum_{j=1}^{\infty} U_{i,j+1} \frac{\Gamma(i+\alpha j+\alpha+1)}{\Gamma(i+\alpha j+1)} x^{i+\alpha j} = \sum_{i=0}^{\infty} \frac{A^{(i)}(0)}{i!} x^i$$

$$+ \sum_{i=0}^{\infty} \left(\sum_{s=0}^i \frac{B^{(s)}(0)}{s!} U_{i-s,0} \right) x^i + \sum_{i=0}^{\infty} \sum_{j=1}^{\infty} \left(\sum_{s=0}^i \frac{B^{(s)}(0)}{s!} U_{i-s,j} \right) x^{i+\alpha j} + \sum_{i=0}^{\infty} \left(\sum_{s=0}^i \sum_{r=0}^{i-s} \frac{C^{(s)}(0)}{s!} U_{r,0} U_{i-s-r,0} \right) x^i$$

$$+ \sum_{i=0}^{\infty} \sum_{j=1}^{\infty} \left(\sum_{s=0}^i \sum_{r=0}^{i-s} \sum_{k=0}^j \frac{C^{(s)}(0)}{s!} U_{r,j-k} U_{i-s-r,k} \right) x^{i+\alpha j}$$

$$U_{i,1} = \frac{\Gamma(i+1)}{\Gamma(i+\alpha+1)} \left(\frac{A^{(i)}(0)}{i!} + \left(\sum_{s=0}^i \frac{B^{(s)}(0)}{s!} U_{i-s,0} \right) + \left(\sum_{s=0}^i \sum_{r=0}^{i-s} \frac{C^{(s)}(0)}{s!} U_{r,0} U_{i-s-r,0} \right) \right), \quad \forall i = 0, 1, 2, \dots$$

$$U_{i,j+1} = \frac{\Gamma(i+\alpha j+1)}{\Gamma(i+\alpha j+\alpha+1)} \left(\left(\sum_{s=0}^i \frac{B^{(s)}(0)}{s!} U_{i-s,j} \right) + \left(\sum_{s=0}^i \sum_{r=0}^{i-s} \sum_{k=0}^j \frac{C^{(s)}(0)}{s!} U_{r,j-k} U_{i-s-r,k} \right) \right),$$

$$\forall i = 0, 1, 2, \dots, \quad j = 1, 2, 3, \dots$$

3. Illustrated examples

In this section, we used the formula $u(x) = \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} U_{i,j} x^{i+\alpha j}$ in the main theorem to find an analytic or approximate solution for any value of α , where α is the order of FRDEs. For any integer number, M , we compute the value of the coefficients $U_{i,j}$, $\forall i, j = 0, 1, 2, \dots, M$ by substituting the value of $U_{i,0}$, $\forall i = 0, 1, 2, \dots, M$ in Eq. (8) to compute $U_{i,1}$, $\forall i = 0, 1, 2, \dots, M$. Then, we use Eq. (9), to compute the rest of coefficients value $U_{i,j}$, $i, j = 0, 1, 2, \dots, M$. This procedure is carried out to have $u(x) = \sum_{i=0}^M \sum_{j=0}^M U_{i,j} x^{i+\alpha j}$. In fact, this procedure is not entailed any integration or any complex manipulations.

So that it is a very easy method to find the solution of the Cauchy problems for FRDEs type Eq. (1). The accuracy and effectiveness of the method will depend on the number of M . The forward linear algebraic system Eq. (6) - Eq. (9) is easy to compute by using Maple software or by setting a computer code to get explicit an approximate solution to Eq. (1). Several examples are adopted to illustrate the advantage of the proposed Method.

Example 1. Consider the following nonlinear FRDE

$$D^\alpha u(x) = 1 + 2u(x) - u^2(x), \quad x \geq 0, \quad 0 < \alpha \leq 1, \quad u(0) = 0 \quad (10)$$

Where the exact solution for $\alpha = 1$ is

$$u(x) = 1 + \sqrt{2} \tanh \left(\sqrt{2}t + \frac{1}{2} \log \left(\frac{\sqrt{2}-1}{\sqrt{2}+1} \right) \right) \quad (11)$$

Using $M = 4$ and solve Eq. (6) - Eq. (9) for any value of $0 < \alpha \leq 1$, we find a solution of FRDE Eq. (10) as

$$u(x) = \frac{x^\alpha}{\Gamma(\alpha+1)} + \frac{2x^{2\alpha}}{\Gamma(1+2\alpha)} - \frac{(\Gamma(1+2\alpha) - 4\Gamma(\alpha+1)^2)x^{3\alpha}}{\Gamma(\alpha+1)^2 \Gamma(1+3\alpha)} \\ - \frac{\sqrt{\pi}(1+4\alpha)(\Gamma(1+2\alpha)^2 - 4\Gamma(\alpha+1)^2 \Gamma(1+2\alpha) + 2\Gamma(1+3\alpha)\Gamma(\alpha+1))16^{-\alpha} x^{4\alpha}}{\Gamma\left(\frac{3}{2}+2\alpha\right)\Gamma(\alpha+1)^2 \Gamma(1+2\alpha)^2} + \dots$$

If $M = 8$ and $\alpha = 1$ then the solution of Eq. (10) is

$$u(x) = x + x^2 + \frac{1}{3}x^3 - \frac{1}{3}x^4 - \frac{7}{15}x^5 - \frac{7}{45}x^6 + \frac{53}{315}x^7 + \frac{71}{315}x^8 - \dots$$

This result is agreement with the exact result in Eq. (11). Also If $M = 6$ and $\alpha = \frac{1}{2}$ then the solution of Eq. (10) is

$$u(x) = \frac{2\sqrt{x}}{\sqrt{\pi}} + 2x + \frac{16(\pi-1)x^{3/2}}{3\pi^{3/2}} + \frac{(\pi-4)x^2}{\pi} - \frac{32(3\pi^2+44\pi-32)x^{5/2}}{45\pi^{5/2}} \\ - \frac{1(333\pi^2+284\pi-512)x^3}{36\pi^2} + \dots \\ = 1.128379167\sqrt{x} + 2.x + 2.051213127x^{3/2} - 0.2732395433x^2 - 5.521879267x^{5/2} \\ - 10.32009894x^3 - \dots$$

This result is agreement with the result in, example 6 [1], example 3.2 [15].

Example 2. Consider the following nonlinear FRDE

$$D^\alpha u(x) = 1 - u^2(x), \quad x \geq 0, \quad 0 < \alpha \leq 1, \quad u(0) = 0 \quad (12)$$

Where the exact solution for $\alpha = 1$ is

$$u(x) = \frac{e^{2x} - 1}{e^{2x} + 1} \quad (13)$$

Using $M = 8$ and solve Eq. (6) - Eq. (9) for any value of $0 < \alpha \leq 1$, we find a solution of FRDE Eq. (12) as

$$u(x) = \frac{x^\alpha}{\Gamma(\alpha+1)} - \frac{\Gamma(1+2\alpha)x^{3\alpha}}{\Gamma(1+3\alpha)\Gamma(\alpha+1)^2} + \frac{2\Gamma(1+4\alpha)\Gamma(1+2\alpha)x^{5\alpha}}{\Gamma(1+5\alpha)\Gamma(\alpha+1)^3\Gamma(1+3\alpha)}$$

$$- \frac{2(64)^\alpha \Gamma\left(\frac{3}{2} + 3\alpha\right) \Gamma(1+2\alpha) (\Gamma(1+2\alpha)\Gamma(1+5\alpha) + 4\Gamma(1+4\alpha)\Gamma(1+3\alpha)) x^{7\alpha}}{\sqrt{\pi} (1+6\alpha)\Gamma(1+3\alpha)\Gamma(1+7\alpha)\Gamma(\alpha+1)^4\Gamma(1+5\alpha)} + \dots$$

If $M = 8$ and $\alpha = 1$ then the solution of Eq. (12) is

$$u(x) = x - \frac{1}{3}x^3 + \frac{2}{15}x^5 - \frac{17}{315}x^7 + \dots$$

This result is agreement with the exact result in Eq. (13). Also If $M = 8$ and $\alpha = \frac{1}{2}$ then the solution of Eq. (12) is

$$u(x) = \frac{2\sqrt{x}}{\sqrt{\pi}} - \frac{16x^{3/2}}{3\pi^{3/2}} + \frac{1024x^{5/2}}{45\pi^{5/2}} - \frac{8192x^{7/2}}{75\pi^{7/2}} + \dots$$

$$= 1.128379167\sqrt{x} - 0.9577979845x^{3/2} + 1.300806688x^{5/2} - 1.987486218x^{7/2} + \dots$$

This result is agreement with the result in, example 2 [8].

Example 3. Consider the following nonlinear FRDE

$$D^\alpha u(x) = x^2 + 2u(x) + u^2(x), \quad x \geq 0, \quad 0 < \alpha \leq 1, \quad u(0) = 0 \quad (14)$$

Using $M = 4$ and solve Eq. (6) - Eq. (9) for any value of $0 < \alpha \leq 1$, we find a solution of FRDE Eq. (14) as

$$u(x) = \frac{2x^{2+\alpha}}{\Gamma(3+\alpha)} + \frac{4x^{2+2\alpha}}{\Gamma(3+2\alpha)} + \frac{8x^{2+3\alpha}}{\Gamma(3+3\alpha)} + \frac{4\Gamma(5+2\alpha)x^{4+3\alpha}}{\Gamma(5+3\alpha)\Gamma(3+\alpha)^2} + \frac{16x^{2+4\alpha}}{\Gamma(3+4\alpha)}$$

$$+ \left(\frac{8\Gamma(5+2\alpha)\Gamma(5+3\alpha)}{\Gamma(5+3\alpha)\Gamma(5+4\alpha)\Gamma(3+\alpha)^2} + \frac{16\Gamma(5+3\alpha)}{\Gamma(3+\alpha)\Gamma(5+4\alpha)\Gamma(3+2\alpha)} \right) x^{4+4\alpha} + \dots$$

If $M = 6$ and $\alpha = \frac{1}{4}$ then the solution of Eq. (14) is

$$u(x) = 0.7845423289x^{9/4} + 1.203604445x^{5/2} + 1.808731848x^{11/4} + 0.4089298840x^{19/4} + 2.666666667x^3$$

$$+ 1.776863848x^5 + 3.862362235x^{13/4} + 5.089518943x^{21/4} + 0.3860668648x^{29/4} + 5.502191747x^{7/2}$$

$$+ 12.01899082x^{11/2} + 2.712447261x^{15/2} + \dots$$

This result is agreement with the result in, example 1 [2].

If $M = 6$ and $\alpha = \frac{1}{2}$ then the solution of Eq. (14) is

$$u(x) = 0.6018022226x^{5/2} + 0.6666666667x^3 + 0.6877739684x^{7/2} + 0.1509625991x^{11/2} \\ + 0.6666666667x^4 + 0.4415553048x^6 + 0.6113546389x^{9/2} + 0.8293160657x^{13/2} \\ + 0.06141305036x^{17/2} + 0.5333333333x^5 + 1.254210816x^7 + 0.2812575859x^9 + \dots$$

This result is agreement with the result in, example 2 [2].

If $M = 6$ and $\alpha = \frac{3}{4}$ then the solution of Eq. (14) is

$$u(x) = 0.4521829621x^{11/4} + 0.3438869842x^{7/2} + 0.2271977785x^{17/4} + 0.05094746950x^{25/4} \\ + 0.1333333333x^5 + 0.09465289534x^7 + 0.07063849782x^{23/4} + 0.1091095254x^{31/4} \\ + 0.008269955518x^{39/4} + 0.03420165810x^{13/2} + 0.09834685897x^{17/2} + 0.02330746635x^{21/2} + \dots$$

This result is agreement with the result in, example 3 [2].

Example 4. Consider the following nonlinear FRDE type

$$D^\alpha u(x) = 1 - u^2(x), \quad x \geq 0, \quad 1 < \alpha \leq 2, \quad u(0) = \sqrt{2}, \quad u'(0) = -1 \quad (15)$$

If $M = 3$ and $\alpha = 2$ then the solution of Eq. (15) is

$$u(x) = \sqrt{2} - x + \frac{1}{2}x^2 + \frac{1}{3}\sqrt{2}x^3 - \frac{1}{12}x^4 - \frac{1}{12}\sqrt{2}x^4 - \frac{1}{60}x^5 + \frac{1}{36}\sqrt{2}x^6 - \frac{1}{252}x^7 + \frac{1}{360}x^6 \\ - \frac{1}{90}\sqrt{2}x^7 - \frac{17}{3360}x^8 + \frac{5}{3024}\sqrt{2}x^9 + \dots$$

If $M = 8$ and $\alpha = \frac{3}{2}$ then the solution of Eq. (15) is

$$u(x) = 1.414213562 - 1.x + 0.7522527778x^{3/2} + 0.8510768648x^{5/2} - 0.1719434921x^{7/2} \\ - 0.4714045206x^3 - 0.1250000000x^4 + 0.2121320343x^5 - 0.0250000000x^6 \\ + 0.08797191783x^{9/2} - 0.1558708833x^{11/2} - 0.08436962655x^{13/2} + 0.04040835286x^{15/2} + \dots$$

This result is agreement with the result in, example 3.3 [15].

Example 5. Consider the following nonlinear FRDE type

$$D^\alpha u(x) = \sin(x) + \cos(2x)u(x) + \tan(x)u^2(x), \quad x \geq 0, \quad 1 < \alpha \leq 2, \quad u(0) = 1, \quad u'(0) = 1 \quad (16)$$

If $M = 3$ and $\alpha = 2$ then the solution of Eq. (16) is

$$u(x) = 1 + x + \frac{1}{2}x^3 - \frac{1}{60}x^5 + \frac{1}{30}x^6 - \frac{1}{2520}x^7 + \frac{1}{672}x^8 + \frac{1591}{362880}x^9 + \dots$$

If $M = 3$ and $\alpha = \frac{3}{2}$ then the solution of Eq. (15) is

$$u(x) = 1 + x + \frac{8}{5} \frac{x^{5/2}}{\sqrt{\pi}} - \frac{32}{189} \frac{x^{9/2}}{\sqrt{\pi}} + \frac{1}{8} x^4 + \frac{7}{40} x^5 - \frac{1}{144} x^6 + \frac{64}{3465} \frac{x^{11/2}}{\sqrt{\pi}} + \frac{2176}{45045} \frac{x^{13/2}}{\sqrt{\pi}} \\ + \frac{4096}{45045} \frac{\left(\frac{247}{720} + \frac{64}{25\pi}\right) x^{15/2}}{\sqrt{\pi}} + \dots$$

If $M = 3$ and $\alpha = \sqrt{\pi}$ then the solution of Eq. (16) is

$$u(x) = 1 + x + 0.6604508265x^{\sqrt{\pi}+1} - 0.06113973675x^{3+\sqrt{\pi}} + 0.05330353742x^{1+2\sqrt{\pi}} \\ + 0.07252965997x^{2+2\sqrt{\pi}} - 0.002447974756x^{3+2\sqrt{\pi}} + 0.002282238135x^{1+3\sqrt{\pi}} \\ + 0.005812050616x^{2+3\sqrt{\pi}} + 0.01477733722x^{3+3\sqrt{\pi}} + \dots$$

4. Conclusion

In this paper, we introduce a useful formula to compute a solution of the Cauchy problem for FRDEs type Eq. (1). This formula is not entailed any integration or any complex manipulations even if the FRDEs is content high nonlinearity terms as in the example (5). Also, it is explicit and easy to compute by using Maple software or by setting a computer code to get an explicit solution to this problem. It should be emphasized that this formula can be used to find a solution of high order Ricatti ordinary differential equations. The accuracy and effectiveness of the considering method depend on the number of terms as shown in illustrated examples. For more accuracy and effectiveness, we advise to use Padé approximation as (Khan at el. 2013) and (Momani at el. 2006).

References

- [1] Arikoglu A., Ozkol I. (2007) Solution of fractional differential equations by using differential transform method. *Chaos, Solitons and Fractals* 34, 1473–1481.
- [2] Bansal M. K. and Jain R.,(2015) Application of generalized differential transform method to fractional-order Ricatti differential equation and numerical result . *International Journal of Pure and Applied Mathematics*, 99, 3, 355-366.
- [3] Balaji S., (2015) Legendre wavelet operational matrix method for solution of fractional order Ricatti differential equation. *Journal of the Egyptian Mathematical Society* 23, 263–270.
- [4] Bonilla, B., Rivero, M., Trujillo, J.J. (2007) On systems of linear fractional differential equations with constant coefficients, *Applied Mathematics and Computation* 187(1 SPEC. ISS.), pp. 68-78.
- [5] Debnath L, Bhatta D (2007) *Integral transforms and their applications*. Chapman& Hall, London
- [6] Eidelman SD, Kochubei AN (2004) Cauchy problem for fractional diffusion equations. *J Diffe Equ* 199:211–255.
- [7] Guo S, Mei L, Ling Y, Sun Y (2012) The improved fractional sub-equation method and its applications to the space-time fractional differential equations in fluid mechanics. *Phys Lett A* 376:407–411.
- [8] Khan N. A., Ara A. and Khan N. A.,(2013) Fractional-order Ricatti differential equation: Analytical approximation and numerical results, *Advances in Difference Equations*. 185.
- [9] Krishnaveni K., Kannan K. and Balachandar S. R., (2014) Solving fractional Ricatti differential equation based on operational matrices. *Journal of Computational Methods in Sciences and Engineering* 14 ,229–243.
- [10] Kilbas AA, Srivastava HM, Trujillo JJ (2006) *Theory and applications of fractional differential equations*. Elsevier, Amsterdam.
- [11] Kilbas, A.A., Rivero, M., Rodríguez-Germá, L., Trujillo, J.J, (2007) α -Analytic solutions of some linear fractional differential equations with variable coefficients, *Applied Mathematics and Computation* 187 (1 SPEC. ISS.), pp. 239-249.
- [12] Khudair A. R.,(2013) On solving non-homogeneous fractional differential equations of Euler type. *Computational and Applied Mathematics*, 32, 3, 577-584.

- [13] LI Y., HU L., 2010, Solving fractional Ricatti differential equations using Haar wavelet ,Third International Conference on Information and Computing, IEEE.
- [14] Machado JT, Kiryakova V, Mainardi F (2011) Recent history of fractional calculus. Commun Nonlinear Sci Numer Simul 16(3):1140–1153.
- [15] Momani S., Shawagfeh N. (2006) Decomposition method for solving fractional Ricatti differential equations, Applied Mathematics and Computation 182 1083–1092.
- [16] Molliq RY, Noorani MSM, Hashim I (2009) Variational iteration method for fractional heat- and wave-like equations. Nonlinear Anal Real World Appl 10:1854–1869.
- [17] Mophou GM (2010) Existence and uniqueness of mild solutions to impulsive fractional differential equations. Nonlinear Anal Theory Methods Appl 72:1604–1615.
- [18] Podlubny I (1999) Fractional differential equations. In: An Introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications. Academic Press.
- [19] San Diego Rajeev, Kushwaha MS (2013) Homotopy perturbation method for a limit case Stefan problem governed by fractional diffusion equation. Appl Math Model 37:3589–3599.
- [20] Xue C, Nie J, Tan W (2008) An exact solution of start-up flow for the fractional generalized Burgers' fluid in a porous half-space. Nonlinear Anal Theory Methods Appl 69:2086–2094.
- [21] Yong Zhou (2014) Basic Theory of Fractional Differential Equations, World Scientific Publishing Co. Pte. Ltd.
- [22] Zhukovskaya NV, Kilbas AA (2011) Solving homogeneous fractional differential equations of Euler type. Diff Equ 47(12) 1714–1725.