

Rare Case of Klinefelters Syndrome With 48XXYY

Vasudev G. Magaji MD, MS

Lehigh Valley Health Network, vasudev_g.magaji@lvhn.org


Deborah Feden CRNP

Lehigh Valley Health Network, Deborah.Feden@lvhn.org

Gretchen Perilli MD

Lehigh Valley Health Network, gretchen_a.perilli@lvhn.org

Follow this and additional works at: <http://scholarlyworks.lvhn.org/medicine>

 Part of the [Congenital, Hereditary, and Neonatal Diseases and Abnormalities Commons](#),
[Endocrinology, Diabetes, and Metabolism Commons](#), and the [Medical Sciences Commons](#)

Published In/Presented At

Magaji, V., Feden, D., & Perilli, G. (2014, May 14-18). *Rare case of Klinefelters syndrome with 48XXYY*. Poster presented at: The AACE 23rd Annual Scientific and Clinical Congress, Las Vegas, NV.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Rare Case of Klinefelters Syndrome With 48XXYY

Vasudev Magaji MD, Debra Feden CRNP and Gretchen Perilli, MD

Division of Endocrinology, Department of Medicine, Lehigh Valley Health Network, Allentown, Pennsylvania

OBJECTIVE

Describe Klinefelters Syndrome with 48XXYY.

CASE PRESENTATION

A 28 year old male with klinefelter syndrome (KF) diagnosed during learning disability evaluation at age of 13 presented for follow up. Intriguingly his medical history was very complicated for his young age. He was on testosterone supplementation and his examination revealed tall stature, atrophic testis, gynecomastia. Intriguingly his medical history for his young age was very complicated.

- He had diagnosis of childhood asthma that was poorly controlled until the initiation of immunoglobulin therapy for common variable immunodeficiency.
- He had stroke at age 14
- Atrial septal defect repaired at age of 20
- Pulmonary embolism managed with IVC filter and anticoagulation.
- On examination he weighed 366lbs (166.36kgs), was 77 inches tall and had bilateral small sized measuring ~10cc.
- Underdeveloped right frontal sinus and undeveloped left frontal sinus along with poor dentition had been noted on head imaging **Figure 1**. Imaging of his cervical spine showed loss of cervical lordosis **Figure 2**.

Since his records of KF diagnosis was unavailable and he had multiple complex medical problems laboratory testing to confirm diagnosis karyotyping was done. Karyotyping revealed 48,XXYY that explained many of his diverse medical problems which are typical of Klinefelters 48, XXYY syndrome. Despite his multiple medical problems and learning difficulties he has been working as high school basketball manager.



Figure 1. Craniofacial abnormalities.




Figure 2. Loss of normal cervical lordosis.

DISCUSSION

KF is the most common congenital cause of primary hypogonadism, occurring in 1:1000 males¹. While KF results from extra X chromosome in 80% of cases, in the remaining cases this results from numeric sex chromosome abnormalities (48,XXX, 48,XXYY,49,XXXXY) or 46,XY/47,XXY mosaicism, or structurally abnormal sex chromosomes greater and lesser numbers of X chromosomes². 48, XXYY syndrome, that occurs in 1:18000 males, is usually caused by double non-disjunction during spermatogenesis resulting in extra sex chromosomes can also rarely occur from non-disjunction of sex chromosomes in initial stages of a normal embryo³. The physical features of KF XXYY are similar to KF 47,XXY but with more pronounced phenotypic abnormalities including mild craniofacial dysmorphism, skeletal anomalies such as radioulnar synostosis and clinodactyly, lower cognitive function resulting in developmental delays, congenital cardiac anomalies and asthma manifest at younger age⁵. Autism spectrum disorders have been reported in more than 1/3rd of 48XXYY KF patients⁴. Their behavior abnormalities include hyperactivity, attention problems, impulsivity, aggression, mood instability, autistic-like behaviors, and poor social function. In addition to hypogonadism during adulthood, deep vein thrombosis and type II diabetes, both of which were seen in our patient occur in about 18% of patients with KF 48,XXYY⁵. Behavioural issues can make insulin and anticoagulation management extremely challenging.

CONCLUSION

KF 48, XXYY is a severe form of KF with additional clinical, neurodevelopmental & behavioural manifestation that make the management of KF challenging. Care of these individuals requires a multidisciplinary approach.

References:

1. The Klinefelter syndrome of testicular dysgenesis. Schwartz ID et al Endocrinol Metab Clin North Am. 1991;20(1):153.
2. Chromosomal Variants in Klinefelter Syndrome. Kotzot D et al, Sexual Development 1661-5425/11/0053-0109.
3. Tremor in XYY and XXY men. Boisen E et al, Acta Neurol Scand 1978;58:66-73.
4. Social Deficits in Male Children and Adolescents with Sex Chromosome Aneuploidy: A Comparison of XXY, XYY, and XXYY syndromes. Tartaglia N et al, Res Dev Disabil. 2012 July ; 33(4): 1254-1263.
5. A new look at XXYY syndrome: medical and psychological features. Tartaglia N et al, Am J Med Genet A 2008, 15(12):1509-1522. 146A.

© 2014 Lehigh Valley Health Network

A PASSION FOR BETTER MEDICINE.™

Lehigh Valley
Health Network

610-402-CARE LVHN.org