Lehigh Valley Health Network LVHN Scholarly Works

Patient Care Services / Nursing

Development and Utilization of an Acuity Scale for Oncology Patient Navigation

Raizalie Roman-Rosado BSN, OCN, CMSRN Lehigh Valley Health Network, Raizalie.RomanRosado@lvhn.org

Laura Beaupre BSN, OCN, CBPN-IC Lehigh Valley Health Network, Laura.Beaupre@lvhn.org

Kathy Sevedge RN, MA, AOCN Lehigh Valley Health Network, Kathleen.Sevedge@lvhn.org

Jane Zubia RN, OCN, CBPN I-C Lehigh Valley Health Network, Jane.Zubia@lvhn.org

Martiza Y. Chicas RN, PCCN Lehigh Valley Health Network, Maritza Y.Chicas@lvhn.org

See next page for additional authors

Follow this and additional works at: http://scholarlyworks.lvhn.org/patient-care-services-nursing Part of the <u>Nursing Commons</u>

Published In/Presented At

Roman, R., Beaupre, L., Sevedge, K., Zubia, J., Chicas, M., McHugh, D., Morrone, D. & Kichline, T. (2014, September, 18-21). *Development and Utilization of an Acuity Scale for Oncology Patient Navigation*. Poster presented at the Academy of Oncology Nurse & Patient Navigators Fifth Annual Navigation and Survivorship Conference, Orlando, FL.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Authors

Raizalie Roman-Rosado BSN, OCN, CMSRN; Laura Beaupre BSN, OCN, CBPN-IC; Kathy Sevedge RN, MA, AOCN; Jane Zubia RN, OCN, CBPN I-C; Martiza Y. Chicas RN, PCCN; Diane McHugh BSN, OCN, CBPN-IC; Dorothy Morrone RNC, MS, OCN; and Tracy Kichline

Development and Utilization of an Acuity Scale for Oncology Patient Navigation Raizalie Roman RN OCN CMSRN, Laura Beaupre RN OCN CBPN-IC, Kathleen Sevedge RN MA AOCN, Jane Zubia RN OCN CBPN-IC, Maritza Chicas RN OCN PCCN, Diane McHugh RN OCN CBPN-IC, Dorothy

Background and Purpose:

The oncology navigation team at LVHN developed a Needs Assessment and Acuity Scale to help in assessing newly referred patients. The tools are used to determine which components of care can be addressed by other internal cancer support resources, which patients are appropriate for navigation, and the level of navigation needed.

During the quarter from March 1, 2014-May 31, 2014, 4% of patients had an acuity score of 0, 21% had an acuity score of 1, 24% had an acuity score of 2, 27% had an acuity score of 3, and 23% of patient had an acuity score of 4.


- 0 = Not appropriate for Navigation ie: no diagnosis of cancer, not incurrent treatment.
- 1 = Stable navigation needs, no active navigation needed at present.
- 2 = Low acuity for navitation
- 3 = Moderate acuity for navigation
- 4 = High acuity for navigation


Conclusion:

By utilizing the needs assessment and acuity scale, the navigators are able to determine if patients are appropriate for navigation, have stable needs, or require active navigation. The intensity of navigation is also indicated. In the future, we hope to include the acuity measurements to help determine workload.

Morrone RN MS OCN, Tracy Kichline - Data Systems Analyst Lehigh Valley Health Network, Allentown, PA

Acuity Scale Pilot





Navigator Needs Assessment

Needs Assessment	
none() uninsured/MA pending/Prescription	assistance
cial coordinator/datecomment	_
none() oblems Transportation needs conment ld/caregiver Lack of social on-citizen support eds No job/income ds Diff. paying bills	
Worker/date comment none()	
y/on psych meds	
ing depression/ anxiety uality help needed currently/ history Prescription meds()) Recreational drugs() ubstance Abuse/date comment	
None ()	
wing	
ian/ Speech Therapist/ Date	
)	





A PASSION FOR BETTER MEDICINE.

610-402-CARE LVHN.org