### Lehigh Valley Health Network LVHN Scholarly Works

Department of Medicine

### Outcomes of Peginterferon alfa-2a and Ribavirin Combination Therapy in a Resident-Initiated, Multidisciplinary, Hepatitis C Clinic

Nicole M. Agostino DO Lehigh Valley Health Network, Nicole M. Agostino@lvhn.org

Erini Vasiliadis DO Lehigh Valley Health Network

K Nadeem Ahmed MD Lehigh Valley Health Network

Suzanne J. Templer DO Lehigh Valley Health Network, Suzanne \_J.Templer@lvhn.org

Edward R. Norris MD, FAPA, FAPM Lehigh Valley Health Network, edward.norris@lvhn.org

### See next page for additional authors

Follow this and additional works at: http://scholarlyworks.lvhn.org/medicine

Part of the <u>Chemicals and Drugs Commons</u>, <u>Diseases Commons</u>, <u>Gastroenterology Commons</u>, <u>Hematology Commons</u>, <u>Infectious Disease Commons</u>, <u>Medical Sciences Commons</u>, <u>Oncology</u> <u>Commons</u>, <u>Psychiatry Commons</u>, <u>Psychiatry and Psychology Commons</u>, <u>and the Therapeutics</u> <u>Commons</u>

### Published In/Presented At

Agostino, N., Vasiliadis, E., Ahmed, K., Templer, S., Norris, E., Brooks, C., Gertner, E., & Yozviak, J. (2007, November 7). *Outcomes of peginterferon alfa-2a and ribavirin combination Therapy in a resident-Initiated, multidisciplinary, hepatitis c clinic*. Poster presented at: The Pennsylvania College of Internal Medicine Meeting, Reading, PA.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

### Authors

Nicole M. Agostino DO; Erini Vasiliadis DO; K Nadeem Ahmed MD; Suzanne J. Templer DO; Edward R. Norris MD, FAPA, FAPM; Charles M. Brooks MD; Eric J. Gertner MD, MPH; and Joseph L. Yozviak DO, FACP

# **Outcomes of Peginterferon alfa-2a and Ribavirin Combination** Therapy in a Resident-Initiated, Multidisciplinary, Hepatitis C Clinic

Nicole Agostino DO, Erini Vasiliadis DO, K. Nadeem Ahmed MD, Suzanne J. Templer DO, Edward R. Norris MD, Charles M. Brooks MD, Eric J. Gertner MD, MPH, and Joseph L. Yozviak DO Lehigh Valley Hospital and Health Network (LVH), Allentown, Pennsylvania, U.S.A.

# Introduction

**Clinical trials have shown that at least 40% of patients with chronic** hepatitis C (HCV) are able to achieve sustained virologic response (SVR) with peginterferon-alfa/ribavirin (PegIFN/RBV) therapy along with adequate medical and psychiatric support. Those with a history of mental illness or multiple medical comorbidities are often excluded from clinical trials<sup>1,2</sup>. Hence, SVR is achieved less often in clinical practice. We proposed that an integrated, multidisciplinary approach to PegIFN/ **RBV** therapy would result in SVR rates similar to those of clinical trials in a medically and psychiatrically complex cohort of patients.

## Methods:

A pilot Hepatitis C Clinic was established in 2004, staffed by internal medicine residents, an attending gastroenterologist and psychiatrist, and a registered-nurse coordinator. All patients were underinsured/uninsured and underwent treatment for HCV with PegIFN/RBV combination therapy according to evidence-based protocols within the confines of managed-care formularies. Charts were retrospectively abstracted for demographics and baseline characteristics, virologic response, side effects, and reasons for discontinuation or non-initiation of therapy.

# Results:

Forty-eight patients were evaluated. None were co-infected with human immunodeficiency virus (HIV). Twenty-two (46%) were not treated, and twenty-six (54%) were treated. Of the patients who were treated, twentytwo (84%) were genotype 1, three (12%) were genotype 2b, and one (3%) was genotype 4a.

### Table 1: Demographic data for all patients

Gender	Male 15 Female 33
Average Age	45.0 year
Average Weight of Treated Patients	191.3 lbs
<b>Baseline Viral Load</b>	> 400,000 IU 2 < 400,000 IU 1 Not Available
Pre-existing Psychiatric Diagnosis	38 (79%)

### Table 2: Reasons not to treat with PegIFN/RBV

Personal Choice	6 (27%)
Unstable Psychiatric Illness	6 (27%)
Comorbidities	4 (18%)
Substance Abuse	2 (9%)
Viral Clearance	2 (9%)
Lack of follow up	1 (4%)
Minimal Fibrosis	1 (4%)

(31%) (69%)

9 (60%) 6 (33%) 3 (6%)

### Table 3: Outcomes of PegIFN/RBV n=26

Nonresponders (NR)	5 (19%)
Early Viralogic Response (EVR)	13 (50%)
End of Treatment Response (ETR)	12 (46%)
Sustained Viralogic Response (SVR)	8/23 (34%)*
Relapse after ETR	2 (7%)
Dropouts	8 (30%)

\* Three patients were omitted from the total number for evaluation of SVR. Two patients had not reached 24 weeks beyond the end of treatment. One patient was lost to follow up.

### Table 4: Side Effects

Hyperuricemia	12 (46%)
Fatigue	10 (38%)
Myalgia	8 (31%)
Anemia	8 (31%)
Arthralgia	5 (19%)
Insomnia	5 (19%)
Neutropenia	5 (19%)

### **References:**

- Treatment of Chronic Hepatitis C: A Randomised Trial. Lancet 2001;358:958-65.

### Table 5: Reasons for **Discontinuation of PegIFN/RBV**

LEHIGH VALLEY HOSPITAL

Side Effects	6 (23%)
Viralogic Nonresponse	5 (19%)
Psychiatric Side Effects	1 (3%)
Poor Adherence	1 (3%)
Ongoing Substance Abuse	1 (3%)
Incarceration	1 (3%)

# Conclusions

ANCC MAGNET RECOGNITION

**Despite a patient population with mostly** genotype 1 HCV and significant medical/ psychiatric comorbidities, a similar number of patients achieved ETR/SVR as compared to those in clinical trials<sup>1,2</sup>. These data suggest that an integrative medicine practice can safely and effectively manage PegIFN/RBV therapy and serve as a model to expand access to antiviral therapy for many individuals with chronic HCV. Additionally, by incorporating a dedicated group of internal medicine residents, the pool of capable and willing providers of **PegIFN/RBV** therapy will be expanded.

1. MW Fried, ML Schiffman, KR Reddy, et al. Peginterferon alfa-2a Plus Ribavirin for Chronic Hepatitis C Virus Infection. NEJM 2002;347:975-82. 2. MP Manns, JG McHutchison, SC Gordon, et al. Peginterferon alfa-2b Plus Ribavirin Compared with Interferon alfa-2b Plus Ribavirin for Initial