

A Case of Three Patients with Severe Hypothermia Rewarmed with ECMO

Matthew Niehaus DO

Lehigh Valley, Matthew.Niehaus@lvhn.org

Rita Pechulis MD, FCCP

Lehigh Valley Health Network, Rita_M.Pechulis@lvhn.org

James K. Wu MD

Lehigh Valley Health Network, james.wu@lvhn.org

Follow this and additional works at: <http://scholarlyworks.lvhn.org/emergency-medicine>



Part of the [Emergency Medicine Commons](#), and the [Surgery Commons](#)

Published In/Presented At

Niehaus, M., Pechulis, R., Wu, J. (2015, September 18). *A Case of Series of Three Patients with Severe Hypothermia Rewarmed with ECMO*. Poster presented at Extracorporeal Life Support Organization Conference, Atlanta, GA.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

A Case Series of Three Patients with Severe Hypothermia Rewarmed with ECMO

Matthew Niehaus, DO; Rita Pechulis, MD; James Wu, MD
Lehigh Valley Health Network, Allentown, Pennsylvania

Introduction

Severe accidental hypothermia, defined as a core body temperature $<28^{\circ}\text{C}$, is associated with multiple neurologic, cardiac, and metabolic complications.¹ Here we report outcomes of three patients with severe hypothermia and cardiovascular collapse rewarmed and supported on ECMO.

Title			
	PATIENT 1	PATIENT 2	PATIENT 3
Etiology of Hypothermia	Intoxicated, found in woods	Intoxicated, found in snow bank	Heroin overdose, found outside his house
Gender	F	M	M
Age	25 yo	26 yo	21 yo
Initial Temperature	26°C	21.7°C	27.2°C
Initial Presentation	Cardiac arrest	Cardiac arrest	Refractory shock
Rhythm in ED	Ventricular fibrillation	Asystole	Atrial fibrillation
Total Time of CPR	37 min	265 min.	NA
Type of ECMO	VA	VA	VV
Cannula Site	Single Lumen Fem-Fem	Single Lumen Fem-Fem	Dual Lumen R U
Time to Target Temperature	Attained 36°C in 30 min	Attained 28.6°C in 46 min	Attained 37°C in 360 min
Total Time on ECMO	10 hours	36 hours	120 hours
Complications	<ul style="list-style-type: none"> Rhabdomyolysis Bilateral transmetatarsal amputations 	<ul style="list-style-type: none"> Acute renal failures Transmetatarsal/finger amputations Tracheostomy PEG tube placement 	<ul style="list-style-type: none"> Aspiration pneumonia ARDS Tracheostomy PEG tube placement
Length of Stay	37 days	57 days	45 days
Outcome	Neurologically intact	Neurologically intact	Neurologically intact

Discussion

Rewarming therapies are classified as either passive (warm blankets, overhead heaters) or active (infusion of warm fluids, lavage of the peritoneal and thoracic cavities),² with active methods recommended for severe hypothermia. In hypothermic cardiac arrest, the time to regain normal body temperature is the most influential parameter for a good recovery.¹

The most common cause of death from severe hypothermia is cardiac arrest, resulting from slowing of conduction velocity in the myocardium and subsequent ventricular fibrillation (VF).² The myocardium will quickly recover normal, organized electrical activity when rewarmed; consequently, aggressive warming techniques should accompany standard CPR in hypothermic cardiac arrest. While body temperature can be raised $2\text{-}4^{\circ}\text{C}$ using warm saline infusion, increases as high as 12°C per hour have been noted with the use of extracorporeal membrane oxygenation (ECMO).^{1,3}

In hypothermic cardiac arrest, the time to regain normal body temperature is the most influential parameter for a good recovery.¹ A recent study found an association between ECMO in CPR and higher survival rates with good neurologic outcomes.⁴

In two of our cases, the use veno-arterial of ECMO rapidly rewarmed our patient's core body temperature and provided hemodynamic support in the peri-arrest period. In our third case, veno-venous ECMO was used to rewarm our patient and prevent continued cardiac deterioration.

ECMO should be considered early on as a primary means for rewarming in patients who present with severe accidental hypothermia.

References:

- Sawamoto K, Bird SB, Katayama Y, et al. Outcome from severe accidental hypothermia with cardiac arrest resuscitated with extracorporeal cardiopulmonary resuscitation. *Am J Emerg Med* 2014;32(4):320-
- American College of Surgeons. *Advanced Trauma Life Support: Student Course Manual (9th ed.)* 2012. Chicago, IL: Author.
- Freude T, Gillen S, Ehnert S, et al. Therapeutic peritoneal lavage with warm saline solution as an option for a critical hypothermic trauma patient. *Wein Klin Wochenschr* 2014;126(1-2):56-61.
- Stub D, Bernard S, Pellegrino V, et al. Refractory cardiac arrest treated with mechanical CPR, hypothermia, ECMO and early reperfusion (the CHEER trial). *Resuscitation* 2015; 86:88-94.

© 2015 Lehigh Valley Health Network

A PASSION FOR BETTER MEDICINE.™

610-402-CARE LVHN.org