

# A Retrospective Analysis of the Outcomes of Extracorporeal Cardiopulmonary Resuscitation (ECPR).

Rachel Wills

Kennedy Gallagher

James A. Burke MD, PhD

*Lehigh Valley Health Network, James\_A.Burke@lvhn.org*

Tim S. Misselbeck MD

*Lehigh Valley Health Network, Timothy\_S.Misselbeck@lvhn.org*

James K. Wu MD

*Lehigh Valley Health Network, james.wu@lvhn.org*

Follow this and additional works at: <https://scholarlyworks.lvhn.org/research-scholars-posters>

---

## Published In/Presented At

Wills, R., Gallagher, K., Burke, J., Misselbeck, T., Wu, J., (2017, July, 31) *A Retrospective Analysis of the Outcomes of Extracorporeal Cardiopulmonary Resuscitation (ECPR)*. Poster presented at LVHN Research Scholar Program Poster Session, Lehigh Valley Health Network, Allentown, PA.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact [LibraryServices@lvhn.org](mailto:LibraryServices@lvhn.org).

# A Retrospective Analysis of the Outcomes of Extracorporeal Cardiopulmonary Resuscitation (ECPR)

Rachel Wills<sup>1</sup>; Kennedy Gallagher<sup>1</sup>; James Burke, MD<sup>2</sup>; Timothy S. Misselbeck, MD<sup>1</sup>; James Wu, MD<sup>1</sup>

<sup>1</sup>Division of Cardiothoracic Surgery, <sup>2</sup>Division of Cardiology  
Lehigh Valley Health Network, Allentown, Pennsylvania

## BACKGROUND

- Extracorporeal cardiopulmonary resuscitation (ECPR) is a form of veno-arterial extracorporeal membrane oxygenation (VA-ECMO) wherein the support is administered during cardiac arrest to externally sustain circulation <sup>1</sup>
- Once it is determined that a patient undergoing cardiac arrest has the potential for recovery and eventual weaning, ECPR is utilized in attempt to increase survival to discharge, which is otherwise low with traditional cardiopulmonary resuscitation (CPR) <sup>2</sup>
- ECPR is a temporary treatment to allow for a duration of cardiopulmonary rest due to anticipated system healing with time, yet is not a curative measure

## OBJECTIVE

- As a single-center, retrospective chart analysis, this study aims to determine the overall outcomes of patients who underwent ECPR at Lehigh Valley Health Network from 2013-2017

## METHODS

- 1 • Creation of a database in Microsoft Excel to collect demographic and outcome information of patients
- 2 • Categorization of patients undergoing VA ECMO and exclusion of those not for the purpose of ECPR
- 3 • Analysis of 45 ECPR patient outcomes and correlational study with date-based cohorts
- 4 • Consultation and comparison with previous studies to find support or refutation for findings

## OUTCOMES

Occurrences of VA ECMO (102)    Thorough Information Available (92)    Occurrences of ECPR for analysis (45)




Figure 1: Good outcomes, shown in blue, were defined as survival at the time of discharge with no consideration for post-discharge outcomes, and thus all not included within this definition were defined as bad outcomes, shown in red. The distribution of overall outcomes of ECPR patients demonstrates 29% survival to discharge.




Figure 2: All patients who underwent ECPR were split into either one of two cohorts based upon the date of ECPR initiation, with 23 patients in the first and 22 in the second. The distribution of Cohort 1 indicated 22% survival to discharge, whereas Cohort 2 reported 36% survival to discharge.




Figure 3: Disposition at discharge of patients were reported based upon the number of days ECPR support was needed prior to weaning, regardless of weaning success. All patients were weaned from ECPR or care was withdrawn within 8 days of their administration with the exception of two cases.

## RESULTS

- With 29% (95%CI 16-42%) overall survival outcomes, LVHN is consistent with published survival rates <sup>2 3</sup>
- There was a 14% increase in survival to discharge between the cohorts (Cohort 1: 2013-2015, Cohort 2: 2016-2017)
- All ECPR patients had been weaned by or on day 8 with the exception of two patients on day 24 and 26, which were converted to veno-venous (VV) ECMO on day 8 and 2, respectively
- No correlation could be seen between days on ECPR and survival to discharge

## CONCLUSIONS

- Though outcomes for ECPR patients are poor, the difference between cohorts shows that improvement is achievable
- Based on current duration of support data, patients should continue to be weaned, converted, or withdrawn from care by day 8 to avoid unnecessarily long durations
- Future studies should conduct correlational analyses between demographic or treatment based variables of ECPR and patient outcomes

## References

- 1 Abrams, D., & Brodie, D. (2015). Novel uses of extracorporeal membrane oxygenation in adults. *Clinics in Chest Medicine*, 36(3), 373-384. doi: 10.1016/j.ccm.2015.05.014
- 2 Chen, Y.S., Chao, A., Yu, H.Y., Ko, W.J., Wu, I.H., Chen, J.C., Huang, S.H.,... Wang, S.S. (2003). Analysis and results of prolonged resuscitation in cardiac arrest patients rescued by extracorporeal membrane oxygenation. *Journal of the American College of Cardiology*, 41(2), 197-203. https://doi.org/10.1016/S0735-1097(02)02716-X
- 3 Richardson, A., Schmidt, M., Bailey, M., Pellegrino, V., Rycus, P., & Pilcher, D. (2017). ECMO cardio-Pulmonary resuscitation (ECPR), trends in survival from an international multicentre cohort study over 12-years. *Resuscitation*, 112, 34-40. doi:10.1016/j.resuscitation.2016.12.009

© 2017 Lehigh Valley Health Network

610-402-CARE LVHN.org