#### **Lehigh Valley Health Network LVHN Scholarly Works**

Department of Medicine

## Impact of Atrial Fibrillation on Outcomes in Patients Hospitalized with ST-Segment Elevation Myocardial Infarction (Poster).

Lohit Garg MD Lehigh Valley Health Network, lohit.garg@lvhn.org

Sahil Agrawal MD

Manyoo Agarwal MD

Aakash Garg MD

Amitoj Singh

See next page for additional authors

Follow this and additional works at: https://scholarlyworks.lvhn.org/medicine



**Part of the** Cardiology Commons, and the Medical Sciences Commons

#### Published In/Presented At

Garg, L., Agrawal, S., Agarwal, M., Garg, A., Singh, A., Bhatia, N., & ... Cox, D. (2017, March 19). Impact of Atrial Fibrillation on Outcomes in Patients Hospitalized with ST-Segment Elevation Myocardial Infarction (Poster). Poster Presented at: American College of Cardiology Annual Conference 2017, Washington D.C.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Authors Lohit Garg MD; Sahil Agrawal MD; Manyoo Agarwal MD; Aakash Garg MD; Amitoj Singh; Nirmanmoh Bhatia; Mahek Shah MD; Brijesh Patel MD; Sudip Nanda MD, FACP; and David A. Cox MD						

# Impact of Atrial Fibrillation on Outcomes in Patients Hospitalized with ST-Segment Elevation Myocardial Infarction

Lohit Garg, Sahil Agrawal, Manyoo Agarwal, Aakash Garg, Amitoj Singh, Nirmanmoh Bhatia, Mahek Shah, Brijesh Patel, Sudip Nanda, David Cox Lehigh Valley Health Network, Allentown, Pennsylvania



- Atrial fibrillation (AF) is the most common persistent cardiac arrhythmia with an increasing incidence in the United States (US)
- AF is present in 12-15% patients presenting with STEMI
- Previous reports have suggested that STEMI patients who have AF have worse outcomes.

### OBJECTIVE

 We sought to determine the impact of AF on in-hospital outcomes of patients admitted with STEMI in the United States

## METHODS

- Data were obtained from the Nationwide Inpatient Sample database for years 2003- 2013
- ICD 9 codes were used to identify patients with STEMI and AF
- Baseline demographic and clinical features were studied and compared between groups of patients with and without AF.
- The primary outcome of interest was all-cause in-hospital mortality.
- Utilization of percutaneous coronary intervention (PCI) and coronary artery surgery (CABG), and the incidences of complications (periprocedural hemorrhage, gastrointestinal bleeding, blood transfusion, and acute ischemic stroke) were also compared in the 2 groups.
- Results are expressed as frequencies (%) for categorical variables and mean ± SD for continuous variables.
- Differences between groups were analyzed using the Student's t test for continuous variables and the chi-square test for categorical variables.
- Statistical analyses were performed using IBM SPSS, Statistics, version 20.0 (IBM Corp., Armonk, New York).

#### RESULTS

- Of the total 2,632,447 STEMI hospitalizations, AF was documented in 339,987 (12.9%) patients
- At baseline, patients with AF were older (mean 74 vs 63 years, p<0.001) and more likely to be female (42% vs 34%, p<0.001) or white (85% vs78%; p<0.001). (Table 1)</li>
- Hypertension, chronic renal failure, diabetes mellitus and congestive heart failure were all more prevalent among those with AF (p<0.001 for all). (Table 1)
- AF patients were less likely to undergo coronary angiography or any coronary revascularization than non-AF patients. They were more likely to undergo CABG but less likely to undergo PCI.
- Patients with AF had higher risk adjusted in- hospital mortality (16.3% vs 7.9%; OR: 1.16; Cl: 1.15-1.18; p<0.001). and an increased risk of bleeding complications after both PCI (12.2% vs 5.3%; OR: 1.18; Cl: 1.16-1.21; p<0.001) and CABG (40.0% VS 33.5%; OR: 1.11; Cl: 1.08-1.13; p<0.001). (Table 2)</li>
- Average length of stay was also longer among those with AF (mean of 7 days vs 4 days; p<0.001).</li>

Table 1. Baseline Characteristics					
Characteristics	AF (339,988)	Non-AF (2,292,259)			
Age	74.4 +/- 12.5 years	63.6 +/- 14.3 years			
Male	196,455 (57.8%)	1,506,360 (65.7%)			
White	228,017 (85.3%)	1,391,804 (77.9%)			
African American	12,680 (4.7%)	146,865 (8.2%)			
Insurance - Medicare	247,890 (73%)	102,4303 (44.8%)			
Insurance - Medicaid	10,715 (3.2%)	143,467 (6.3%)			
Private	63,246 (18.6%)	837,280 (36.6%)			
Obesity	25,089 (7.4%)	222,451 (9.8%)			
Hypertension	199,029 (59.1%)	1,312,941 (57.7%)			
Diabetes Mellitus	93,338 (27.7%)	613,886 (27%)			
Dyslipidemia	131,068 (38.6%)	1,162,491 (50.7%)			
Current or Former Smoker	71,638 (21.1%)	874,986 (38.2%)			
Prior Myocardial Infarction	25,685 (7.6%)	167,260 (7.3%)			
Prior PCI	25,216 (7.4%)	206,090 (9%)			
Prior CABG	17,697 (5.2%)	91,920 (4%)			
Chronic Kidney Disease	51,047 (15.2%)	177,995 (7.8%)			
Chronic Lung Disease	73,306 (21.8%)	355,026 (15.6%)			
History of Cerebrovascular Accident	10,529 (3.1%)	39,254 (1.7%)			
Peripheral Vascular Disease	32,878 (9.8%)	160,974 (7.1%)			

Table 2. Outcomes					
Treatment and Outcomes	AF	Non-AF	Adjusted Odds Ratio	P-value	
Coronary Angiograophy	58.2 %	72.7 %	1.00 (0.98-1.01)	.50	
Percutaneous Coronary Intervention	41 %	61 %	0.92 (0.91-0.93)	<0.001	
Coronary Artery Bypass Grafting	15%	7%	2.37 (2.34-2.40)	<0.001	
All-cause In-hospital Mortality	16.3%	7.9 %	1.16 (1.15-1.18)	<0.001	
Cardiogenic Shock	14.2%	7.6%	1.27 (1.25-1.28)	<0.001	
Bleeding Complications (PCI)*	12.2%	5.3%	1.18 (1.16-1.21)	<0.001	
Bleeding Complications (CABG)	40.0%	33.5%	1.11 (1.08-1.13)	<0.001	
Acute Ischemic Stroke	1.7%	0.8%	1.4 (1.35-1.44)	<0.001	

\* GI bleed, retroperitoneal hematoma, transfusion or acute anemia.

### LIMITATIONS

- Retrospective design of the study and possibility of varying coding practices among hospitals.
- Inability to differentiate paroxysmal from persistent or permanent AF.
- Medication related data are unavailable due to administrative nature of the database

#### CONCLUSIONS

- AF is common among patients presenting with STEMI.
- AF is associated with older age, and higher prevalence of comorbidities at admission in STEMI patients.
- AF is independently associated with higher in-hospital mortality, and periprocedural complications in STEMI patients.

**Disclosures:** Authors have no conflict of interest.

© 2017 Lehigh Valley Health Network

610-402-CARE LVHN.org

