

Impact of sarcopenia on outcomes of locally advanced esophageal cancer patients treated with neoadjuvant chemoradiation followed by surgery

Gilbert Z. Murimwa

USF MCOM- LVHN Campus, Gilbert.Murimwa@lvhn.org

P S. Venkat

W Jin

USF MCOM- LVHN Campus

S Leuthold

K Latifi

See next page for additional authors

Follow this and additional works at: <https://scholarlyworks.lvhn.org/select-program>

Part of the [Medical Education Commons](#)

Published In/Presented At

Murimwa, G. Z., Venkat, P. S., Jin, W. Leuthold, S. Latifi, K. Almhanna, K. Pimiento, J. M., Fontaine, J. P., Hoffe, S. E., Frakes, J. M. (2019, March). *Impact of sarcopenia on outcomes of locally advanced esophageal cancer patients treated with neoadjuvant chemoradiation followed by surgery*. Poster Presented at: 2019 SELECT Capstone Posters and Presentations Day. Kasych Family Pavilion, Lehigh Valley Health Network, Allentown, PA

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Authors

Gilbert Z. Murimwa, P S. Venkat, W Jin, S Leuthold, K Latifi, K Almhanna, J M. Pimiento, J P. Fontaine, Sarah E. Hoffe, and J M. Frakes

Impact of sarcopenia on outcomes of locally advanced esophageal cancer patients treated with neoadjuvant chemoradiation followed by surgery

G. Z. Murimwa¹, P. S. Venkat², W. Jin¹, S. Leuthold³, K. Latifi², K. Almhanna⁴, J. M. Pimiento⁴, J. P. Fontaine⁴, S. E. Hoffe², and J. M. Frakes²

¹University of South Florida Morsani College of Medicine, Tampa, FL, ²H. Lee Moffitt Cancer Center and Research Institute, Department of Radiation Oncology, Tampa, FL, ³University of South Florida, Tampa, FL, United States ⁴H. Lee Moffitt Cancer Center and Research Institute, Tampa, FL

Lehigh Valley Health Network, Allentown, Pennsylvania

INTRODUCTION

- Sarcopenia is the progressive and generalized loss of skeletal muscle
- Its significance is increasingly recognized and reported within the surgical and medical oncology fields
- It is now reported as an independent predictor of clinical outcomes in multiple gastrointestinal cancers
- Sarcopenia is being explored in surgical and medical oncology as a prognostic factor before treatment
- This study evaluated whether sarcopenia could be prognostic for grade 3 or greater toxicity, pathologic response, or overall survival in patients treated neoadjuvantly for esophageal cancer

METHODS

- Moffitt esophageal cancer patients treated with neoadjuvant chemoradiation followed by surgery from 2008-2012
- All patients received IMRT/IGRT utilizing dose painting to a total dose of 50.4/56 Gy in 28 fractions along with cisplatin and 5FU chemotherapy
- Sarcopenia was defined as the presence of a psoas area less than the median of the cohort
- ROC curve, logistic regression, chi square and Kaplan Meier estimates were used when appropriate

Eligibility Criteria

- CT imaging for planning including the L4 vertebral body
- 56 patients were included


RESULTS

Table 1

		Total Psoas Area		
		<1373.6	>1373.6	
		No. of patients (%)		P
Total Patients		23	33	
Median age (range)		64 (47-81)	62 (30-81)	
Median Follow-up (months)		56.73	65.37	
Gender	Male	14 (25%)	33 (59%)	<0.001
	Female	9 (16%)	0	
Pathologic Complete Response	Not pathCR	11 (20%)	17 (30%)	0.786
	pathCR	12 (21%)	16 (29%)	
Favorable Pathologic Response	TRG 2/3	3 (5%)	7 (13%)	0.432
	TRG 0/1	20 (36%)	26 (46%)	
Acute Toxicity	Grade ≤2	8 (14%)	25 (45%)	0.002
	Grade 3+	15 (27%)	8 (14%)	

T1. Patient Characteristics

Figure 1



F1. ROC Analysis of Grade 3+ Toxicity


- Sarcopenia was predictive of
 - Grade 3 Toxicity
 - The smaller the psoas cross sectional area, the higher the chance of any grade 3 toxicity
 - Patients below our cutoff were 5.86 times more likely to develop a grade 3 or higher toxicity (p=0.003)
- Sarcopenia was not predictive of
 - Pathologic Response
 - Overall Survival (p=0.124)

Table 2

	Sarcopenic	Non-sarcopenic
Total Patients	23	33
Patients with G3 Toxicity	15	8
Dysphagia requiring feeding tube	10	6
Neutropenia	3	1
Hospitalization	2	0
Radiation Pneumonitis	0	1

T2. Toxicity

Figure 2



F2. Kaplan Meier Curve for Overall Survival

CONCLUSIONS / FUTURE DIRECTIONS

Sarcopenia may be a useful prognostic marker for radiation therapy, especially in esophageal cancer

- Risk Stratification
- Supportive/Nutritional Management
- Dose Escalation
- Quality of life benefit
- Further Validation

REFERENCES

- Harada K, Ida S, Baba Y, et al. Prognostic and clinical impact of sarcopenia in esophageal squamous cell carcinoma. *Dis Esophagus*. 2016;29(6):627-33.
- Cintosun U, Altun B, Tasci I. Sarcopenia Is a Condition With Increasing Importance in Medical Oncology. *Oncologist*. 2016;21(2):e1.
- Joglekar S, Nau PN, Mezhir JJ. The impact of sarcopenia on survival and complications in surgical oncology: A review of the current literature-- Author response. *J Surg Oncol*. 2015;112(8):910.
- Mourtzakis M, Prado CM, Lieffers JR, Reiman T, Mccargar LJ, Baracos VE. A practical and cise approach to quantification of body composition in cancer patients using computed tomography images acquired during routine care. *Appl Physiol Nutr Metab*. 2008;33(5):997-1006.

© 2018 Lehigh Valley Health Network