

Assessing Appropriateness of Magnetic Resonance Imaging for Headaches within the Lehigh Valley Health Network

Manveer Dilts-Garcha MA BSc

Follow this and additional works at: <https://scholarlyworks.lvhn.org/select-program>

Part of the [Medical Education Commons](#)

Published In/Presented At

Dilts-Garcha, M. (2018, March). *Assessing Appropriateness of Magnetic Resonance Imaging for Headaches within the Lehigh Valley Health Network*. Poster Presented at: Poster Presented at: 2018 SELECT Capstone Posters and Presentations Day. Kasych Family Pavilion, Lehigh Valley Health Network, Allentown, PA.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Assessing Appropriateness of Magnetic Resonance Imaging for Headaches within the Lehigh Valley Health Network

Manveer Dilts-Garcha MA BSc, Mentor: Gary Clauser MD
Lehigh Valley Health Network, Allentown, PA

Background

- Migraines cost the healthcare system and society at large more than \$14 billion annually
- Medicare spends \$146 to \$211 million on imaging for headaches annually
- Office based neuroimaging costs at approximately \$1 billion annually
- Only 1% to 3% of neuroimaging studies for chronic headaches reveal significant abnormalities
- Neuroimaging rates have increased despite multiple guidelines aimed at reducing overuse

Problem Statement

To what degree do MRIs - ordered for the evaluation of headaches in the outpatient setting at LVHN - meet the guideline recommendations as laid out by the American College of Radiology (ACR) in 2013 and the AAN.

Methods

A retrospective chart review was performed looking at MRIs ordered with the following associated diagnosis codes:

- headache (ICD10-R51)
- chronic headache (ICD10- G43.719)
- migraine without aura (ICD10-43.009)
- migraine with aura (ICD10-43.109)

Charts were manually reviewed and assigned an appropriateness score per the 2013 ACR Guidelines and per commonly cited red flag criteria.


Key outcome measures included percentage of studies ordered appropriately as well as the percentage of studies with clinically significant findings.

Results


- 37% of studies were determined to be appropriate
- 36% were inappropriate
- 27% lacked sufficient data to make a determination
- On average 4% of the studies showed significant abnormalities on neuroimaging

	Family	IM	Neuro	All
Count	85	51	66	202
Insignificant	78	48	60	186
Significant	3	3	2	8
% Significant	4%	6%	3%	4%


Appropriateness of Neuroimaging



ACR Score Unknown Relative Frequency



Significant Findings on Imaging



Discussion

- The percentage of clinically significant abnormalities found (4%) is in line with prior research on imaging for chronic headaches
- Results indicate significant room for improvement with only a third of studies being ordered appropriately. The remaining two-thirds were either inappropriate or lacked sufficient information in the note to make a determination
- Absence of a note, lack of chronicity, prior history, and medication use history were the most common information lacking
- All significant findings were associated with appropriate indications for neuroimaging findings. In one case, the wrong kind of imaging was ordered. This is reassuring as to the validity of the guidelines.

Conclusions

- We can do better in terms of the neuroimaging we order at LVHN
- Areas with potential significant potential for improvement include
 - guideline adherence
 - documentation
- Future efforts for improvement may focus on educating providers on key red flag features, and which history/physical components are most important for determining whether imaging is warranted

References:

1. Callaghan, B. C., Kerber, K. A., Pace, R. J., Skolarus, L. E., & Burke, J. F. (2014). Headaches and neuroimaging: High utilization and costs despite guidelines. *JAMA Internal Medicine*, 174(5), 819-821. doi:10.1001/jamainternmed.2014.173
2. Clarke, C. E., Edwards, J., Nicholl, D. J., & Sivaguru, A. (2010). Imaging results in a consecutive series of 530 new patients in the Birmingham Headache Service. *Journal of Neurology*, 257(8), 1274-1278. doi:10.1007/s00415-010-5506-7
3. Douglas, A. C., Wippold, F. J., Broderick, D. F., Aiken, A. H., Amin-Hanjani, S., Brown, D. C., Corey, A. S., Germano, I. M., Hadley, J. A., Jagadeesan, B. D., and Jurgens, J. S. (2014). ACR appropriateness criteria headache. *Journal of the American College of Radiology*, 11(7), pp.657-667.
4. Frisberg, B. M., Rosenberg, J. H., Matchar, D. B., McCrory, D. C., Pietrzak, M. P., Rozen, T. D., & Silberstein, S. D. (2000). Evidence-based guidelines in the primary care setting: neuroimaging in patients with nonacute headache. *St Paul, MN: US Headache Consortium*.
5. Headache Classification Committee of the International Headache Society (IHS). (2013). The international classification of headache disorders, (beta version). *Cephalalgia*.
6. Katz, M. (2016). The Cost-Effective Evaluation of Uncomplicated Headache. *The Medical Clinics Of North America*, 100(5), 1009-1017. doi:10.1016/j.mcn.2016.04.009
7. Loder, E., Weizenbaum, E., Frisberg, B., Silberstein, S. and American Headache Society Choosing Wisely Task Force (2013). Choosing Wisely in Headache Medicine: The American Headache Society's List of Five Things Physicians and Patients Should Question. *Headache: The Journal of Head and Face Pain*, 53: 1651-1659. doi:10.1111/head.12233
8. Rasmussen, B. K., Jensen, R., Schroll, M., & Olesen, J. (1991). Epidemiology of headache in a general population -- A prevalence study. *Journal of Clinical Epidemiology*, 44(11), 1147-1157. doi:10.1016/0895-4356(91)90147-2
9. Schwartz, A. L., Landon, B. E., Elshaug, A. G., Chermew, M. E., & McWilliams, J. (2014). Measuring low-value care in medicare. *JAMA Internal Medicine*, 174(7), 1067-1076. doi:10.1001/jamainternmed.2014.1541
10. Sempere, A. P., Porta-Etessam, J., Medrano, V., Garcia-Morales, I., Concepcion, L., Ramos, A., Botella, C. (2005). Neuroimaging in the evaluation of patients with non-acute headache. *Cephalalgia*, 25(1), 30-35. doi:10.1111/j.1468-2982.2004.00798.x
11. Urbaniak, G. C., & Plous, S. (2015). Research Randomizer (Version 4.0) [Computer software]. Retrieved on July 3, 2017, from <http://www.randomizer.org/>
12. Wang, H.Z., Simonson, T.M., Greco, W.R. and Yuh, W.T., 2001. Brain MR imaging in the evaluation of chronic headache in patients without other neurologic symptoms. *Academic radiology*, 8(5), pp.405-408.

© 2018 Lehigh Valley Health Network