



Scientia Agropecuaria

Website: <http://revistas.unitru.edu.pe/index.php/scientiaagrop>

Facultad de Ciencias
Agropecuarias

Universidad Nacional de
Trujillo

HS-SPME-GC-MS detection of volatile compounds in *Myrciaria jaboticaba* Fruit

Gerby Giovanna Rondán Sanabria^{1,2,*}; Ausberta Jesus Cabezas Garcia²; Antonio William Oliveira Lima³; Brousett-Minaya Magaly Alejandra⁴; Narendra Narain²

¹ Dirección de Investigación, Universidad Tecnológica del Perú (UTP), Av. Tacna y Arica 160, Arequipa, Peru.

² Laboratory of Flavor and Chromatographic Analysis, Federal University of Sergipe, 49100-000 São Cristóvão, SE, Brazil.

³ Institute of Chemistry, São Paulo University, Av. Prof. Lineu Prestes, 748 – Butantã, 05508-000, São Paulo, SP, Brazil.

⁴ Facultad de Ciencias Naturales y formales, Universidad Nacional de San Agustín, Av. Independencia s/n, Arequipa, Peru.

Received December 10, 2017. Accepted August 13, 2018.

Abstract

This study was performed to investigate the volatile compounds for the characteristic aroma in jaboticaba fruit distributed in southern and central regions of Brazil. The present work combines headspace solid phase microextraction (HS-SPME) and high resolution gas chromatography-mass spectrometry (GC-MS) techniques to identify and quantify the volatile compounds. The influence of different SPME fibers (CAR/PDMS and DVB/CAR/PDMS) in extraction of volatile compounds was evaluated. The effects of extraction temperature and salt concentration (NaCl) in the extraction medium were studied using the response surface methodology in order to achieve the highest extraction efficiency. The better extraction of volatile compounds was achieved by using a CAR/PDMS fiber and the optimum adsorption conditions were at 42 °C for 30 min and 5% NaCl concentration. A total of 71 compounds were identified, among these, 57% were terpenes which was the most representative class of compounds, followed by esters (19%), aldehydes (10%), alcohols (5.5%) and aromatics compounds (4.4%) and other organic compounds 2.8%. Limonene and ethyl acetate were the volatile compounds that showed highest relative concentration and these could contribute to the characteristic aroma of the jaboticaba fruit along with other compounds such as β -pinene, δ -cadinene, linalool, β -guaiene, and α -caryophyllene.

Keywords: jaboticaba; *Myrciaria*; volatile compounds; headspace solid phase microextraction; gas chromatography mass spectrometry.

1. Introduction

Jaboticaba is a Brazilian native fruit from the Atlantic rainforest that belongs to the Myrtaceae family, grape-like in appearance and texture. Its economic importance has been continuously growing in Brazil because of the sweet and slightly acidic flavor of the pulp. Jaboticaba is a small fruit, grape-like in appearance and is one of Brazil's richest sources in anthocyanins. The most common varieties are *Myrciaria cauliflora* (DC) O. Berg, *Myrciaria jaboticaba* (Vell) Berg and jaboticaba-assú. The fruit grows directly on the main trunk and branches, has a diameter varying from 1-4 centimeters, with 1 - 4 large seeds per fruit and richly colorful epicarpio covering the white sweet pulp and jelly-like in appearance (Reynertson *et al.*, 2006). It is mostly consumed as fresh fruit or turned

into various industrial products are also available such as juice, jellies, wines and liquors.

The interest of this fruit has grown in recent years because it has in its chemical composition phenolic constituents such as anthocyanins, flavonoids and ellagitannins. The fruit is also reported to possess well defined biological properties including antioxidant and anti-inflammatory activities (Middleton *et al.*, 2000). Although the jaboticaba is a fruit with important functional properties for health the components that give it a pleasant taste and aroma have not yet been investigated. The aroma is one of the most important quality attributes of tropical and subtropical fruits and numerous analytical techniques have been developed to characterize the compounds responsible

* Corresponding author
E-mail: C16238@utp.edu.pe (G. Rondán).

for its aroma quality. Most of the volatile compounds of the fruit are composed of saturated and unsaturated molecules from aliphatic compounds that have functional oxygen groups such as ester, alcohol, acid, aldehyde, ketone or lactone, which originate during the maturation process by different metabolic pathways. A recent study reported that the high sugar content associated with terpenes contribute to result in sweet, acidic and slight resinous flavor of the jaboticaba fruits (Plagemann *et al.*, 2012).

The flavour impact is a complex mixture of volatile organic chemicals and taste contributors. Quality perception cannot be analysed and it is seldom possible to trace the complex composition of flavour and taste. However, one of the most interesting techniques for studying flavour quality is HS-SPME (Souza-Silva *et al.*, 2015). There is growing interest in this field to elucidate whether the flavour profiles of fresh fruits and vegetables are affected by the cultivation system (Vallverdú-Queralt and Lamuela-Raventós, 2016).

The development of fast and simple methods for volatiles extraction is important because it can reduce the sampling time and analysis can be performed quickly. A fast sampling technique that can be adapted for gas chromatography (GC) analysis of volatile compounds is the solid phase microextraction (SPME). This technique has been applied for the analysis of volatile and nonvolatile compounds in liquid and gaseous samples (Arthur and Pawliszyn, 1990) and to analyze the flavor in juices and vegetable oils (Yang and Peppard, 1994). SPME is a pretty fast and reliable technique for the determination of volatile compounds in complex matrices. It is based on the extraction of analytes from the sample matrix using a fused silica fiber (1 - 2 cm), coated with a thin film of absorbent material which in most cases is of polymeric nature. Subsequently, in order to separate and identify the organic compounds, the desorption of analytes is carried out by increasing the temperature, where the fiber is held into the GC injection port for its instrumental analysis (Lasekan and Abbas, 2010). HS-SPME technique has been used to evaluate the pitomba fruit (*Talisia esculenta* Radlk.) (De Souza *et al.*, 2016); strawberries, raspberries, blackberries and bananas (Ibanez *et al.*, 1998); Bayberry (Cheng *et al.*, 2015); passion fruit, cashew apple, tamarind, guava and acerola (Carasek and Pawliszyn, 2006); grape berry (Parker *et al.*, 2007) and mango (Liu *et al.*, 2013).

In this study a new system was developed consisting of HS-SPME for the capture of volatile compounds on fiber containing different porous material, followed by desorption of volatile compounds onto a system of GC-MS, wherein a capillary column was used to separate the volatile compounds, identify and quantify these compounds present in jaboticaba fruit pulp. The study was performed to select the best fiber for the volatile analysis, along with the optimization of the extraction conditions such as the temperature and ionic strength of extraction media and analyzing the data obtained by response surface methodology interpretation.

2. Materials and methods

Sample

Fresh ripe jaboticaba fruits having no apparent injury or microbial spoilage were purchased from a local market of Aracaju city in the state of Sergipe, Brazil. The fruits were washed with water and disinfected using 1% sodium hypochlorite solution whereafter the fruits were again washed with water. The pulp of jaboticaba fruit was removed manually using a stainless steel knife. After the extraction of the seed, the pulp and peel were homogenized in a waring blender without addition of water, then it was filtered and the resulting homogenate was kept in polythene bags and stored in a freezer maintained at -20 °C until further analysis. For the extraction of volatiles compounds, the samples were thawed at room temperature and diluted with distilled water (Milli-Q Integral Water Purification System) in an aqueous solution of sodium chloride (Sigma Chemical Co., Aldrich Chemical Co.) at different concentrations. The final proportion of pulp and the solution was of 3.5:10.

Selection of SPME fibers in the extraction of volatile compounds

Tests were performed using different fibers such as carboxen/polydimethylsiloxane (CAR/PDMS; 85 µm) and divinylbenzene/Carboxen/polydimethylsiloxane (DVB/CAR/PDMS; 50/30 µm), both supplied by Supelco (Bellefonte, PA, USA) and these were evaluated to verify the ability to collect and concentrate the headspace volatiles of jaboticaba homogenate, mainly due to their sorption characteristics. These fibers were exposed to HS using SPME holder in 40 ml vials containing a mixture of 3.5 g of fruit homogenate and 10 ml of deionized water. The vials were sealed with a polypropylene plug and extraction perfor-

med for 30 min at 25 °C under constant magnetic stirring. After the extraction, SPME fibers were removed from the vials and inserted into the GC injection system (Agilent GC system model 7820A) - MS (Agilent MS model 5975), for the separation and identification of volatiles. Experiments containing only fruit homogenate with distilled water were performed between the extraction processes also to verify the results that there was no interference from earlier runs.

Extraction Conditions

After the selection of the best SPME fiber for sorption of volatiles of jabuticaba homogenate, the best experimental conditions for the extraction of volatiles by HS-SPME were determined. The Response Surface methodology (RSM) was used wherein the independent variables were salt concentration or ionic strength (IS) and extraction temperature (T). The data were interpreted on the basis of the dependent variable being the number of detected peaks (NP). A Central Composite Design (CCD) with two variables at five levels (-1, 1.41, 0, +1, 1.41) was used and the center point was replicated six times, totalizing 14 experiments. The experimental conditions in the center point were: salt concentration 15% NaCl and 35 °C. The experimental values varied were: X_1 = ionic strength (0, 5, 15, 26 and 30%) and X_2 = temperature (25, 28, 35, 42 and 45 °C). For HS-SPME analysis, 3.5 g of homogenate was mixed with 10 ml aqueous solution of NaCl at different concentrations in a hermetically sealed 40 mL vials with a polypropylene plug. Subsequently the mixture was heated at different temperatures (25, 28, 35, 42 and 45 °C) in thermostatic bath for 15 min in order to reach thermal equilibrium with constant magnetic stirring. Later the selected fiber was inserted into the vial and was exposed to HS for 30 minutes to adsorb the volatiles. The SPME fiber was immediately taken to the GC injection port and thermally desorbed for 8 min at 250 °C in split mode. The number of detected peaks in each run was used as a response variable of the experimental design and a quadratic model was constructed to describe the observed data.

Chromatographic analysis

Chromatographic analysis was performed using GC-MS system equipped with a DB-5 capillary column (inner diameter: 30 m x 0.25 mm, film thickness: 0.25 µm). The oven temperature was programmed initially at 40 °C for 4 min, then increased at the rate of 7 °C/min to 160 °C wherein held for 8 min,

subsequently the temperature raised at 15 °C/min up to 250 °C and maintained for 3 min at this temperature, totalizing 38 min of chromatographic analysis. Helium was used as carrier gas at a constant rate of 1.0 mL/min. The mass spectrometer was operated in scan mode from 29 – 550 m/z, at 70 eV ionization.

The identification of volatiles compounds was performed by calculating the linear retention index (LRI) of each compound using a series of alkane standards analyzed under identical conditions as that of the samples and comparing the mass spectrum of the compound peaks with mass spectra library of NIST/EPA (NIH version 2.0). The values of LRI of each compound were also compared with those described in the literature as well as from the data base of compounds identification in our laboratory wherein standards analysis was performed in identical analytical conditions. The relative concentrations of individual components were expressed as a percentage of the relative peak area to that of the total peak area.

Statistical analysis

The statistical significance of the model was evaluated by analysis of variance (ANOVA) with significance level of 5% ($p < 0,05$). Statistica V.7 (Statsoft Inc., Tulsa, OK, USA) was used in order to generate ANOVA and response surface methodology interpretations.

3. Results and discussion

Optimization of extraction conditions using HS-SPME

The extraction time, extraction temperature, mass of the sample and sodium chloride concentration are the factors which influence the vapor pressure of the volatiles in the headspace. The optimization method evaluated only considered the effect of the salt addition associated with the adsorption temperature, while the other variables remained constant during the tests. The response was evaluated to determine the best extraction conditions for HS-SPME based on the analysis of chromatograms relating to higher number of identified volatile compounds and the intensity of the response. However, before optimization of the extraction conditions, the adsorption profile of the 3 SPME fibers was evaluated and in later experiments only the selected fiber was used.

SPME Fibers

The selection of the most suitable SPME fiber depends on the target compounds and the sample under study (Gioti *et al.*, 2007).

Two semi polar fibers - *CAR/PDMS* and *DVB/CAR/PDMS* were tested, and compared individually. The results reported in [Figure 1A](#) show that the *CAR/PDMS* fiber resulted in a better profile having high extraction efficiency of volatile compounds in jaboticaba as it resulted in greater number of detected compounds and their signal strength. *CAR/PDMS* fiber coating is most suitable for the extraction of small volatile molecules. The sensitivity of this fiber into smaller molecules, such as esters, acids and non-polar compounds, allows the extraction of a broad range of aromatic compounds ([Érica et al., 2017](#)). The compounds with the highest intensity were the ethyl acetate (22.04%) and limonene (11.47%) ([Figure 1B](#)). Furthermore, the main components adsorbed were the terpenoids, including α -pinene, β -pinene, β -ocimene, β -caryophyllene, α -

muurolene, δ -cadinene, calamenene and limonene, which could be responsible for the characteristic aroma of the fruit. Considering better results obtained by using the *CAR/PDMS* fiber, it was selected for further studies of the characterization of volatile compounds of jaboticaba fruit.

Effects of the combination of sodium chloride and extraction temperature

The response surface methodology (RSM) is a powerful tool that enables quickly and efficiently to interpret the data while having only a minimum number of experiments. A total of 14 experiments ([Table 1](#)) were performed in random combinations of temperature and ionic strength to check the effects of the combination of sodium chloride and the extraction temperature, the results are presented in [Figure 2](#).




Figure 1. Influence of the fiber (*PDMS/DVD/CAR* and *PDMS/CAR*) on the extraction of volatile compounds from jaboticaba homogenate: A - Relative concentration (%) of the number of detected peaks; B - Concentration (%) of the main detected compounds.

Table 1

Description of independent variables analyzed in Central Composite Design showing the experimental and predicted values of number of peaks in chromatogram

Experiment/N°	Coded Levels		Independent variables		Peak number	
	X1	X2	IS	T / °C	Experimental	Predicted
1	-1	-1	5	28	53	55.011
2	-1	1	5	42	89	87.052
3	1	-1	26	28	46	50.700
4	1	1	26	42	81	81.741
5	0	0	15	35	63	62.167
6	0	0	15	35	63	62.167
7	0	0	15	35	62	62.167
8	-1.414	0	0	35	78	78.525
9	1.414	0	30	35	75	71.722
10	0	-1.414	15	25	44	39.824
11	0	1.414	15	45	83	84.423
12	0	0	15	35	62	62.167
13	0	0	15	35	62	62.167
14	0	0	15	35	61	62.167

(-1.41) Low level, (1.41) High level, (0) Central point, (IS) Ionic Strength and (T) Temperature.

The response surface (Figure 2A) was obtained using mathematical model expressed in coded variables by interpreting the data using Statistica software, which resulted in the following equation:

$$Y_{(NP)} = 62.1669 - 2.4050 X_1 + 15.7700 X_2 + 6.4800 X_1 X_1 - 0.0150 X_2 X_2 - 0.2500 X_1 X_2 \text{ (eq.1)}$$

Salt concentration (X_1) and temperature (X_2), as well as quadratic term ionic strength ($X_1 X_1$) were significant ($p < 0.05$) and these influenced the volatile compounds extraction as shown in the Pareto Chart (Figure 1B). It can be observed that the temperature was the most significant parameter ($p < 0.05$), showing a strong positive influence. The model had a value $R = 0.97$, considered appropriate for extraction and values greater than 0.90 are considered statistically significant. It was also observed that the maximum number of detected compounds were found at the surface of the peak, this fact could indicate the best experimental conditions. According to this model, the best combination of extraction was at 42 °C with 5% NaCl, wherein 89 volatile compounds were separated as shown in the chromatogram (Figure 3).

CCD was an important tool in the optimization of the extraction method of volatiles because traditional optimization methods evaluate only the effect of a variable while keeping the other variables constant during the experiments. However, this type of test suffers from the limitation that it does not take into account as to what would happen if other variables also change. The experimental design (response surface + central composite design) used in this study allowed the estimation of the effects of two variables simultaneously.

The experimental design used in this work has been earlier used in studies for the extraction of volatile compounds from

several fruits such as cupuassu (Oliveira et al., 2004), orange juice (Mirhosseini and Tan, 2009; Charve et al., 2011), cherimoya (*Annona cherimila*), soursop (*Annona muricata*) (Ferreira et al., 2009; Cheong, et al., 2011) and umbu (*Spondias tuberosas*) (Galvão et al., 2011).





Figure 2. A - Response surface generated by quadratic model (eq. 1) in the optimization of the temperature (T, °C) and ionic strength (IS, %) in the extraction of volatile compounds of jaboticaba fruit using *HS-SPME*; B - Pareto chart showing statistically significant variables ($p < 0.05$) influencing the response during the extraction of volatile compounds.

Identification and quantification of volatile compounds

Figure 3 shows the typical total ion chromatogram obtained under best extrac-

tion conditions. The 15 compounds which are numbered in the chromatogram represent the compounds with higher intensity. Table 2 shows the data on areas of the detected peaks, expressed in percentage, for the different compounds that were used to indicate the relative concentration of each substance along with their characteristic odor note. A total of 71 compounds were identified according to the similarity of their mass spectra and by comparison with the standard values described in the literature.

Among the identified compounds, 41 (57.7%) were terpenes, 14 (19.7%) esters, 7 (9.90%) aldehydes, four (5.60%) alcohols, three (4.20%) aromatic compounds. The compounds that showed the highest concentration were: limonene (17.7%) followed by ethyl acetate (10.7%), 1,3-dichlorobenzene (9.7%), eucalyptol (5.8%) caryophyllene (5.0%), ocimene (3.6%), β -pinene (3.0%), β -guaiene (2.9%), 2-butanolic acid ethyl ester (2.5%) and α -pinene (2.2%).

Plagemann et al. (2012) identified only 45 volatile compounds from jaboticaba pulp, among which the β -pinene, δ -cadinene, 2-phenylethanol and linalool were the most prominent compounds in the fruit flavor. These authors also attributed to terpenes as the major contributors to the flavor, indicating that the most odorous compounds were β -caryophyllene and limonene.

In the essential oil of jaboticaba leaves (*Myrciaria cauliflora*), Duarte et al. (2010) identified compounds similar to those detected in this study, including α -pinene, β -pinene, limonene, δ -elemene, β -elemene, β -caryophyllene, δ -cadinene and α -muurolene, which may be responsible for the aroma of the fruit.

Comparing the results of this study with those found in other typical fruits of the Amazon of the same genus (*Myrciaria*), there was similarity in the presence of several compounds, such as in the fruit of camu-camu (*Myrciaria dúbia*) showing α -pinene (17.5), limonene (10.8), β -caryophyllene and β -pinene were reported. The β -pinene is also the most important volatile compound in umbu-cajá (*Spondias cintherea*) (Franco and Shibamoto, 2000). In pitanga (*Eugenia inflora* L.) fruit, the presence of β -pinene and β -ocimene as important constituents for aroma of the fruit was also reported (Oliveira et al., 2006). While chinese bayberry had a stronger "pine" odor, which is caused mainly by α -pinene (Cheng et al., 2015).

Based on the identification and quantification of volatile compounds present in the jaboticaba fruit, it could be concluded that the main compounds such as limonene, ethyl acetate, eucalyptol, caryophyllene, ocimene, β -pinene and α -pinene could contribute to the characteristic aroma quality of this fruit.




Figure 3. Representative chromatogram of the volatile compounds of jaboticaba homogenate. The predominant peaks and the most striking aroma compounds are numbered as follows: Ethyl acetate (1), 2-butanolic acid, ethyl ester, (E) - (2), α -pinene (3), β -Pinene (4), Benzene 1,3-dichloro- (5), Benzene, 1-methyl-3-(1-methylethyl) - (6), limonene (7), Eucalyptol (8), β -ocimene, (9), β -caryophyllene (10), β - guaieno (11), α -muuroleno (12), δ -cadinene (13), calameneno (14) and linalool (15).

Table 2

Volatile compounds identified in jaboticaba fruit pulp

Compounds ^a	LRI		Relative area percent	Odor description ^c	Identification ^d
	DB-5	Ref.			
Alcohols					
Ethanol	369,59	---	1,75%	sweet	MS
3-methyl-1-butanol	730,45	1,2,3,7	0,05%	whiskey, malt, burnt	MS, LRI
(Z)-3-Hexenol	857,60	1,2,3,7	0,21%	grass	MS, LRI
1-Hexanol	871,34	1,2,3,10	0,18%	resin, flower, green	MS, LRI
Aldehydes					
3-Methyl-1-butanol	581,58	---	0,12%		MS
Pentanal	603,85	---	0,11%	almond, malt, pungent	MS
Hexanal	801,18	1,3,4,5,12	0,79%	grass, tallow, fat	MS, LRI
(E)-2-Hexenal	854,72	1,3,4,5	0,18%	green, leaf	MS, LRI
Nonanal	1105,58	3,4,7,10,12	0,39%	fat, citrus, green	MS, LRI
Decanal	1206,82	1,3,4,6,7,12	0,07%	soap, orange peel, tallow	MS, LRI
Aromatic compounds					
Styrene	890,20	4,10,11	0,14%	balsamic, gasoline	MS, LRI
Benzaldehyde	961,45	1,3,4,7,12	0,02%	almond, burnt sugar	MS, LRI
1,3-dichlorobenzene	1011,55	---	9,70%		MS
p-cymene	1025,73	3,6,11,16	3,13%	solvent, gasoline, citrus	MS, LRI
Esters					
Ethyl Acetate	503,21	---	10,71%	pineapple	MS
Ethyl propanoate	711,06	8	0,05%	Fruit	MS, LRI
Propyl acetate	713,12	8	0,11%	celery, ethereal, fruity, pear, powerful, raspberry	MS, LRI
Methyl (E)-2-butenolate	762,14	3	0,24%		MS
2-Methylpropyl acetate	773,63	---	0,07%	fruit, apple, banana	MS
Ethyl butanoate	803,98	2,3,6,8	0,44%	apple	MS, LRI
Ethyl (E)-2-butenolate	846,42	3,7	2,48%	Fruity	MS, LRI
3-methylbutyl acetate	879,47	---	0,39%	banana	MS
Methyl (E)-2-hexanoate	968,28	9,8,10	0,08%		MS, LRI
Ethyl hexanoate	1001,02	2,3,7,8,9	0,21%	apple peel, fruit	MS, LRI
(Z)-3-hexenyl acetate	1008,73	7,8,11	0,22%	Unripe banana, fruit, green	MS, LRI
(Z)-3-Hexenyl butanoate	1187,44	10	0,09%	wine, green	MS, LRI
Ethyl octanoate	1197,93	2,3	0,09%	fruit, fat, floral, pear	MS, LRI
Methyl 3-phenylpropanoate	1387,43	11	0,08%	strawberry	MS, LRI
Ether					
Ethyl ether	395,29	---	0,19%		MS
Ketones					
2-Heptanone	892,56	11	0,10%	soap	MS, LRI
Terpenoids					
Tricyclene	927,89	3	0,49%		MS, LRI
α-Pinene	934,01	1,4,3,6	2,15%	Terpenic, woody	MS, LRI
Sabinene	974,44	3,6,7,13	0,08%	pepper, turpentine, wood	MS, LRI
β-Pinene	976,80	1,3,4,13	2,98%	Sweet, green	MS, LRI
β-myrcene	992,14	3,6,7	1,35%	Herbaceous, woody	MS, LRI
α-Phellandrene	1004,30	3,6	1,41%	Citrus, spicy	MS, LRI
Limonene	1029,99	1,3,4,6,12,13	17,71%	Citrus, terpenic	MS, LRI
Eucalyptol	1032,28	5,13,16	5,79%	mint, sweet	MS, LRI
(E)-β-Ocimene	1050,18	3,6,4,13	3,57%	sweet, herb	MS, LRI
γ-Terpinene	1060,63	3,4,6	0,34%	gasoline, turpentine	MS, LRI
Terpinolene	1089,88	3,6,12	0,16%	Lime, terpenic	MS, LRI
Linalool	1100,45	1,3,7,10,13	0,32%	flower, lavender	MS, LRI
1,3,8-p-Menthatriene	1132,22	3,6,11	0,27%		MS, LRI
(4E,6Z)-allo-Ocimene	1143,70	5,6	0,25%	Green, fresh	MS, LRI
α-Terpineol	1193,52	3,7	0,21%	oil, anise, mint	MS, LRI
δ-Elemene	1343,45	1,11,14	0,61%	wood	MS, LRI
α-Cubebene	1355,82	1,5,6	0,45%	herb, wax	MS, LRI
α-Copaene	1378,82	1,3,5,6	0,09%	Woody, spicy	MS, LRI
β-Cubebene	1383,31	---	0,99%		MS
β-elemene	1398,23	3,4,11,13,15	1,41%	herb, wax, fresh	MS, LRI
β-Caryophyllene	1429,10	1,4,3,5	5,01%	Spicy, woody	MS, LRI
Aromadendrene	1438,10	16	0,36%	Dusty, honey, wood	MS, LRI
α-Guaiene	1440,14	11,15	0,25%	wood, balsamic	MS, LRI
β-Gurjunene	1444,12	6	0,21%		MS, LRI
β-farnesene	1448,81	11	0,47%	wood, citrus, sweet	MS, LRI
α-Elemene	1454,94	15	0,38%	wood	MS, LRI
Allo-Aromadendrene	1459,36	5,14	0,36%	wood	MS, LRI
α-Caryophyllene	1463,50	3,5,14	0,80%	wood	MS, LRI
γ-Murolene	1472,28	3,6,5,16	1,33%	herb, wood, spice	MS, LRI
Eremophilene	1484,32	3,6	0,84%		MS, LRI
Germacrene D	1490,50	3,6,13,14	2,92%	wood, spice	MS, LRI
Valencene	1496,47	3,11	1,00%	green, oil	MS, LRI
α-Selinene	1499,12	3,6,13	0,20%		MS, LRI
Epizonarene	1502,28	16	0,62%		MS, LRI
α-Murolene	1505,26	13,14	3,92%	wood	MS, LRI
γ-Cadinene	1513,49	1,5,16	0,27%	wood	MS, LRI
δ-Cadinene	1520,29	3,5,6,13	0,83%	thyme, medicine, wood	MS, LRI
Calamenene	1528,06	3,11	2,20%	herb, spice	MS, LRI
Naphthalene, 1,2,3,4,4a,7-hexahydro-1,6-dimethyl-4-(1-methylethyl)-	1537,04	---	0,15%		MS
Germacrene B	1564,31	1,10,11,14	0,27%	wood, earth, spice	MS, LRI
Caryophyllene oxide	1593,21	1,3,5	0,38%	herb, sweet, spice	MS, LRI
Saturated hydrocarbons					
n-Hexane	478,80	---	0,26%	Alkane	MS

^a Compounds are listed in their elution order on a DB-5 column;^b LRI: Linear retention indices on column (DB-5) determined with n-alkanes and literature and from the data-base of our laboratory;^c Odor description based on the Leffingwell's Flavor-Base (Leffingwell, 2007) and Flavornet (<http://www.flavornet.org/flavornet.html>);^d Identification method: MS = mass spectrum; LRI is compared with references from standards or literature values as well as from the data base of Volatile compounds analyzed in our laboratory;Ref: ¹Erica et al., (2017); ²Boulangier and Crouzet (2001); ³Pino (2012); ⁴Cheng et al. (2015); ⁵Porat et al. (2011); ⁶Liu et al. (2013); ⁷Lasekan et al. (2013); ⁸Christof et al., (2014); ⁹Márquez et al. (2011); ¹⁰Liqin et al., (2017); ¹¹Acree and Arn (2004); ¹²Cuevas et al. (2016); ¹³Plagemann et al. (2012); ¹⁴Duarte et al. (2010); ¹⁵Huang et al. (2009); and ¹⁶Parker et al. (2007).

4. Conclusions

The combination of HS-SPME technique for capture and GC-MS system for identification of volatile compounds in jaboticaba fruit resulted in identification of a large number of volatile compounds belonging to the organic classes of terpenes, esters, aldehydes, alcohols and aromatic compounds, many of these have not been reported earlier. Among the 2 SPME fibers (CAR/PDMS and DVB/CAR/PDMS) evaluated for the capture of volatile compounds, the fiber CAR/PDMS resulted in higher number of compounds in jaboticaba fruit. Based on this study, it could be concluded that the combination of HS-SPME with GC-MS is suitable for the extraction of volatile compounds from jaboticaba fruit. A large number of esters and terpenic compounds could be responsible for the typical aroma of the jaboticaba fruit, the principal compounds being limonene, ethyl acetate, eucalyptol, caryophyllene, ocimene, β -pinene and α -pinene. Based on the data obtained, the authors recommended further studies taking into accounts the harvest season. In addition to chromatography gases coupled with mass spectroscopy analysis olfactory to confirm the volatile compounds responsible for the aroma of the fruit.

Acknowledgments

All authors thank the financial support provided from the project, National Institute of Science and Technology (INCT) for Tropical Fruits from CNPq, Brazil offered for promoting research work undertaken in this study. The co-authors Gerby and Ausberta thank the CAPES (Ministry of Education, Brazil) for providing fellowship to her.

References

- Acree, T.; Arn, H. 2004. Gas chromatography - olfactometry (GCO) of natural products Sponsored by DATU Inc. Flavornet retrieved from: <http://www.flavornet.org>.
- Arthur, C.L.; Pawliszyn, J. 1990. Solid phase microextraction with thermal desorption using fused silica optical fibers. *Analytical Chemistry* 62: 2145-2148.
- Boulanger, R.; Crouzet, J. 2001. Identification of the aroma components of acerola (*Malpighia glabra* L.): free and bound flavor compounds. *Food Chemistry* 74: 209-216.
- Carasek, E.; Pawliszyn, A. 2006. Screening of tropical fruit volatile compounds using Solid-Phase Microextraction (SPME) fibers and internally cooled SPME fiber. *Journal Agricultural and Food Chemistry* 54: 8688-8696.
- Charve, J.; Chen, Ch.; Hegeman, A.D.; Reineccius, G.A. 2011. Evaluation of instrumental methods for the untargeted analysis of chemical stimuli of orange juice flavour. *Journal of Flavour and Fragrance* 26: 429-440.
- Cheng, H.; Chen, J.; Chen, S.; Wu, D.; Liu, D.; Ye, X. 2015. Characterization of aroma-active volatiles in three Chinese bayberry (*Myrica rubra*) cultivars using GC-MS-olfactometry and an electronic nose combined with principal component analysis. *Food Research International* 72: 8-15.
- Cheong, K.W.; Tan, C.P.; Mirhosseini, H.; Chin, S.T.; Man, Y.B.C.; Hamid, N.S.A. et al. 2011. Optimization of equilibrium headspace analysis of volatile flavor compounds of Malaysian soursop (*Annona muricata*): Comprehensive two-dimensional gas chromatography time-of-flight mass spectrometry (GC/GC-TOFMS). *Food Chemistry* 125: 1481-1489.
- Christof, B.S.; Tara, G.; Reinhold, C. 2014. Influence of harvest maturity and fruit logistics on pineapple (*Ananas comosus* [L.] Merr.) Volatiles assessed by headspace solid phase microextraction and gas chromatography-mass spectrometry (HS-SPME-GC/MS). *Food Chemistry* 150: 382-391.
- Cuevas, F.J.; Moreno-Rojas, J.M.; Arroyo, F.; Daza, A.; Ruiz-Moreno, M.J. 2016. Effect of management (organic vs conventional) on volatile profiles of six plum cultivars (*Prunus salicina* Lindl.). A chemometric approach for varietal classification and determination of potential markers. *Food Chemistry* 199: 479-484.
- De Souza, M.P.; Bataglion, G.A.; da Silva, F.M.A.; de Almeida, R. A.; Paz, W.H.P.; Nobre, T.A.; Marinho, J.V.N.; Salvador, M.J.; Fidelis, C.H.V.; Acho, L.D.R.; de Souza, A.D.L.; et al. 2016. Phenolic and aroma compositions of pitomba fruit (*Talisia esculenta* Radlk.). *Food Research International* 87: 87-94.
- Duarte, R.A.; Santos, S.C.; Seraphin, J.C.; Ferri, P.H. 2010. Environmental influence on phenols and essential oils of *Myrciaria cauliflora* leaves. *Journal of the Brazilian Chemical Society* 21(9): 1672-1680.
- Érica, A.; Souza, S.; Saboia, G.; Nina, C.J.; Hoffmann, C.; Dos Santos, R.S.I.; Geraldo, L. G. Soares, G.L.; Zini, C.A. 2017. Development of a HS-SPME-GC/MS protocol assisted by chemometric tools to. *Talanta* 175: 9-20.
- Ferreira, L.; Perestrello, R.; Câmara, J.S. 2009. Comparative analysis of the volatile fraction from Annona cherimola Mill. cultivars by solid-phase microextraction and gas chromatography-quadrupole mass spectrometry detection. *Talanta* 77: 1087-1096.
- Franco, M.R.B.; Shibamoto, T. 2000. Volatile composition of some Brazilian fruits: Umu-cajá (*Spondias cytherea*), camu-camu (*Myrciaria dubia*), araçá-boi (*Eugenia stipitata*), and cupuaçu (*Theobroma grandiflorum*). *Journal of Agricultural and Food Chemistry* 48: 1263-1265.
- Galvão, S.M.; Narain, N.; dos Santos, P.S.M.; Nunes, M.L. 2011. Volatile compounds and descriptive odor attributes in umbu (*Spondias tuberosa*) fruits during maturation. *Food Research International* 44: 1919-1926.
- Gioti, M.E.; Fiamegos, C.Y.; Skalkos, C.D.; Atalikas, D.C. 2007. Improved method for the in vitro assessment of antioxidant activity of plant extracts by headspace solid-phase microextraction and gas chromatography-electron capture detection. *J. of Chromatography A* 1152(1): 150-155.
- Huang, B.; Qin, L. Chu, Q.; Zhang, Q.; Gao, L.; Zheng, H. 2009. Comparison of headspace SPME with hydrodistillation and SFE for analysis of the volatile components of the roots of *Valeriana officinalis* var. latifolia. *Chromatographia* 69: 489-496.
- Ibanez, E.; Lopez-Sebastian, S.; Ramos, E.; Tabera, J. Reglero, G. 1998. Analysis of volatile fruit components by headspace solid-phase microextraction. *Food Chemistry* 63: 281-286.
- Lasekan, O.; Khatib, A.; Juhari, H.; Patiram, P.; Lasekan, S. 2013. Headspace solid-phase microextraction gas chromatography-mass spectrometry determination of volatile compounds in different

- varieties of African star apple fruit (*Chrysophyllum albidum*). Food Chemistry 141: 2089–2097.
- Lasekan, O.; Abbas, K. 2010. Flavour chemistry of palm toddy and palm juice: a review. Trends in Food Science & Technology 21: 494-501.
- Leffingwell. 2007. Flavor Base. Leffingwell & Associates. Available in: www.leffingwell.com/flavbase.htm
- Liqin, Y.; Chunxiang, Y.; Weidong, L.; Jiangbo, H.; Sun, M.; Jingru, Z.; Zhongshuang, Z. 2017 Evaluation of volatile compounds from Chinese dwarf cherry (*Cerasus humilis* (Bge.) Sok.) germplasms by headspace solid-phase microextraction and gas chromatography–mass spectrometry. Food Chemistry 217: 389–397.
- Liu, F.X.; Fu, S.F.; Bi, X.F.; Chen, F. et al. 2013. Physico-chemical and antioxidant properties of four mango (*Mangifera indica* L.) cultivars in china. Food Chemistry 138: 396-405.
- Márquez, C.J.; Jimenez, A.M.; Osorio, C.; Cartagena, J.R. 2011. Volatile compounds during the ripening of colombian soursop (*Annona muricata* L. cv. Elita). Vitae, Revista de la Facultad de Química Farmacéutica 18(3):245-250.
- Middleton, E.J.R.; Kandaswami, C.; Theoharides, T.C. 2000. The Effects of Plant Flavonoids on Mammalian Cells: Implications for Inflammation, Heart Disease, and Cancer. Pharmacological Reviews 52: 673–751.
- Mirhosseini, H.; Tan, C.P. 2009. Response surface methodology and multivariate analysis of equilibrium headspace concentration of orange beverage emulsion as function of emulsion composition and structure. Food Chemistry 115: 324–333.
- Oliveira, L.A.; Lopes, B.R.; Fernando, A.; Cabral, M.N.; Eberlin. 2006. Volatile compounds from pitanga fruit (*Eugenia uniflora* L.). Food Chemistry 99: 1–5.
- Oliveira, M.A.; Pereira, N.R.; Marsaioli, Jr.A.; Augusto, F. 2004. Studies on the aroma of cupuassu liquor by headspace solid-phase microextraction and gas chromatography. Journal of Chromatography A 1025: 115–124.
- Parker, M.; Pollnitz, A.P.; Cozzolino, D.; Leigh-Francis, I.; Herderich, M.J. 2007. Identification and quantification of a marker compound for 'pepper' aroma and flavor in shiraz grape berries by combination of chemometrics and gas chromatography-mass spectrometry. Journal of Agricultural and Food Chemistry 55: 5948-5955.
- Pino, J.A. 2012. Odour-active compounds in mango (*Mangifera indica* L. cv. Corazón). International Journal of Food Science and Technology 47: 1944 – 1950.
- Plagemann, I.; Krings, I.; Berger, R.G.; Maróstica, Jr. M.R. 2012. Volatile constituents of jabuticaba (*Myrciaria jabuticaba* (Vell.) O. Berg) fruits. Journal of Essential Oil Research 24(1): 45–51.
- Porat, R.; Tietel, Z.; Zippori, I.; Dag, A. 2011. Aroma volatile compositions of high-and low-aromatic guava varieties. Journal of the Science and Food Agriculture 91: 2794–2798.
- Reynertson, K.A.; Wallace, A.M, Adachi, S.; Gil, R.R.; Yang, H.; Basile, M.J. 2006. Bioactive depsides and anthocyanins from jabuticaba (*Myrciaria cauliflora*). Journal of Natural Products 69: 1228–1230. .
- Souza-Silva, É.A.; Gionfriddo, E.; Pawliszyn, J. 2015. A critical review of the state of the art of solid-phase microextraction of complex matrices II. Food analysis. TrAC Trends in Analytical Chemistry 71: 236-248.
- Vallverdú, Q.A.; Lamuela-Raventós, R.M. 2016. Foodomics: A new tool to differentiate between organic and conventional foods. Electrophoresis. 13:1784-94.
- Yang, X.; Peppard, T. 1994. Solid-phase microextraction for flavor analysis. Journal Agricultural of Food Chemistry 42: 1925-1930.