

SIGNED STAR k -DOMATIC NUMBER OF A GRAPH

SEYED MAHMOUD SHEIKHOESLAMI AND LUTZ VOLKMANN

ABSTRACT. Let G be a simple graph without isolated vertices with vertex set $V(G)$ and edge set $E(G)$ and let k be a positive integer. A function $f : E(G) \rightarrow \{-1, 1\}$ is said to be a signed star k -dominating function on G if $\sum_{e \in E(v)} f(e) \geq k$ for every vertex v of G , where $E(v) = \{uv \in E(G) \mid u \in N(v)\}$. A set $\{f_1, f_2, \dots, f_d\}$ of signed star k -dominating functions on G with the property that $\sum_{i=1}^d f_i(e) \leq 1$ for each $e \in E(G)$, is called a signed star k -dominating family (of functions) on G . The maximum number of functions in a signed star k -dominating family on G is the signed star k -domatic number of G , denoted by $d_{kSS}(G)$.

In this paper we study the properties of the signed star k -domatic number $d_{kSS}(G)$. In particular, we determine the signed star k -domatic number of some classes of graphs. Some of our results extend these one given by Atapour et al. [1] for the signed star domatic number.

1. INTRODUCTION

Let G be a graph with vertex set $V(G)$ and edge set $E(G)$. We use [8] for terminology and notation which are not defined here and consider simple graphs without isolated vertices only. For every nonempty subset E' of $E(G)$, the subgraph $G[E']$ induced by E' is the graph whose vertex set consists of those vertices of G incident with at least one edge of E' and whose edge set is E' .

Two edges e_1 and e_2 of G are called *adjacent* if they are distinct and have a common vertex. The *open neighborhood* $N_G(e)$ of an edge $e \in E(G)$ is the set of all edges adjacent to e . Its *closed neighborhood* is $N_G[e] = N_G(e) \cup \{e\}$. For a function $f : E(G) \rightarrow \{-1, 1\}$ and a subset S of $E(G)$ we define $f(S) = \sum_{e \in S} f(e)$. The *edge-neighborhood* $E_G(v)$ of a vertex $v \in V(G)$ is the set of all edges incident with the vertex v . For each vertex $v \in V(G)$, we also define $f(v) = \sum_{e \in E_G(v)} f(e)$.

Received by the editors September 2, 2010, and in revised form May 25, 2011.

2000 *Mathematics Subject Classification.* 05C69.

Key words and phrases. Signed star k -domatic number; Signed star domatic number; Signed star k -dominating function; Signed star dominating function; Signed star k -domination number; Signed star domination number; Regular graphs.

Let k be a positive integer. A function $f : E(G) \rightarrow \{-1, 1\}$ is called a *signed star k -dominating function* (SSkDF) on G , if $f(v) \geq k$ for every vertex v of G . The *signed star k -domination number* of a graph G is $\gamma_{kSS}(G) = \min\{\sum_{e \in E} f(e) \mid f \text{ is a SSkDF on } G\}$. The signed star k -dominating function f on G with $f(E(G)) = \gamma_{kSS}(G)$ is called a $\gamma_{kSS}(G)$ -*function*. As assuming $\delta(G) \geq k$ is clearly necessary, we will always assume that when we discuss $\gamma_{kSS}(G)$ all graphs involved satisfy $\delta(G) \geq k$. The signed star k -domination number, introduced by Xu and Li in [11], has been studied by several authors (see for instance [2, 7]). The signed star 1-domination number is the usual signed star domination number which has been introduced by Xu in [9] and has been studied by several authors (see for instance [5, 6, 10]).

A set $\{f_1, f_2, \dots, f_d\}$ of signed star k -dominating functions on G with the property that $\sum_{i=1}^d f_i(e) \leq 1$ for each $e \in E(G)$, is called a *signed star k -dominating family* (of functions) on G . The maximum number of functions in a signed star k -dominating family on G is the *signed star k -domatic number* of G , denoted by $d_{kSS}(G)$. The signed star k -domatic number is well-defined and $d_{kSS}(G) \geq 1$ for all graphs G with $\delta(G) \geq k$, since the set consisting of any one SSkD function forms a SSkD family on G . A d_{kSS} -*family* of a graph G is a SSkD family containing $d_{kSS}(G)$ SSkD functions. The signed star 1-domatic number $d_{1SS}(G)$ is the usual signed star domatic number $d_{SS}(G)$ which was introduced by Atapour et al. in [1].

Our purpose in this paper is to initiate the study of signed star k -domatic number in graphs. We first study basic properties and bounds for the signed star k -domatic number of a graph, some of which are analogous to those of the signed star domatic number $d_{SS}(G)$ in [1]. In addition, we determine the signed star k -domatic number of some classes of graphs.

Observation 1.1. *Let G be a graph of order $n \geq 3$ and size m . If $k \in \{n-2, n-1\}$ and $\delta(G) \geq k$, then $\gamma_{kSS}(G) = m$ and hence $d_{kSS}(G) = 1$.*

Observation 1.2. *Let G be a graph of size m with $\delta(G) \geq k$. Then $\gamma_{kSS}(G) = m$ if and only if each edge $e \in E(G)$ has an endpoint u such that $\deg(u) = k$ or $\deg(u) = k+1$.*

Proof. If each edge $e \in E(G)$ has an endpoint u such that $\deg(u) = k$ or $\deg(u) = k+1$, then trivially $\gamma_{kSS}(G) = m$.

Conversely, assume that $\gamma_{kSS}(G) = m$. Suppose to the contrary that there exists an edge $e = uv \in E(G)$ such that $\min\{\deg(u), \deg(v)\} \geq k+2$. Define $f : E(G) \rightarrow \{-1, 1\}$ by $f(e) = -1$ and $f(e') = 1$ for $e' \in E(G) \setminus \{e\}$. Obviously, f is a signed star k -dominating function of G with weight less than m , a contradiction. This completes the proof. \square

2. BASIC PROPERTIES OF THE SIGNED STAR k -DOMATIC NUMBER

In this section we study basic properties of $d_{kSS}(G)$.

Theorem 2.1. *Let G be a graph of size m with $\delta(G) \geq k$, signed star k -domination number $\gamma_{kSS}(G)$ and signed star k -domatic number $d_{kSS}(G)$. Then*

$$\gamma_{kSS}(G) \cdot d_{kSS}(G) \leq m.$$

Moreover, if we have $\gamma_{kSS}(G) \cdot d_{kSS}(G) = m$, then for each d_{kSS} -family $\{f_1, f_2, \dots, f_d\}$ of G , each function f_i is a γ_{kSS} -function and $\sum_{i=1}^d f_i(e) = 1$ for all $e \in E(G)$.

Proof. If $\{f_1, f_2, \dots, f_d\}$ is a signed star k -dominating family on G such that $d = d_{kSS}(G)$, then the definitions imply

$$\begin{aligned} d \cdot \gamma_{kSS}(G) &= \sum_{i=1}^d \gamma_{kSS}(G) \leq \sum_{i=1}^d \sum_{e \in E(G)} f_i(e) \\ &= \sum_{e \in E(G)} \sum_{i=1}^d f_i(e) \leq \sum_{e \in E(G)} 1 = m \end{aligned}$$

as desired.

If $\gamma_{kSS}(G) \cdot d_{kSS}(G) = m$, then the two inequalities occurring in the proof become equalities. Hence for the d_{kSS} -family $\{f_1, f_2, \dots, f_d\}$ of G and for each i , $\sum_{e \in E(G)} f_i(e) = \gamma_{kSS}(G)$, thus each function f_i is a γ_{kSS} -function, and $\sum_{i=1}^d f_i(e) = 1$ for all $e \in E(G)$. \square

Corollary 2.2. *If G is a graph of size m and $\delta(G) \geq k$, then*

$$\gamma_{kSS}(G) + d_{kSS}(G) \leq m + 1.$$

Proof. By Theorem 2.1,

$$(2.1) \quad \gamma_{kSS}(G) + d_{kSS}(G) \leq d_{kSS}(G) + \frac{m}{d_{kSS}(G)}.$$

Using the fact that the function $g(x) = x + m/x$ is decreasing for $1 \leq x \leq \sqrt{m}$ and increasing for $\sqrt{m} \leq x \leq m$, this inequality leads to the desired bound immediately. \square

Corollary 2.3. *Let G be a graph of size m and $\delta(G) \geq k$. If $2 \leq \gamma_{kSS}(G) \leq m - 1$, then*

$$\gamma_{kSS}(G) + d_{kSS}(G) \leq m.$$

Proof. Theorem 2.1 implies that

$$(2.2) \quad \gamma_{kSS}(G) + d_{kSS}(G) \leq \gamma_{kSS}(G) + \frac{m}{\gamma_{kSS}(G)}.$$

If we define $x = \gamma_{kSS}(G)$ and $g(x) = x + m/x$ for $x > 0$, then because $2 \leq \gamma_{kSS}(G) \leq m - 1$, we have to determine the maximum of the function

g in the interval $I : 2 \leq x \leq m - 1$. It is easy to see that

$$\begin{aligned} \max_{x \in I} \{g(x)\} &= \max\{g(2), g(m-1)\} \\ &= \max\left\{2 + \frac{m}{2}, m-1 + \frac{m}{m-1}\right\} \\ &= m-1 + \frac{m}{m-1} < m+1, \end{aligned}$$

and we obtain $\gamma_{kSS}(G) + d_{kSS}(G) \leq m$. This completes the proof. \square

Corollary 2.4. *Let $k \geq 1$ be an integer, and let G be a graph of size m and $\delta(G) \geq k$. If $\min\{\gamma_{kSS}(G), d_{kSS}(G)\} \geq 2$, then*

$$\gamma_{kSS}(G) + d_{kSS}(G) \leq \frac{m}{2} + 2.$$

Proof. Since $\min\{\gamma_{kSS}(G), d_{kSS}(G)\} \geq 2$, it follows by Theorem 2.1 that $2 \leq d_{kSS}(G) \leq m/2$. By (2.1) and the fact that the maximum of $g(x) = x + m/x$ on the interval $2 \leq x \leq m/2$ is $g(2) = g(m/2)$, we see that

$$\gamma_{kSS}(G) + d_{kSS}(G) \leq d_{kSS}(G) + \frac{m}{d_{kSS}(G)} \leq \frac{m}{2} + 2.$$

\square

Observation 1.1 demonstrates that Corollary 2.4 is no longer true in the case that $\min\{\gamma_{kSS}(G), d_{kSS}(G)\} = 1$.

Theorem 2.5. *Let G be a graph with $\delta(G) \geq k$ and let $v \in V(G)$. Then*

$$d_{kSS}(G) \leq \begin{cases} \frac{\deg(v)}{k} & \text{if } \deg(v) \equiv k \pmod{2}, \\ \frac{\deg(v)}{k+1} & \text{if } \deg(v) \equiv k+1 \pmod{2}. \end{cases}$$

Moreover, if the equality holds, then for each function f_i of a SSkD family $\{f_1, f_2, \dots, f_d\}$ and for every $e \in E(v)$,

$$\sum_{e \in E(v)} f_i(e) = \begin{cases} k & \text{if } \deg(v) \equiv k \pmod{2}, \\ k+1 & \text{if } \deg(v) \equiv k+1 \pmod{2}, \end{cases}$$

and $\sum_{i=1}^d f_i(e) = 1$.

Proof. Let $\{f_1, f_2, \dots, f_d\}$ be a SSkD family of G such that $d = d_{kSS}(G)$. If $\deg(v) \equiv k \pmod{2}$, then

$$d = \sum_{i=1}^d 1 \leq \sum_{i=1}^d \frac{1}{k} \sum_{e \in E(v)} f_i(e) = \frac{1}{k} \sum_{e \in E(v)} \sum_{i=1}^d f_i(e) \leq \frac{1}{k} \sum_{e \in E(v)} 1 = \frac{\deg(v)}{k}.$$

Similarly, if $\deg(v) \equiv k + 1 \pmod{2}$, then

$$\begin{aligned} d &= \sum_{i=1}^d 1 \leq \sum_{i=1}^d \frac{1}{k+1} \sum_{e \in E(v)} f_i(e) \\ &= \frac{1}{k+1} \sum_{e \in E(v)} \sum_{i=1}^d f_i(e) \leq \frac{1}{k+1} \sum_{e \in E(v)} 1 = \frac{\deg(v)}{k+1}. \end{aligned}$$

If $d_{kSS}(G) = \deg(v)/k$ when $\deg(v) \equiv k \pmod{2}$ or $d_{kSS}(G) = \deg(v)/(k+1)$ when $\deg(v) \equiv k+1 \pmod{2}$, then the two inequalities occurring in the proof of each corresponding case become equalities, which gives the properties given in the statement. \square

Corollary 2.6. *Let G be a graph and $1 \leq k \leq \delta(G)$. Then*

$$d_{kSS}(G) \leq \begin{cases} \frac{\delta(G)}{k} & \text{if } \delta(G) \equiv k \pmod{2}, \\ \frac{\delta(G)}{k+1} & \text{if } \delta(G) \equiv k+1 \pmod{2}. \end{cases}$$

Theorem 2.7. *The signed star k -domatic number is an odd integer.*

Proof. Let G be an arbitrary graph, and assume that $d = d_{kSS}(G)$ is even. Let $\{f_1, f_2, \dots, f_d\}$ be the corresponding signed star k -dominating family on G . If $e \in E(G)$ is an arbitrary edge, then $\sum_{i=1}^d f_i(e) \leq 1$. On the left-hand side of this inequality, a sum of an even number of odd summands occurs. Therefore it is an even number, and we obtain $\sum_{i=1}^d f_i(e) \leq 0$ for each $e \in E(G)$. This forces

$$kd = \sum_{i=1}^d k \leq \sum_{i=1}^d \sum_{e \in E(v)} f_i(e) = \sum_{e \in E(v)} \sum_{i=1}^d f_i(e) \leq 0,$$

which is a contradiction. \square

An immediate consequence of Theorems 2.5, 2.7 and Corollary 2.6 is the following result.

Corollary 2.8. *Let G be a graph with $\delta(G) \geq k$. If $\delta(G) < 3k$ or if G has a vertex v of degree $\deg(v) = 3k + 1$, then $d_{kSS}(G) = 1$.*

Proof. If $\delta(G) < 3k$, then Corollary 2.6 implies that

$$d_{kSS}(G) \leq \frac{\delta(G)}{k} < \frac{3k}{k} = 3.$$

Applying Theorem 2.7, we deduce that $d_{kSS}(G) \leq 1$ and thus $d_{kSS}(G) = 1$. If G has a vertex v of degree $\deg(v) = 3k + 1$, then $\deg(v) \equiv k + 1 \pmod{2}$ and thus it follows from Theorem 2.5 that

$$d_{kSS}(G) \leq \frac{\deg(v)}{k+1} = \frac{3k+1}{k+1} < 3.$$

Again Theorem 2.7 leads to the desired result $d_{kSS}(G) = 1$. \square

Corollary 2.9. *Let G be a graph of size m . Then $\gamma_{kSS}(G) + d_{kSS}(G) = m + 1$ if and only if each edge $e \in E(G)$ has an endpoint u such that $\deg(u) = k$ or $\deg(u) = k + 1$.*

Proof. If each edge $e \in E(G)$ has an endpoint u such that $\deg(u) = k$ or $\deg(u) = k + 1$, then $\gamma_{kSS}(G) = m$ by Observation 1.2. Hence $d_{kSS}(G) = 1$ and the result follows.

Conversely, let $\gamma_{kSS}(G) + d_{kSS}(G) = m + 1$. The result is obviously true for $m = 1, 2, 3$. Assume $m \geq 4$. By Corollary 2.4, we may assume that $\min\{\gamma_{kSS}(G), d_{kSS}(G)\} = 1$. If $\gamma_{kSS}(G) = 1$, then $d_{kSS}(G) = m$, which is a contradiction to Corollary 2.6. If $d_{kSS}(G) = 1$, then $\gamma_{kSS}(G) = m$ and the result follows by Observation 1.2. \square

As an application of Corollary 2.6 and Theorem 2.7, we will prove the following Nordhaus-Gaddum type result.

Theorem 2.10. *For every graph G of order n with $\delta(G) \geq k$ and $\delta(\overline{G}) \geq k$,*

$$(2.3) \quad d_{kSS}(G) + d_{kSS}(\overline{G}) \leq \frac{n-1}{k}.$$

If $d_{kSS}(G) + d_{kSS}(\overline{G}) = (n-1)/k$, then G is regular, k and $\delta(G)$ are even and n is odd such that $n-1 \equiv 0 \pmod{4}$.

Proof. Since $\delta(G) + \delta(\overline{G}) \leq n-1$, Corollary 2.6 leads to

$$d_{kSS}(G) + d_{kSS}(\overline{G}) \leq \frac{\delta(G)}{k} + \frac{\delta(\overline{G})}{k} \leq \frac{n-1}{k}.$$

If G is not regular, then $\delta(G) + \delta(\overline{G}) \leq n-2$ and hence we obtain the better bound $d_{kSS}(G) + d_{kSS}(\overline{G}) \leq (n-2)/k$. Thus assume now that G is $\delta(G)$ -regular.

Case 1: Assume that k is odd. If $\delta(G)$ is even, then it follows from Corollary 2.6 that

$$\begin{aligned} d_{kSS}(G) + d_{kSS}(\overline{G}) &\leq \frac{\delta(G)}{k+1} + \frac{\delta(\overline{G})}{k} = \frac{\delta(G)}{k+1} + \frac{n-\delta(G)-1}{k} \\ &< \frac{\delta(G)}{k} + \frac{n-\delta(G)-1}{k} = \frac{n-1}{k}. \end{aligned}$$

If $\delta(G)$ is odd, then n is even and thus $\delta(\overline{G}) = n - \delta(G) - 1$ is even. Using Corollary 2.6, we find that

$$\begin{aligned} d_{kSS}(G) + d_{kSS}(\overline{G}) &\leq \frac{\delta(G)}{k} + \frac{\delta(\overline{G})}{k+1} = \frac{\delta(G)}{k} + \frac{n-\delta(G)-1}{k+1} \\ &< \frac{\delta(G)}{k} + \frac{n-\delta(G)-1}{k} = \frac{n-1}{k}. \end{aligned}$$

Combining these two bounds, we conclude that $d_{kSS}(G) + d_{kSS}(\overline{G}) < (n-1)/k$ when k is odd.

Case 2: Assume that k is even. If $\delta(G)$ is odd, then Corollary 2.6 implies $d_{kSS}(G) + d_{kSS}(\overline{G}) < (n-1)/k$ as above. If $\delta(G)$ is even and n is even, then

$\delta(\overline{G}) = n - \delta(G) - 1$ is odd, and we obtain the bound $d_{kSS}(G) + d_{kSS}(\overline{G}) < (n - 1)/k$ as above.

Finally, assume that $\delta(G)$ is even and n is odd such that $n - 1 = 4p + 2$. If $d_{kSS}(G) + d_{kSS}(\overline{G}) = (n - 1)/k$, then we observe that

$$d_{kSS}(G) = \frac{\delta(G)}{k} \quad \text{and} \quad d_{kSS}(\overline{G}) = \frac{\delta(\overline{G})}{k}.$$

According to Theorem 2.7, these two values are odd integers, say

$$d_{kSS}(G) = \frac{\delta(G)}{k} = 2s + 1 \quad \text{and} \quad d_{kSS}(\overline{G}) = \frac{\delta(\overline{G})}{k} = 2t + 1.$$

If $k = 2i$, then we arrive at the contradiction

$$d_{kSS}(G) + d_{kSS}(\overline{G}) = \frac{\delta(G)}{k} + \frac{\delta(\overline{G})}{k} = 2(s + t + 1) = \frac{4p + 2}{2i}.$$

This contradiction completes the proof of Theorem 2.10. \square

The following examples will demonstrate that $d_{kSS}(G) + d_{kSS}(\overline{G}) = (n - 1)/k$ in Theorem 2.10 is possible when G is regular, k and $\delta(G)$ are even and n is odd such that $n - 1 \equiv 0 \pmod{4}$.

Let $k \geq 2$ be an even integer and $n \geq 5$ such that $n - 1 = 2k$. Now let H be a k -regular graph of order n . Then \overline{H} is also k -regular. Corollary 2.6 implies that $d_{kSS}(H) \leq 1$ and thus $d_{kSS}(H) = 1$. It follows that

$$d_{kSS}(H) + d_{kSS}(\overline{H}) = 2 = \frac{n - 1}{k}.$$

Corollary 2.11. *Let G be a graph of order n with $\delta(G) \geq k$ and $\delta(\overline{G}) \geq k$. If $\delta(G) < 3k$ and $\delta(\overline{G}) < 3k$ or $n < 4k + 1$, then*

$$d_{kSS}(G) + d_{kSS}(\overline{G}) = 2.$$

Proof. If $\delta(G) < 3k$ and $\delta(\overline{G}) < 3k$, then Corollary 2.8 implies the desired result immediately. If $n < 4k + 1$, then it follows from (2.3) that

$$d_{kSS}(G) + d_{kSS}(\overline{G}) \leq \frac{n - 1}{k} < \frac{4k}{k} = 4,$$

and thus Theorem 2.7 leads to $d_{kSS}(G) + d_{kSS}(\overline{G}) = 2$. \square

3. SIGNED STAR k -DOMATIC NUMBER OF REGULAR GRAPHS

In this section we determine values of the signed star k -domatic number for some classes of regular graphs.

Theorem 3.1. *Let G be an r -regular and 1-factorable graph and let $1 \leq k \leq r$ be an integer. Then*

$$d_{kSS}(G) = \begin{cases} \left\lfloor \frac{r}{k} \right\rfloor & \text{when } r \equiv k \pmod{2} \text{ and } \left\lfloor \frac{r}{k} \right\rfloor \text{ is odd,} \\ \left\lfloor \frac{r}{k} \right\rfloor - 1 & \text{when } r \equiv k \pmod{2} \text{ and } \left\lfloor \frac{r}{k} \right\rfloor \text{ is even,} \\ \left\lfloor \frac{r}{k+1} \right\rfloor & \text{when } r \equiv k+1 \pmod{2} \text{ and } \left\lfloor \frac{r}{k+1} \right\rfloor \text{ is odd,} \\ \left\lfloor \frac{r}{k+1} \right\rfloor - 1 & \text{when } r \equiv k+1 \pmod{2} \text{ and } \left\lfloor \frac{r}{k+1} \right\rfloor \text{ is even.} \end{cases}$$

Proof. By Observation 1.2 and Theorem 2.1 we may assume $k \leq r - 2$. Let $\{M_0, M_1, \dots, M_{r-1}\}$ be a 1-factorization of G . We distinguish two cases.

Case 1: Assume that $r \equiv k \pmod{2}$. Suppose that $r = kq + t$, where q is a positive integer and $0 \leq t \leq k - 1$. By Corollary 2.6 and Theorem 2.7, $d_{kSS}(G) \leq q$ if q is odd and $d_{kSS}(G) \leq q - 1$ if q is even.

Subcase 1.1: Assume that q is odd. Then t is even. Define the functions f_1, f_2, \dots, f_q as follows.

$$f_1(e) = \begin{cases} 1 & \text{if } e \in M_i \text{ where } 0 \leq i \leq \frac{k(q-1)}{2} + k - 1, \\ -1 & \text{if } e \in M_i \text{ and } \frac{k(q-1)}{2} + k \leq i \leq kq - 1, \end{cases}$$

and for $2 \leq j \leq q$ and $0 \leq i \leq kq - 1$,

$$f_j(M_i) = f_{j-1}(M_{i+2k}),$$

where the sum is taken modulo kq . In addition, if $t > 0$,

$$f_j(M_i) = (-1)^{i+j} \text{ for } 1 \leq j \leq q \text{ and } kq \leq i \leq r - 1.$$

It is easy to see that f_j is a signed star k -dominating function of G for each $1 \leq j \leq q$ and $\{f_1, f_2, \dots, f_q\}$ is a signed star k -dominating family of G . Hence $d_{kSS}(G) \geq q$. Therefore $d_{kSS}(G) = q$, as desired.

Subcase 1.2: Assume that q is even. Then $t + k$ is even. Define the functions f_1, f_2, \dots, f_{q-1} as follows.

$$f_1(M_i) = \begin{cases} 1 & \text{if } 0 \leq i \leq \frac{k(q-2)}{2} + k - 1, \\ -1 & \text{if } \frac{k(q-2)}{2} + k \leq i \leq k(q-1) - 1, \end{cases}$$

and for $2 \leq j \leq q - 1$ and $0 \leq i \leq k(q-1) - 1$,

$$f_j(M_i) = f_{j-1}(M_{i+2k}),$$

where the sum is taken modulo $k(q-1)$. In addition,

$$f_j(M_i) = (-1)^{i+j} \text{ for } 1 \leq j \leq q \text{ and } k(q-1) \leq i \leq r - 1.$$

It is easy to see that f_j is a signed star k -dominating function of G for each $1 \leq j \leq q-1$ and $\{f_1, f_2, \dots, f_{q-1}\}$ is a signed star k -dominating family on G . Hence $d_{kSS}(G) \geq q-1$ and so $d_{kSS}(G) = q-1$.

Case 2: Assume that $r \equiv k+1 \pmod{2}$. Suppose that $r = (k+1)q + t$, where q is a positive integer and $0 \leq t \leq k$. By Corollary 2.6 and Theorem 2.7, $d_{kSS}(G) \leq q$ if q is odd and $d_{kSS}(G) \leq q-1$ if q is even.

Subcase 2.1: Assume that q is odd. Then t is even. Define the functions f_1, f_2, \dots, f_q as follows.

$$f_1(M_i) = \begin{cases} 1 & \text{if } 0 \leq i \leq \frac{(k+1)(q-1)}{2} + k, \\ -1 & \text{if } \frac{(k+1)(q-1)}{2} + k + 1 \leq i \leq (k+1)q - 1, \end{cases}$$

and for $2 \leq j \leq q$ and $0 \leq i \leq (k+1)q - 1$,

$$f_j(M_i) = f_{j-1}(M_{i+2(k+1)}),$$

where the sum is taken modulo $(k+1)q$. In addition, if $t > 0$,

$$f_j(M_i) = (-1)^{i+j} \text{ for } 1 \leq j \leq q \text{ and } (k+1)q \leq i \leq r-1.$$

It is easy to see that f_j is a signed star k -dominating function of G for each $1 \leq j \leq q$ and $\{f_1, f_2, \dots, f_q\}$ is a signed star k -dominating family of G . Hence $d_{kSS}(G) \geq q$. Therefore $d_{kSS}(G) = q$, as desired.

Subcase 2.2: Assume that q is even. Then $t+k+1$ is even. Define the functions f_1, f_2, \dots, f_{q-1} as follows.

$$f_1(M_i) = \begin{cases} 1 & \text{if } 0 \leq i \leq \frac{(k+1)(q-2)}{2} + k, \\ -1 & \text{if } \frac{(k+1)(q-2)}{2} + k + 1 \leq i \leq (k+1)(q-1) - 1, \end{cases}$$

and for $2 \leq j \leq q-1$ and $0 \leq i \leq (k+1)(q-1) - 1$,

$$f_j(M_i) = f_{j-1}(M_{i+2(k+1)}),$$

where the sum is taken modulo $(k+1)(q-1)$. In addition,

$$f_j(M_i) = (-1)^{i+j} \text{ for } 1 \leq j \leq q \text{ and } (k+1)(q-1) \leq i \leq n-1.$$

It is easy to see that f_j is a signed star k -dominating function of G for each $1 \leq j \leq q-1$ and $\{f_1, f_2, \dots, f_{q-1}\}$ is a signed star k -dominating family of G . Hence $d_{kSS}(G) \geq q-1$ and so $d_{kSS}(G) = q-1$, as desired. \square

Applying Theorem 3.1 and the well-known classical Theorem of König [4] that a k -regular bipartite graph is 1-factorable, we obtain the next result.

Corollary 3.2. *If G is an r -regular bipartite graph and $1 \leq k \leq r$ is an integer, then*

$$d_{kSS}(G) = \begin{cases} \left\lfloor \frac{r}{k} \right\rfloor & \text{when } r \equiv k \pmod{2} \text{ and } \left\lfloor \frac{r}{k} \right\rfloor \text{ is odd,} \\ \left\lfloor \frac{r}{k} \right\rfloor - 1 & \text{when } r \equiv k \pmod{2} \text{ and } \left\lfloor \frac{r}{k} \right\rfloor \text{ is even,} \\ \left\lfloor \frac{r}{k+1} \right\rfloor & \text{when } r \equiv k+1 \pmod{2} \text{ and } \left\lfloor \frac{r}{k+1} \right\rfloor \text{ is odd,} \\ \left\lfloor \frac{r}{k+1} \right\rfloor - 1 & \text{when } r \equiv k+1 \pmod{2} \text{ and } \left\lfloor \frac{r}{k+1} \right\rfloor \text{ is even.} \end{cases}$$

Theorem 3.3. *Let G be a graph of order n and factorable into r Hamiltonian cycles and let $1 \leq k \leq 2r$ be an integer. Then*

$$d_{kSS}(G) = \begin{cases} \left\lfloor \frac{2r}{k} \right\rfloor & \text{when } k \text{ is even and } \left\lfloor \frac{2r}{k} \right\rfloor \text{ is odd,} \\ \left\lfloor \frac{2r}{k} \right\rfloor - 1 & \text{when } k \text{ and } \left\lfloor \frac{2r}{k} \right\rfloor \text{ are even,} \\ \left\lfloor \frac{2r}{k+1} \right\rfloor & \text{when } k \text{ and } \left\lfloor \frac{2r}{k+1} \right\rfloor \text{ are odd,} \\ \left\lfloor \frac{2r}{k+1} \right\rfloor - 1 & \text{when } k \text{ is odd and } \left\lfloor \frac{2r}{k+1} \right\rfloor \text{ is even.} \end{cases}$$

Proof. Let G be a Hamiltonian factorable graph, and let $\{C_0, C_1, \dots, C_{r-1}\}$ be a Hamiltonian factorization of G . We distinguish two cases.

Case 1: Assume that k is even. Suppose that $2r = kq + t$, where q is a positive integer and $0 \leq t \leq k - 1$. By Corollary 2.6 and Theorem 2.7, $d_{kSS}(G) \leq q$ if q is odd and $d_{kSS}(G) \leq q - 1$ if q is even.

Subcase 1.1: Assume that q is odd. Then t is even and $r = (k/2)q + (t/2)$. Define the functions f_1, f_2, \dots, f_q as follows.

$$f_1(C_i) = \begin{cases} 1 & \text{if } 0 \leq i \leq \frac{k(q-1)}{4} + \frac{k}{2} - 2, \\ -1 & \text{if } \frac{k(q-1)}{4} + \frac{k}{2} - 1 \leq i \leq \frac{k}{2}q - 1, \end{cases}$$

and for $2 \leq j \leq q$ and $0 \leq i \leq \frac{k}{2}q - 1$,

$$f_j(C_i) = f_{j-1}(C_{i+k}),$$

where the sum is taken modulo $(k/2)q$. In addition, if $t > 0$,

$$f_j(C_i) = (-1)^{i+j} \text{ for } 1 \leq j \leq q \text{ and } \frac{k}{2}q \leq i \leq r - 1.$$

It is easy to see that f_j is a signed star k -dominating function of G for each $1 \leq j \leq q$ and $\{f_1, f_2, \dots, f_q\}$ is a signed star k -dominating family of G . Hence $d_{kSS}(G) \geq q$. Therefore $d_{kSS}(G) = q$, as desired.

Subcase 1.2: Assume that q is even. Then $(k/2) + (t/2)$ is even. Define the functions f_1, f_2, \dots, f_{q-1} as follows.

$$f_1(C_i) = \begin{cases} 1 & \text{if } 0 \leq i \leq \frac{k(q-2)}{4} + \frac{k}{2} - 2, \\ -1 & \text{if } \frac{k(q-2)}{4} + \frac{k}{2} - 1 \leq i \leq \frac{k}{2}(q-1) - 1, \end{cases}$$

and for $2 \leq j \leq q-1$ and $0 \leq i \leq (k/2)(q-1) - 1$,

$$f_j(M_i) = f_{j-1}(M_{i+k}),$$

where the sum is taken modulo $(k/2)(q-1)$. In addition,

$$f_j(C_i) = (-1)^{i+j} \text{ for } 1 \leq j \leq q \text{ and } \frac{k}{2}(q-1) \leq i \leq r-1.$$

It is easy to see that f_j is a signed star k -dominating function of G for each $1 \leq j \leq q-1$ and $\{f_1, f_2, \dots, f_{q-1}\}$ is a signed star k -dominating family on G . Hence $d_{kSS}(G) \geq q-1$ and so $d_{kSS}(G) = q-1$.

Case 2: Assume that k is odd. Suppose that $2r = (k+1)q + t$, where q is a positive integer and $0 \leq t \leq k$. By Corollary 2.6 and Theorem 2.7, $d_{kSS}(G) \leq q$ if q is odd and $d_{kSS}(G) \leq q-1$ if q is even.

Subcase 2.1: Assume that q is odd. Then t is even. Define the functions f_1, f_2, \dots, f_q as follows.

$$f_1(C_i) = \begin{cases} 1 & \text{if } 0 \leq i \leq \frac{(k+1)(q-1)}{4} + \frac{k+1}{2} - 2, \\ -1 & \text{if } \frac{(k+1)(q-1)}{4} + \frac{k+1}{2} - 1 \leq i \leq \frac{(k+1)}{2}q - 1, \end{cases}$$

and for $2 \leq j \leq q$ and $0 \leq i \leq (k+1)q/2 - 1$,

$$f_j(C_i) = f_{j-1}(C_{i+(k+1)}),$$

where the sum is taken modulo $(k+1)q/2$. In addition, if $t > 0$,

$$f_j(C_i) = (-1)^{i+j} \text{ for } 1 \leq j \leq q \text{ and } \frac{(k+1)}{2}q \leq i \leq r-1.$$

It is easy to see that f_j is a signed star k -dominating function of G for each $1 \leq j \leq q$ and $\{f_1, f_2, \dots, f_q\}$ is a signed star k -dominating family of G . Hence $d_{kSS}(G) \geq q$. Therefore $d_{kSS}(G) = q$, as desired.

Subcase 2.2: Assume that q is even. Then $t/2 + (k+1)/2$ is even. Define the functions f_1, f_2, \dots, f_{q-1} as follows.

$$f_1(C_i) = \begin{cases} 1 & \text{if } 0 \leq i \leq \frac{(k+1)(q-2)}{4} + \frac{k+1}{2} - 2, \\ -1 & \text{if } \frac{(k+1)(q-2)}{4} + \frac{k+1}{2} - 1 \leq i \leq \frac{(k+1)}{2}(q-1) - 1, \end{cases}$$

and for $2 \leq j \leq q - 1$ and $0 \leq i \leq (k + 1)(q - 1)/2 - 1$,

$$f_j(C_i) = f_{j-1}(C_{i+(k+1)}),$$

where the sum is taken modulo $(k + 1)(q - 1)/2$. In addition,

$$f_j(C_i) = (-1)^{i+j} \text{ for } 1 \leq j \leq q \text{ and } \frac{(k+1)}{2}(q-1) \leq i \leq r-1.$$

It is easy to see that f_j is a signed star k -dominating function of G for each $1 \leq j \leq q - 1$ and $\{f_1, f_2, \dots, f_{q-1}\}$ is a signed star k -dominating family of G . Hence $d_{kSS}(G) \geq q - 1$ and so $d_{kSS}(G) = q - 1$, as desired. \square

According to Theorems 3.1, 3.3 and the following two well-known results, we can determine the signed star k -domatic number of complete graphs.

Theorem. *The complete graph K_{2r} is 1-factorable.*

Theorem. *For every positive integer r , the graph K_{2r+1} is Hamiltonian factorable.*

REFERENCES

1. M. Atapour, S. M. Sheikholeslami, A. N. Ghameshlou, and L. Volkmann, *Signed star domatic number of a graph*, Discrete Appl. Math. **158** (2010), 213–218.
2. A. N. Ghameshlou, A. Khodkar, R. Saei, and S. M. Sheikholeslami, *Signed (b, k) -edge covers in graphs*, Intelligent Information Management **2** (2010), 143–148.
3. T. W. Haynes, S. T. Hedetniemi, and P. J. Slater, *Fundamentals of domination in graphs*, Marcel Dekker, Inc., New York, 1998.
4. D. König, *Über Graphen und ihre Anwendung auf Determinantentheorie und Mengenlehre*, Math. Ann. **77** (1916), 453–465.
5. R. Saei and S. M. Sheikholeslami, *Signed star k -subdomination numbers in graphs*, Discrete Appl. Math. **156** (2008), 3066–3070.
6. C. P. Wang, *The signed star domination numbers of the Cartesian product*, Discrete Appl. Math. **155** (2007), 1497–1505.
7. C.P. Wang, *The signed b -matchings and b -edge covers of strong product graphs*, Contrib. Discrete Math. **5** (2010), 1–10.
8. D. B. West, *Introduction to graph theory*, Prentice-Hall, Inc., 2000.
9. B. Xu, *On edge domination numbers of graphs*, Discrete Math. **294** (2005), 311–316.
10. ———, *Two classes of edge domination in graphs*, Discrete Appl. Math. **154** (2006), 1541–1546.
11. B. Xu and C. H. Li, *Signed star k -domination numbers of graphs*, Pure Appl. Math. (Xi'an) **25** (2009), 638–641, in Chinese.

DEPARTMENT OF MATHEMATICS, AZARBAIJAN UNIVERSITY OF TARBIAT MOALLEM,
TABRIZ, I. R. IRAN

E-mail address: s.m.sheikholeslami@azaruniv.edu

LEHRSTUHL II FÜR MATHEMATIK, RWTH-AACHEN UNIVERSITY,
52056 AACHEN, GERMANY

E-mail address: volkm@math2.rwth-aachen.de