

# Petrology and geochemistry of Devono-Carboniferous volcanic rocks in Nova Scotia

*J. Dostal, Department of Geology  
Saint Mary's University, Halifax, N.S. Canada B3H 3C3*

*J.D. Keppie, Nova Scotia Department of Mines and Energy  
P.O. Box 1087, Halifax, N.S. Canada B3J 2X1*

*C. Dupuy, Centre Geologique et Geophysique, U.S.T.L.  
34060 Montpellier Cedex, France*

The Devono-Carboniferous volcanic rocks of Nova Scotia include Middle Devonian basalts from the McAras Brook Formation in the northern Antigonish Highlands (Ballantynes Cove and McAras Brook areas), Middle Devonian and Carboniferous basalts and rhyolites of the Fountain Lake Group in the Cobequid Highlands and the Upper Devonian-Lower Carboniferous basalts and rhyolites of the Fisset Brook Formation in Cape Breton Island. The volcanic rocks were extruded upon the continental crust in an intraplate setting. The basalts are tholeiitic except those from Ballantynes Cove which are alkaline. All these basalts could have been derived from a similar upper mantle source - garnet peridotite. Rhyolites were probably generated by crustal anatexis related to the ascending basaltic magma. The Devono-Carboniferous volcanism is probably connected with rifting along faults bounding the Magadalen pull-apart basin. In the Cobequid Highlands, the volcanism appears to be spatially and temporally associated with plutonism.

Les roches volcaniques Dévono-Carbonifères de la Nouvelle-Ecosse comprennent: les basaltes de la formation McAras Brook (Dévonian moyen) dans la partie nord des hautes-terres d'Antigonish (régions de Ballantynes Cove et de McAras Brook); les basaltes et les rhyolites du groupe Fountain Lake (Dévonian moyen et Carbonifère) des monts Cobequid; et les basaltes et rhyolites (Dévonian supérieur-Carbonifère inférieur) de la formation Fisset Brook sur l'île du Cap-Breton. Les roches volcaniques furent répandues à la surface de la croûte continentale dans un contexte intraplaque. Les basaltes sont tous tholéitiques à l'exception de ceux de Ballantynes Cove qui sont alcalins. Tous ces basaltes pourraient provenir d'une même source située dans la partie supérieure de manteau terrestre - péridotite à grenat. Les rhyolites résultent probablement de l'anatexie de la croûte associée à la montée du magma basaltique. Le volcanisme Dévono-Carbonifère est probablement relié à une fissuration le long des failles qui encadrent le bassin de déchirement des Madeleines. Dans les monts Cobequid, le volcanisme semble associé de façon spatio-temporelle au plutonisme.

[Traduit par le journal]

## INTRODUCTION

The Devonian Acadian Orogeny is one of the more significant events in the Paleozoic history of the northern Appalachians. The effects of the Acadian Orogeny extend from the Atlantic seaboard some 500 km across the trend of the Appalachians and include various degrees of deformation and metamorphism and widespread plutonism. Devono-Carboniferous volcanic rocks are widely scattered along two belts through northern New Brunswick to north-central Maine and from northeastern Massachusetts through coastal Maine to northern Nova Scotia (Ruitenberg *et al.* 1977, Osberg 1978, Keppie and Dostal 1980).

In Nova Scotia, volcanic rocks of Middle Devonian to Carboniferous age occur in the Cobequid Highlands, northern Antigonish Highlands and Cape Breton Island (Fig. 1). There is only a limited amount of information on these rocks (Keppie and Dostal 1980, Smith and MacDonald 1981, Blanchard 1982) although an understanding of their petrogenesis would contribute substantially to any tectonic model for later stages of the Acadian Orogeny. The purpose of this paper is to present some geochemical data on the Middle Devonian basalts from the McAras Brook Formation in the northern Antigonish Highlands (northern Nova Scotia, Fig. 1) and to compare these rocks to other volcanic suites from Nova Scotia containing mafic rocks (basalts) for which a Middle Devonian to Carboniferous age is well documented.

## GEOLOGICAL SETTING

The McAras Brook Formation (Williams 1914, Bell 1926, Boucot *et al.* 1974, Keppie *et al.* 1978) crops out in a few small isolated areas around the periphery of the Antigonish Highlands (Fig. 1) and is well exposed only along the coast of Northumberland Strait. The type section (MAB) occurs near Knoydart Point, west of Arisaig (Fig. 1), where continental red conglomerates and sandstones with intercalated basalt flows lie unconformably upon sedimentary rocks of Late Silurian and Early Devonian age (Boucot *et al.* 1974, Keppie *et al.* 1978). The McAras Brook Formation is, in turn, unconformably overlain by Late Visean sedimentary rocks (Keppie *et al.* 1978). The basalt flows vary in thickness from 4 to 18 m and the aggregate thickness of basalts is about 70 m.

At Ballantynes Cove, near Cape George, the McAras Brook Formation is composed of red conglomerate and about 70m of basalts in five flows, and is overlain by clastic sediments of Middle Devonian age. The basalt flows at Ballantynes Cove (BC) probably lie stratigraphically above all those exposed in the type section (Keppie *et al.* 1978). Thus, the rocks in the type section of the McAras Brook Formation are probably also of early Middle Devonian or late Early Devonian age.

Similar basalt flows occur in the Cobequid Highlands (Donohoe and Wallace 1980) in the Fountain Lake Group (FLG) (Fig. 1) which consists mainly of volcanic rocks, predominantly rhyolites with minor basalts and interbedded clastic rocks. In the Eastern Cobequid Highlands, the sequence is up to 3,000 m




Fig. 1 - Map showing the distribution of Devonian-Carboniferous volcanic rocks in Nova Scotia and indicating numbers of the samples from each locality.

thick. The FLG has been inferred to be Middle Devonian in age (Donohoe and Wallace 1982). However, Cormier (1982) has recently obtained two different Rb/Sr isochron ages ( $341 \pm 4$  Ma and  $387 \pm 2$  Ma) on these volcanics, which indicate that they are both Devonian and mid-Carboniferous.

Volcanic rocks are also present in the Late Devonian-Early Carboniferous Fisset Brook Formation (Kelly and MacKasey 1965, Cormier and Kelley 1964) in Cape Breton Island (Fig. 1). The Fisset Brook Formation (FBF) ranges in thickness from 250 m to 1,000 m of which volcanic rocks form the main part. It invariably rests unconformably upon Precambrian basement rocks and generally passes upwards into Early Carboniferous non-marine and marine sedimentary rocks. The FBF rocks are usually steeply dipping due to deformation associated with the Hercynian Orogeny. The samples were collected from the Cheticamp and Lake Ainslie areas and by Smith and Macdonald (1981) from Lowland Cove (LC) at the northern part of Cape Breton Island (Fig. 1). Smith and Macdonald (1981) showed that the section at Lowland Cove consists of a basal porphyritic rhyolite ( $\sim 75$  m thick) overlain by interbedded conglomerate, sandstone and siltstone ( $\sim 25$  m thick) followed by mafic tuffs and flows (275 m thick). Spores in the sedimentary rocks interbedded with the LC volcanic rocks are of Upper Devonian age. They are unconformably overlain by sedimentary rocks of the Horton Group (Smith and Macdonald 1981). The Fisset Brook Formation in the type section (near Cheticamp) and Lake Ainslie was extensively studied by Kelly and MacKasey (1965) and Blanchard (1982). The FBF rhyolites from the latter areas have Rb/Sr isochron ages of  $376 \pm 12$  Ma and  $370 \pm 20$  Ma, respectively (Cormier and Kelly 1964, Keppie and Smith 1978).

#### PETROGRAPHIC NOTES

The mafic volcanics of the McAras Brook Formation are usually dark greenish gray with amygdules in the upper parts of some flows. The amygdules are

mainly filled with calcite or chlorite. Calcite also commonly occurs in the form of fine veins which cut the rocks. The rocks are massive, aphanitic to porphyritic. Their primary mineral assemblage consists of plagioclase and clinopyroxene (augite) phenocrysts set in a finer-grained matrix of plagioclase, clinopyroxene and opaque minerals. Clinopyroxene phenocrysts are more abundant at Ballantynes Cove whereas plagioclase phenocrysts are predominant in basalts from the type section. Microprobe analyses of fresh plagioclase relics from the type section basalts gave compositions usually ranging from An59Ab39Or2 to An51Ab47Or2. The basalts from both sections have undergone alteration. The plagioclase is commonly saussuritized or albitized and clinopyroxene is replaced by chlorite or actinolite. The variably altered matrix also contains chlorite, calcite and quartz in cracks and small vesicles. In general, the rocks from the type section are more altered and contain fewer fresh relics of clinopyroxene than those from Ballantynes Cove.

The basalts from the Fountain Lake Group and from the Fisset Brook Formation are both composed mainly of clinopyroxene (augite) and plagioclase (andesine-labradorite) with subordinate amounts of Fe-Ti oxides. The rocks may contain phenocrysts of plagioclase enclosed in an ophitic matrix. The primary magmatic minerals of the basalts are replaced to a variable degree by secondary phases. The secondary minerals include albite, chlorite, epidote, zeolites, calcite and sericite. Amygdules are abundant particularly at the top of the sections and are filled mainly by chlorite, zeolites and calcite.

The rhyolites of the Fountain Lake Group and Fisset Brook Formation are usually porphyritic with phenocrysts of quartz and alkali feldspar set in a fine-grained matrix of quartz, potash feldspar and plagioclase with minor or accessory amounts of calcite, chlorite, Fe-Ti oxides and Fe-oxides. The petrography of the Fisset Brook Formation has been recently described by Blan-

chard (1982) for the Cheticamp-Margaree and Lake Ainslie areas and by Smith and Macdonald (1981) for the Lowland Cove area.

#### ANALYTICAL NOTES

The major elements were determined by X-ray fluorescence and supplementary techniques. Analyses of Zr, Nb, Y, Rb, Sr, Ba, La and Ce were also carried out by X-ray fluorescence whereas rare-earth elements (REE) were determined by instrumental neutron activation. The precision and accuracy of the trace element data were given by Dostal and Capedri (1979) and Dupuy *et al.* (1979). The whole rock analyses selected from a suite of 80 analyzed samples are given in Tables 1 and 2. Analyses of clinopyroxene were done using an energy dispersive microprobe. The average compositions of the clinopyroxenes are given in Table 3. The individual analyses of whole-rocks and clinopyroxenes and locations of the samples are deposited in the Open File Library of the Nova Scotia Department of Mines and Energy. Analyzed whole rock samples include 38 basalts from the McAras Brook Formation (19 from the type section and 19 from Ballantynes Cove), 20 samples from the Fisset Brook Formation (7 basalts and 2 rhyolites from Lowland Cove, 9 basalts from the type section and 2 rhyolites from Lake Ainslie), and 7 basalts and 15 rhyolites from the Fountain Lake Group (Fig. 1). Sampling was restricted to these areas because they are the only well dated Devonian-Carboniferous volcanic rocks. Other presumed Devonian-Carboniferous volcanic rocks were not sampled as their significance to tectonic modelling would be difficult to assess or they are covered by other work (Blanchard 1982).

#### GEOCHEMISTRY

The chemical composition of the Devonian-Carboniferous volcanic rocks of Nova Scotia confirms the petrographic observations of the predominance of basalts and rhyolites. The volcanic rocks of the McAras Brook Formation from both the

type section and Ballantynes Cove are basalts with  $\text{SiO}_2 < 55\%$  (on a volatile-free basis). Volcanic suites from the Fisset Brook Formation and Fountain Lake Group are bimodal with respect to  $\text{SiO}_2$  and are composed usually of basalts and rhyolites, although the rare occurrence of intermediate rocks has also been reported (e.g. Smith and Macdonald 1981, Blanchard 1982).




Fig. 2 - CaO-K<sub>2</sub>O-Na<sub>2</sub>O diagram for some Devonian-Carboniferous basalts of Nova Scotia. ● - type section of McAras Brook Formation; x - Ballantynes Cove, ▲ - Fountain Lake Group, o - Lowland Cove.

#### Alteration

All the analyzed samples have been affected by secondary processes which modified their chemical composition. The rocks have variable but high  $\text{H}_2\text{O}$  contents and a low  $\text{FeO}/\text{Fe}_2\text{O}_3$  ratio. In addition, some samples are high in  $\text{CO}_2$ . Among the studied volcanic rocks, the effects of alteration on the chemical compositions are most pronounced in MAB basalts, which have up to 4.3% of  $\text{H}_2\text{O}^+$ . In this section, the variable degree of oxidation is reflected by the wide range of  $\text{FeO}/\text{Fe}_2\text{O}_3$  ratios (0.07 - 1.84). This ratio is the lowest in the basal flow where it is accompanied by abundant hematite in thin section. The effects of secondary processes are also shown by variations of  $\text{Na}_2\text{O}$ ,  $\text{K}_2\text{O}$  and  $\text{CaO}$  (Fig. 2). On average, the basalts are enriched in  $\text{Na}_2\text{O}$  and depleted in  $\text{CaO}$ . The in-

verse variations of  $\text{Na}_2\text{O}$  and  $\text{CaO}$  which are reflected in thin section by albitization of plagioclase is particularly clearly displayed by MAB rocks.

Internal consistencies and overall systematic variations in basalts from BC, LC and FLG show that the composition of these rocks has been less modified during alteration than those from MAB. The composition of the rhyolites also seems to have been affected by secondary processes. However, in spite of alteration, the original geochemical characteristics of the rocks and the magmatic variation trends are still relatively well preserved (Tables 1 and 2).

#### *Basalts*

Major elements. - The mafic rocks have  $\text{SiO}_2$  contents (on a volatile-free basis) ranging from 47 to 55% and  $\text{FeO}_{\text{tot}}/\text{MgO}$  ratios varying from 0.8 to  $>3$ . Most analyzed samples from Ballantynes Cove are nepheline normative, however those from the other areas are mainly Ol- and/or Hy-normative. For a given  $\text{FeO}_{\text{tot}}/\text{MgO}$  ratio, the basalts of Ballantynes Cove have higher contents of  $\text{TiO}_2$  and  $\text{P}_2\text{O}_5$  and lower  $\text{SiO}_2$  and  $\text{Al}_2\text{O}_3$  than those from the other areas. They also have relatively uniform composition with only limited chemical variations ( $\text{FeO}_{\text{tot}}/\text{MgO} = 0.8-1.6$ ). On the other hand, the basalts from the other localities display a large range of  $\text{FeO}_{\text{tot}}/\text{MgO}$  ratios which positively correlate with  $\text{TiO}_2$  and  $\text{P}_2\text{O}_5$ . The increase of  $\text{TiO}_2$  with the  $\text{FeO}_{\text{tot}}/\text{MgO}$  ratio is typical of tholeiitic suites. The presence of rocks with high  $\text{TiO}_2$  contents and  $\text{FeO}_{\text{tot}}/\text{MgO}$  ratios among the tholeiitic basalts indicates that these basaltic suites underwent extensive low-pressure fractional crystallization. Compared to typical calc-alkaline high alumina basalts, the studied basalts are higher in  $\text{TiO}_2$ ,  $\text{FeO}_{\text{tot}}$  and  $\text{MgO}$  but lower in  $\text{SiO}_2$ .

Trace elements - Since secondary processes frequently affect the distribution of many trace elements most of the following discussion deals with the ele-

ments which are generally considered to be rather immobile and the samples selected were the least altered from the collected suites.

The abundances of trace elements corroborate the differences, shown by major elements, in the basalt composition between Ballantynes Cove and the other localities. For a given  $\text{FeO}_{\text{tot}}/\text{MgO}$  ratio, the BC basalts are higher in light REE (LREE), Zr, and Nb. In fact the abundances of these elements in the BC basalts are characteristic of alkali rocks. Their REE patterns (Fig. 3) are strongly fractionated with LREE enrichment and gradual depletion of heavy REE (HREE). Their La/Yb ratio is  $\sim 30$  and is also typical of alkali basalts.

The REE patterns of the basalts from the other localities show moderate LREE enrichment and a smaller degree of fractionation ( $\text{La}/\text{Yb} \sim 4-10$ ) than the BC rocks. The patterns resemble those of continental tholeiites or tholeiites dredged from the Atlantic Ocean near  $43^\circ\text{N}$  (Shibata et al. 1979). However the relative enrichment of incompatible elements and the inter-element relationships such as  $\text{TiO}_2$  vs Zr (Fig. 4) show that all the Devonian-Carboniferous basalts are of the within-plate type, as previously suggested by Keppie and Dostal (1980), and later for FBF basalts by Smith and Macdonald (1981) and Blanchard (1982).

#### *Rhyolites*

On the Ab-Or-Q diagram (Dostal et al. 1983) the scatter of the Devonian-Carboniferous silicic rocks is largely due to alteration, although most of them lie within the granitic field of Winkler and Von Platten (1961). Compared to recent rhyolite sequences (Ewart and Stipp 1968, Zielinski et al. 1977, Dostal et al. 1982, Cameron and Hanson 1982), they are lower in  $\text{CaO}$  and have a relatively high  $\text{K}_2\text{O}/\text{Na}_2\text{O}$  ratio. The studied rhyolites show large variations in the abundances of lithophile elements. The LC rocks are distinctly higher in LREE, Zr and Nb than the other rhyolites. The REE patterns of rhyolites from FLG show

Table 1  
CHEMICAL COMPOSITION OF REPRESENTATIVE SAMPLES FROM THE McARAS BROOK FORMATION

	TYPE SECTION						BALLANTYNES COVE						
	1	2	3	4	5	6	7	8	9	10	11	AV.	s.d.
SiO <sub>2</sub>	50.23	49.91	46.02	47.44	45.38	46.55	42.74	42.98	44.52	46.70	47.18	48.63	1.62
TiO <sub>2</sub>	1.40	1.15	1.84	1.65	2.36	2.31	1.49	1.44	1.46	1.57	1.54	1.70	0.14
Al <sub>2</sub> O <sub>3</sub>	16.73	16.56	17.20	16.27	15.87	16.63	14.71	13.95	13.08	13.95	14.59	15.38	0.70
Fe <sub>2</sub> O <sub>3</sub>	3.28	2.98	11.74	9.37	10.23	6.77	5.66	8.03	9.75	9.73	6.26		
FeO	3.47	2.96	1.07	0.99	3.84	3.26	3.14	2.62	0.49	1.04	2.62	10.12*	0.54
MnO	0.50	0.55	0.75	1.14	0.44	0.14	0.27	0.24	0.18	0.17	0.20	0.21	0.04
MgO	7.39	7.30	5.84	4.38	4.18	3.52	9.83	9.43	8.66	7.98	6.16	8.64	1.27
CaO	9.53	8.92	5.45	6.45	8.12	9.68	9.69	9.71	8.83	8.51	8.71	9.80	1.08
Na <sub>2</sub> O	2.83	3.46	4.92	5.35	3.33	4.82	3.08	2.27	3.74	4.50	3.91	3.67	0.88
K <sub>2</sub> O	0.41	0.74	0.21	0.27	1.31	0.55	0.80	0.79	0.54	0.71	0.96	1.11	0.43
P <sub>2</sub> O <sub>5</sub>	0.20	0.21	0.35	0.37	0.80	0.81	0.74	0.86	0.72	0.69	0.75	0.74	0.07
H <sub>2</sub> O <sup>+</sup>	2.96	3.60	3.12	3.30	1.59	3.29	4.08	4.05	4.10	3.56	3.12		
H <sub>2</sub> O <sup>-</sup>	0.84		0.47		1.24	0.93	1.02			0.44	0.87		
CO <sub>2</sub>	0.43	0.56	1.50	2.60	0.19	0.57	1.38	2.18	2.03	0.88	2.16		
Σ	100.29	98.90	100.48	99.58	98.88	99.83	98.63	98.55	98.10	100.43	99.13	100.00	
Rb(ppm)	21	13	2	6	31	16	62	48	41	61	65	58	8
Sr	295	318	739	870	683	860	2749	2445	2119	2983	2674	2485	442
Ba	272	342	252	1238	582	417	924	885	630	1047	840	1085	335
La	11	12.9	21	15.5	31	26	47	56.1	53.0	52	50.5	48	6
Ce	26	29.8	40	33.5	64	59	110	114.0	113.0	118	113.0	105	16
Nd		16.2						49.3	46.2		47.9		
Sm		4.41		5.18				8.53	7.57		7.98		
Eu		1.43		1.86				2.42	2.32		2.19		
Tb		0.89		1.01				0.82	0.81		0.81		
Yb		3.13		3.17				1.75	1.72		1.66		
Lu		0.48		0.48				0.26	0.26		0.25		
Zr	139	155	172	179	260	255	249	245	226	256	229	246	19
Nb	12	13	14	15	21	21	21	21	20	21	21	21	2

AV.=average of 19 basalt samples from Ballantynes Cove recalculated to 100% (on volatile-free basis); s.d.=standard deviation for the average; \*-total Fe as FeO.




Fig. 3 - A - Chondrite-normalized REE abundances of basalts of McAras Brook Formation from Ballantynes Cove (● - sample 8 and x - sample 9) and from the type section (+ - sample 4 and o - sample 2). B - Chondrite-normalized REE abundances of basalts from the Fisset Brook Formation in Lowland Cove. (o - sample 13 and x - sample 14) from the Fountain Lake Group (+ - sample 20 and ● - sample 19) and from the McAras Brook

LREE enrichment with a La/Yb ratio of about 5 and a small negative Eu anomaly (Fig. 3). The LC rhyolites have more fractionated REE patterns with a La/Yb ratio >10 and a distinct negative Eu anomaly.

#### Clinopyroxene

The average compositions of clinopyroxene from MAB, BC and LC basalts are given in Table 3. The clinopyroxene from the BC rocks are augites with a compositional range of Wo 42.0-47.6 En 38.8-47.4 Fs 9.8-14.0. The relics of clinopyroxene, both phenocrysts and in groundmass also show distinct chemical variations in Ti, Al and Na. Titanium displays a positive correlation with the Fe/Mg ratio whereas SiO<sub>2</sub> correlates negatively with Al<sub>2</sub>O<sub>3</sub> and TiO<sub>2</sub>. The rims of some augite crystals are higher in Ti, Al and Fe and lower in Ca than their cores.

Although clinopyroxene of the MAB basalts is also augite, there are significant differences in the compositions of this mineral from the two sections of the McAras Brook Formation. On the average, clinopyroxenes from MAB basalts are higher in SiO<sub>2</sub>, FeO<sub>tot</sub> and MnO and lower in Al<sub>2</sub>O<sub>3</sub> and CaO than those from BC rocks. They also have a lower Al<sup>iv</sup> and Wo component but a higher Fs component with a composition of Wo 37.4-42.7 En 39.8-42.8 Fs 14.9-22.1. MAB pyroxenes also do not display covariation of TiO<sub>2</sub> vs FeO<sub>tot</sub>/MgO and of SiO<sub>2</sub> vs TiO<sub>2</sub>. Compared to the average compositions of clinopyroxenes from various rock types, MAB clinopyroxenes have very low abundances of Al<sub>2</sub>O<sub>3</sub> (Table 3). Clinopyroxene from LC basalts are similar to those from MAB.

Formation type section (□ - sample 4). C - Chondrite-normalized REE abundances of rhyolites from the Fisset Brook Formation in Lowland Cove (+ - sample 16 and ● - sample 15) and in the Lake Ainslie area (o - sample 17) and from the Fountain Lake Group (x - sample 26). Chondritic values from Frey et al. (1968).



Table 2

## CHEMICAL COMPOSITION OF REPRESENTATIVE VOLCANIC ROCKS FROM THE FISSET BROOK FORMATION AND FOUNTAIN LAKE GROUP

	FISSET BROOK FORMATION						FOUNTAIN LAKE GROUP									
	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	
SiO <sub>2</sub>	49.70	49.10	48.70	73.50	74.90	70.79	45.80	45.70	46.55	46.13	45.80	75.44	76.41	73.35	79.56	
TiO <sub>2</sub>	1.70	2.77	2.86	0.35	0.28	0.30	1.25	2.26	2.45	2.42	3.02	0.20	0.27	0.18	0.10	
Al <sub>2</sub> O <sub>3</sub>	15.90	13.50	13.60	12.40	11.70	14.24	16.20	15.61	16.14	15.50	14.80	13.57	11.05	13.63	10.73	
Fe <sub>2</sub> O <sub>3</sub>	5.72	8.06	9.00	2.46	2.24	0.45	6.54	8.13	9.14	5.35	6.72	1.09	2.47	0.86	0.55	
FeO	5.65	6.65	5.70	1.10	0.85	1.45	2.95	4.69	4.64	7.59	8.25	1.25	1.25	0.67	0.87	
MnO	0.23	0.36	0.24	0.08	0.05	0.03	0.42	0.16	0.24	0.20	0.28	0.04	0.04	0.01	0.02	
MgO	5.36	5.37	4.80	0.46	0.11	0.50	8.27	6.02	4.75	5.62	5.23	1.57	0.28	0.17	0.22	
CaO	7.69	4.98	7.56	0.59	0.63	0.14	10.90	10.13	6.78	9.13	9.48	0.27	0.45	0.21	0.35	
Na <sub>2</sub> O	3.42	3.16	2.92	3.21	2.03	1.57	2.47	2.48	2.48	2.61	2.73	0.00	3.42	2.15	2.72	
K <sub>2</sub> O	1.17	1.80	1.76	4.90	6.60	10.00	0.23	0.62	2.34	1.45	0.68	4.82	4.26	7.94	4.43	
P <sub>2</sub> O <sub>5</sub>	0.47	0.81	0.79	0.04	0.02	0.04	0.19	0.42	0.41	0.33	0.59	0.04	0.02	0.03	0.05	
H <sub>2</sub> O <sup>F</sup>	2.20	3.65	2.55	0.80	0.40	0.79	3.10	3.01	2.73	2.50	2.55	1.96	0.21	0.69	0.52	
H <sub>2</sub> O <sup>-</sup>				0.00	0.02		0.65				0.00					
CO <sub>2</sub>	0.05	0.10	0.10	0.20	0.35	0.17	0.80	0.70	0.48	0.85	0.05	0.46	0.26	0.00	0.27	
Σ	99.26	100.31	100.58	100.09	100.18	100.47	99.77	99.93	99.13	99.68	100.18	100.71	100.39	99.89	100.38	
Rb(ppm)	37	46	38	115	124	240	12	24	74	55	19	201	168	202	120	
Sr	306	202	309	54	70	103	323	285	305	386	259	30	80	167	138	
Ba	342	519	584	267	297	1459	190	205	604	408	207	865	86	536	435	
La	20	30.8	29.1	138	144	13.8	7	17.1	22.5	18	22	37	99	32	47.4	
Ce	48	71.6	71.3	291	293	33.1	16	42.5	54.0	46	53	84	237	69	113	
Nd		38.5	40.3			15.2		24.5	29.4						52.3	
Sm		8.92	9.01	24.1	23.8	3.37		6.17	6.84						11.4	
Eu		2.77	2.93	1.61	1.09	0.68		2.24	2.34						0.46	
Tb		1.58	1.67	3.76	3.45	0.65		1.26	1.31						2.55	
Yb		4.55	4.67	11.8	10.6	2.77		3.22	3.48						8.43	
Lu		0.70	0.73	1.89	1.68	0.52		0.48	0.54						1.23	
Zr	203	278	271	761	689	211	100	225	248	186	253	171	1264	216	214	
Nb	17	22	22	37	33	16	9	19	21	16	22					

DOSTAL, KEPPIE and DUPUY

FISSET BROOK FORMATION: Lowland Cove: samples 12-16 (major elements taken from Smith and Macdonald, 1981); Lake Ainslie: sample 17. FOUNTAIN LAKE GROUP: Devonian: samples 18-24, Carboniferous: 25-26.



Compositions of clinopyroxene relics in the basalts confirm their whole-rock geochemical characteristics and the inferred paleotectonic settings of the host lavas. On the  $\text{SiO}_2$  vs  $\text{Al}_2\text{O}_3$  diagram (Fig. 5), the pyroxenes from MAB and LC volcanics fall into the sub-alkaline field of Le Bas (1962) whereas those from BC basalts plot predominantly into the alkaline field. On the discriminant diagrams of Leterrier *et al.* (1982) the pyroxenes from MAB and LC are characteristic of non-orogenic tholeiitic basalts whereas clinopyroxenes from BC seem to be mainly related to alkaline series.

#### PETROGENESIS

With the exception of rocks from Ballantynes Cove, the analyzed Devonian-Carboniferous basalts are continental tholeiites. The metavolcanics from Ballantynes Cove, which have higher contents of relatively immobile lithophile elements (such as LREE, Zr, Nb) for a comparable  $\text{FeO}_{\text{tot}}/\text{MgO}$  ratio, resemble alkali basalts. The basalts underwent variable degrees of low-pressure fractional crystallization which changed the abundances of many elements. However, some element ratios including Ti/Zr,

Zr/Nb and the shape of the REE patterns change little during such a process and thus they are suitable for evaluating the origin of these rocks, particularly as these elements probably were not affected by secondary processes. Similarities of element ratios such as Zr/Nb among the analyzed basalts suggest a genetic relation between the tholeiitic and alkali basalts, in particular, their derivation from a similar source. Kay and Gast (1973) and Frey *et al.* (1978) have shown that alkali basalts, with highly fractionated REE patterns such as those of the BC basalts can be generated by a small degree of partial melting of upper mantle garnet peridotite with chondritic-relative REE abundances. The model calculations suggest that this mechanism can also produce liquids with the geochemical characteristics of the parental magmas of metabasalts from Ballantynes Cove. Subsequent low-pressure fractional crystallization of such magmas can form melts with the compositions of the BC rocks.

Compared to the BC alkali basalts, the other continental tholeiitic basalts have lower LREE but higher HREE abundances. These rocks could be derived from a similar parental source as the




Fig. 4 - A -  $\text{TiO}_2$ -Zr diagram of Pearce *et al.* (1981) for the basalts of the McAras Brook Formation (o - Ballantynes Cove; x - type section). B -  $\text{TiO}_2$ -Zr diagram of Pearce *et al.* (1981) for the basalts from the Fourn-tain Lake Group (x) and the Fisset Brook Formation at Lowland Cove (●) and the type section and Lake Ainslie (o).




Fig. 5 - Variations of  $\text{SiO}_2$  vs  $\text{Al}_2\text{O}_3$  in clinopyroxenes from the basalts of A-Ballantynes Cove, B-type section of the McAras Brook Formation and C-Fisset Brook Formation at Lowland Cove. Fields of pyroxenes from: S-Subalkaline rocks, A-Alkaline rocks, and P-Peralkaline rocks (after Le-Bas 1962).

alkali basalts but such a process would involve a significantly higher degree of partial melting of upper mantle peridotite containing a phase that preferentially enriches HREE, for example, garnet. In fact, the process involving variable degrees of partial melting of garnet-bearing upper mantle has been frequently invoked to elucidate the genetic relation between associated alkali and tholeiitic basalts (e.g. Frey *et al.* 1974, Shibata *et al.* 1979).

The two most frequently invoked processes for the genesis of rhyolitic rocks are fractional crystallization of basaltic or andesitic magma and partial melting of crustal rocks (Ewart and Stipp 1968). In the Devonian-Carboniferous volcanic rocks there is some evidence that rhyolites might have been produced by differentiation of a basic magma (e.g. high contents of some incompatible elements in LC rhyolites accompanied by a distinct negative Eu anomaly in their REE pattern). However, the lack of intermediate rocks argues against this model. In fact, as recently demonstrated by Cameron and Hanson (1982) it is difficult, on the basis of geochemistry alone, to discriminate between these two

processes. The close spatial and temporal associations of rhyolites and basalts suggest that these rocks are genetically related. However, the overall chemical similarity of the felsic rocks to recent calc-alkali rhyolites suggest that a more plausible mechanism for their origin is by partial melting of the crustal rocks. This process may also account for diversity in the composition of rhyolites from different areas and the lack of intermediate rocks. The crustal anatexis was probably related to the ascending basaltic magma.

#### CONCLUSION

The geochemical results presented here are consistent with the geological interpretation of these volcanic rocks as part of Devonian-Carboniferous intra-plate continental molasse deposits (Keppie and Dostal 1980) formed during the transpression stage of the Acadian Orogeny (Keppie 1982a) in an intramontane or pull-apart basin centered about the Magdalen Islands (Keppie 1982b). The development of the Magdalen pull-apart basin immediately post-dated the compressive phase of the Acadian Orogeny.

Table 3

## AVERAGE COMPOSITION OF CLINOPYROXENES

	BC	MAB	LC	WPT	WPA	VAB
SiO <sub>2</sub>	49.39 (1.01)	51.36 (0.45)	50.60 (0.70)	51.4 (1.1)	49.2 (2.4)	50.7 (1.5)
TiO <sub>2</sub>	1.02 (0.30)	1.16 (0.21)	1.36 (0.28)	0.94 (0.46)	1.77 (1.20)	0.67 (0.37)
Al <sub>2</sub> O <sub>3</sub>	5.62 (0.89)	1.75 (0.42)	1.90 (0.52)	3.2 (1.4)	4.1 (2.7)	3.3 (1.8)
FeO*	7.16 (0.60)	10.81 (1.22)	11.96 (1.31)	11.5 (3.5)	11.3 (5.5)	9.1 (4.5)
MnO	0.15 (0.06)	0.30 (0.10)	0.30 (0.08)	0.42 (0.40)	0.43 (0.38)	0.26 (0.15)
MgO	14.46 (0.80)	14.46 (0.39)	14.13 (0.74)	14.8 (3.1)	10.9 (3.8)	15.3 (1.4)
CaO	21.21 (0.86)	19.71 (0.79)	19.49 (0.59)	17.1 (3.5)	20.5 (2.8)	19.8 (3.2)
Na <sub>2</sub> O	0.34 (0.11)	0.36 (0.11)	0.24 (0.10)	0.47 (0.19)	1.17 (1.36)	0.35 (0.17)
n	42	20	70	26	82	73

BC - pyroxenes from basalts of Ballantynes Cove; MAB - pyroxenes from basalts of McAras Brook type section; LC - pyroxenes from basalts of Lowland Cove; WPT - pyroxenes from within-plate tholeiites (Nisbet and Pearce, 1977); WPA - pyroxenes from within-plate alkalic basalts (Nisbet and Pearce, 1977); VAB - volcanic arc basalts (Nisbet and Perce, 1977). Standard deviations are given in parenthesis; FeO\* = total Fe recalculated as FeO; n = number of analyses.

which produced folds and an attendant cleavage in Emsian and older rocks. It was accompanied by rifting that led to extensive volcanism along the faults bounding the basin. The Devonian-Carboniferous volcanic rocks were emplaced mainly around the southern and eastern margins of the basins. Active rifting appears to have occurred in three main pulses: Middle Devonian, Tournaisian (straddling the Devonian-Carboniferous boundary) and mid-Carboniferous and were separated by periods of passive sedimentation. In the Cobequid Highlands, some of the volcanism appears to be spatially and temporally related to plutonism, as both yield similar 330-340 Ma Rb/Sr whole-rock isochron ages and are juxtaposed (Donohoe and Wallace 1982, Cormier 1982). This geological setting is very similar to the Mount Pleasant area in southern New Brunswick suggesting interesting economic possibilities.

## ACKNOWLEDGEMENTS

The study was supported by grants from NSERC Canada (operating grant A3782) the Faculty of Science of Saint Mary's

University, and the Mineral Resources Division of Nova Scotia Department of Mines and Energy. The authors thank Dr. D. B. Clarke for permitting access to microprobe facilities at Dalhousie University and Dr. A.S. Macdonald and P.K. Smith for allowing use of their samples for further analysis. We would also like to thank Drs. J.B. Murphy, R.P. Raeside and S.M. Barr for critical comments.

BELL, W.A. 1926. Carboniferous formations of Northumberland Strait, Nova Scotia. Geological Survey of Canada, Summary Report 1924, part C, pp. 142-180.

BLANCHARD, M.C. 1982. Geochemistry and petrology of the Fisset Brook Formation, Western Cape Breton Island, Nova Scotia. M.Sc. thesis, Dalhousie University, 198p.

BOUCOT, A.J., DEWEY, J.F., DINELEY, D. L., FLETCHER, R., FYSON, W.K., GRIFFIN, J.G., HICKOX, C.F., MCKERROW, W.S. and ZIEGLER, A.M. 1974. Geology of the Ariesaig area Antigonish County, Nova Scotia. Geological Society of America, Special Paper 139, 191p.

- CAMERON, K.L. and HANSON, G.N. 1982. Rare earth element evidence concerning the origin of voluminous mid-Tertiary rhyolitic ignimbrites and related volcanic rocks, Sierra Madre Occidental, Chihuahua, Mexico. *Geochimica et Cosmochimica Acta*, 46, pp. 1489-1503.
- CORMIER, R.F. 1982. Rb/Sr age data for the Fountain Lake Group volcanics. Nova Scotia Department of Mines and Energy. Reports of Activities 1981, p. 14.
- CORMIER, R.F. and KELLY, A.M. 1964. Absolute age of the Fisset Brook Formation and the Devonian-Mississippian boundary, Cape Breton Island, Nova Scotia. *Canadian Journal of Earth Sciences*, 1, pp. 159-166.
- DONOHUE, H.V. Jr. and WALLACE, P.I. 1980. Structure and stratigraphy of the Cobequid Highlands, Nova Scotia. Geological Association of Canada, Halifax 80; Field Trip 19, 64p.
- DONOHUE, H.V. Jr. and WALLACE, P. I. 1982. Geological Maps of the Cobequid Highlands. Nova Scotia Department of Mines and Energy, Maps 82-6, 7, 8, 9. Scale 1:50,000.
- DOSTAL, J. and CAPEDEI, S. 1979. Rare-earth elements in high-grade metamorphic rocks from the western Alps. *Lithos*, 12, pp. 41-49.
- DOSTAL, J., COULON, C. and DUPUY, C. 1982. Cainozoic andesitic rocks of Sardinia (Italy). In Thorpe, R.S. (Editor). *Andesites: Orogenic Andesites and Related Rocks*. John Wiley and Sons, pp. 353-370.
- DOSTAL, J., DUPUY, C. and KEPPIE, J.D. 1983. Uranium and thorium in Paleozoic rhyolites of Nova Scotia. *Canadian Journal of Earth Sciences*, 20, pp. 266-274.
- DUPUY, C., DOSTAL, J. and COULON, C. 1979. Geochemistry and origin of andesitic rocks from North-Western Sardinia. *Journal of Volcanology and Geothermal Research*, 6, pp. 375-389.
- EWART, A. and STIPP, J.J. 1968. Petrogenesis of the volcanic rocks of the central North Island, New Zealand as indicated by a study of  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios, and Sr, Rb, K, V and Th abundances. *Geochimica et Cosmochimica Acta*, 32, pp. 699-735.
- FREY, F.A., HASKIN, M.A., POETZ, J. and HASKIN, L.A. 1968. Rare earth abundances in some basic rocks. *Journal of Geophysical Research*, 73, pp. 6085-6098.
- FREY, F.A., BRYAN, W.B. and THOMPSON, G. 1974. Atlantic Ocean floor: Geochemistry and petrology of basalts from Legs 2 and 3 of the Deep-Sea Drilling Project. *Journal of Geophysical Research*, 79, pp. 5507-5527.
- FREY, F.A., GREEN, D.H. and ROYS, D. 1978. Integrated models of basalt petrogenesis: A study of quartz tholeiites to olivine melilitites from South Eastern Australia utilizing geochemical and experimental petrological data. *Journal of Petrology*, 19, pp. 463-513.
- KAY, R., and GAST, P.W. 1973. The rare-earth content and origin of alkali-rich basalt. *Journal of Geology*, 81, pp. 653-682.
- KELLEY, D.G. and MACKASEY, W.O. 1965. Basal Mississippian volcanic rocks in Cape Breton Island, N.S. Geological Survey of Canada Paper 64-34, 10p.
- KEPPIE, J.D. 1982a. Tectonic map of Nova Scotia. Nova Scotia Department of Mines and Energy. Scale 1:500,000.
- KEPPIE, J.D. 1982b. The Minas Geofracture. In St. Julien, P. and Beland, J. (Editors). Geological Association of Canada Special Paper 24, pp. 263-280.
- KEPPIE, J.D. and SMITH, P.K. 1978. Compilation of isotopic age data of Nova Scotia. Nova Scotia Department of Mines Report 78-4.
- KEPPIE, J.D., GILES, P.S. and BOEHNER, R.S. 1978. Some Middle Devonian to Lower Carboniferous rocks of Cape George, Nova Scotia. Nova Scotia Department of Mines, Paper 78-4, 37p.

- KEPPIE, J.D. and DOSTAL, J. 1980. Paleozoic volcanic rocks of Nova Scotia. In Wones, R. (Editor). Caledonides in the U.S.A. International Geological Correlation Program, 27(a), pp. 249-256.
- LE BAS, M.J. 1962. The role of aluminium in igneous clinopyroxenes with relation to their parentage. *American Journal of Science*, 260, pp. 267-288.
- LETERRIER, J., MAURY, R.C., THONON, P., GIRARD, D. and MARCHAL, M. 1982. Clinopyroxene composition as a method of identification of the magmatic affinities of paleovolcanic series. *Earth and Planetary Science Letters*, 59, pp. 138-154.
- NISBET, E.G. and PEARCE, J.A. 1977. Clinopyroxene composition in mafic lavas from different tectonic settings. *Contributions to Mineralogy and Petrology*, 63, pp. 149-160.
- OSBERG, P. 1978. Synthesis of the geology of the north-eastern Appalachians (U.S.A.). Geological Survey of Canada Paper 78-13, 27p.
- PEARCE, J.A., ALABASTER, T., SHELTON, A.W. and SEARLE, M.P. 1981. The Oman ophiolite as a Cretaceous arc-basin complex: evidence and implications. *Philosophical Transactions, Royal Society A300*, pp. 299-317.
- RUITENBERG, A.A., FYFFE, L.R., McCUTCHEON, S.R., ST. PETER, C.R., D'ARINKI, R.R. and VENUGOPAL, D.V. 1977. Evolution of Pre-Carboniferous tectonostratigraphic zones in the New Brunswick Appalachians. *Geoscience Canada*, 4, pp. 171-181.
- SHIBATA, T., THOMPSON, G. and FREY, F.A. 1979. Tholeiitic and alkali basalts from the Mid-Atlantic ridge at 43°N. *Contributions to Mineralogy and Petrology*, 70, pp. 127-142.
- SMITH, P.K. and MacDONALD, A.S. 1981. The Fisset Brook Formation at Lowland Cove, Inverness County, Nova Scotia. Nova Scotia Department of Mines and Energy Paper 81-1, 18p.
- WILLIAMS, M.Y. 1914. Arisaig-Antigonish District, Nova Scotia. Geological Survey of Canada, Memoir 60, 173p.
- WINKLER, H.G.F. and VON PLATEN, H. 1961. Experimentelle Gesteinsmetamorphose. V. Experimentelle anatektischer Schmelzen und ihre petrogenetische Bedeutung. *Geochimica et Cosmochimica Acta*, 24, pp. 250-259.
- ZIELINSKI, R.A., LIPMAN, P.W. and MILLARD, H.T. 1977. Minor element abundances in obsidian, perlite and felsite of calc-alkalic rhyolites. *American Mineralogist*, 62, pp. 426-437.

Reviewers: J.B. Murphy  
R.P. Raeside