

# Rose-Hulman Undergraduate Mathematics Journal

---

Volume 14  
Issue 1

Article 5

---

## Trees of Irreducible Numerical Semigroups

Taryn M. Laird

*Northern Arizona University*, [tml94@nau.edu](mailto:tml94@nau.edu)

Jose E. Martinez

*Northern Arizona University*

Follow this and additional works at: <https://scholar.rose-hulman.edu/rhumj>

---

### Recommended Citation

Laird, Taryn M. and Martinez, Jose E. (2013) "Trees of Irreducible Numerical Semigroups," *Rose-Hulman Undergraduate Mathematics Journal*: Vol. 14 : Iss. 1 , Article 5.

Available at: <https://scholar.rose-hulman.edu/rhumj/vol14/iss1/5>

ROSE-  
HULMAN  
UNDERGRADUATE  
MATHEMATICS  
JOURNAL

# TREES OF IRREDUCIBLE NUMERICAL SEMIGROUPS

Taryn M. Laird and José E. Martinez <sup>a</sup>

VOLUME 14, No. 1, SPRING 2013

Sponsored by

Rose-Hulman Institute of Technology

Department of Mathematics

Terre Haute, IN 47803

Email: [mathjournal@rose-hulman.edu](mailto:mathjournal@rose-hulman.edu)

<http://www.rose-hulman.edu/mathjournal>

---

<sup>a</sup>Northern Arizona University

# TREES OF IRREDUCIBLE NUMERICAL SEMIGROUPS

Taryn M. Laird and José E. Martinez

**Abstract.** A 2011 paper by Blanco and Rosales describes an algorithm for constructing a directed tree graph of irreducible numerical semigroups of fixed Frobenius numbers. This paper will provide an overview of irreducible numerical semigroups and the directed tree graphs. We will also present new findings and conjectures concerning the structure of these trees.

---

**Acknowledgements:** We would like to give a special thanks to our mentor, Jeff Rushall, for his help and guidance, and to Ian Douglas for sharing his expertise in programming.

## 1 Introduction

There are many problems in which we unknowingly encounter numerical semigroups in mathematics. Two of the most well known examples of these problems are the *coin problem* and the *postage stamp problem*. Here is one of many such puzzles.

**Example 1.1.** Suppose we are given an infinite number of 4 cent, 6 cent, and 9 cent coins. We would like to determine the largest amount of exact change that we cannot make using combinations of these coins.

In order to find this maximum number, we must consider all possible combinations of the coins. That is,

$$\begin{aligned} \{\text{all possible amounts of change}\} &= \{4k_1 + 6k_2 + 9k_3 \mid k_1, k_2, k_3 \in \mathbb{N}\} \\ &= \{0, 4, 6, 8, 9, 10, 12, 13, 14, 15, 16, \dots\} \end{aligned}$$

Since we have made 4 consecutive integer amounts (12 cents, 13 cents, 14 cents and 15 cents) we know that all integer amounts greater than 15 cents can be made by adding more 4 cent coins to these consecutive amounts. This means that the largest amount of exact change we cannot make is 11 cents.

The next problem is similar to Example 1.1, but it has an important difference.

**Example 1.2.** Suppose we are given an infinite number of 4 cent and 14 cent coins and we again wish to find the largest amount of change that we cannot make with these coins.

We again consider all possible combinations of the coins:

$$\begin{aligned} \{\text{all possible amounts of change}\} &= \{4k_1 + 14k_2 \mid k_1, k_2 \in \mathbb{N}\} \\ &= \{0, 4, 8, 12, 14, 16, 18, \dots\} \end{aligned}$$

Note that we can never make an odd amount of change with our coins. This means that there cannot be a largest un-makeable amount.

The infinitely large set of change amounts in the first example is called a numerical semigroup, and the largest uncreatable number is called the Frobenius number of the numerical semigroup.

We now ask if there are other sets of coin denominations that have a largest un-makeable amount of 11 cents. It turns out that other such sets of denominations do share the same largest unmakeable amount and a special subset of the corresponding numerical semigroups can be recursively constructed in a directed tree graph.

In the following sections of the paper we will first define numerical semigroups and certain properties that are important to both the numerical semigroups themselves and the directed tree graphs. We will then briefly examine how numerical semigroups are connected to commutative ring theory. In section 4 we will present an algorithm that creates the directed tree graphs and construct a few examples. To conclude the paper, we will present some new results that help describe the structure of these directed tree graphs.

## 2 Numerical Semigroups

We first need a solid grasp of what a numerical semigroup is, and more precisely, what it means to be an irreducible numerical semigroup.

**Definition 2.1.** A *numerical semigroup* is a subset  $S$  of the nonnegative integers  $\mathbb{N}$  that satisfies the following:

1.  $S$  contains 0.
2.  $S$  is closed under addition.
3.  $\mathbb{N} \setminus S$  is finite.

We say  $\{a_1, a_2, \dots, a_n\}$  is a *generating set* for  $S$  if  $S = \{k_1 a_1 + k_2 a_2 + \dots + k_n a_n \mid k_1, k_2, \dots, k_n \in \mathbb{N}\}$ , and we call each  $a_i$  a *generator* of  $S$ . If no proper subset is a generating set for  $S$ , we say  $\{a_1, a_2, \dots, a_n\}$  is the *minimal generating set* and we write  $S = \langle a_1, a_2, \dots, a_n \rangle$ ,  $0 < a_1 < a_2 < \dots < a_n$ .

**Example 2.1.** Consider the numerical semigroup generated by 5, 6 and 13. Then

$$\begin{aligned} S = \langle 5, 6, 13 \rangle &= \{5k_1 + 6k_2 + 13k_3 \mid k_1, k_2, k_3 \in \mathbb{N}\} \\ &= \{0, 5, 6, 10, 11, 12, 13, 15, 16, 17, 18, 19, 20, 21, 22, 23, \dots\} \\ &= \{0, 5, 6, 10, 11, 12, 13, 15, \rightarrow\} \end{aligned}$$

The arrow indicates that all integers greater than 15 are in  $S$ .

Associated with every numerical semigroup are two important integers which are important for our studies.

**Definition 2.2.** The *Frobenius number* of a numerical semigroup  $S$ , denoted  $F(S)$ , is the largest integer not in  $S$ . The *multiplicity* of  $S$ , denoted  $m(S)$ , is the smallest positive element of  $S$ .

**Example 2.2.** Consider once again the numerical semigroup generated by 5, 6, and 13. That is,

$$S = \langle 5, 6, 13 \rangle = \{0, 5, 6, 10, 11, 12, 13, 15, \rightarrow\}$$

Here  $F(S) = 14$  as it is the largest integer not in  $S$ , and  $m(S) = 5$ , as 5 is the smallest positive element in  $S$ .

Next we define the two special types of numerical semigroups that are the focus of our research.

**Definition 2.3.** A numerical semigroup  $S$  is said to be *symmetric* if and only if  $F(S)$  is odd and if  $x \in \mathbb{Z} \setminus S$ , then  $F(S) - x \in S$ .

**Example 2.3.** Consider,  $S = \langle 6, 7, 10, 11 \rangle = \{0, 6, 7, 10, 11, 12, 13, 14, 16, \rightarrow\}$ . Here  $F(S) = 15$ , which is odd, as required. In addition, for every integer  $x \notin S$  we have  $F(S) - x \in S$  as depicted in Figure 1 below. For example, if  $x = 8$ , we have  $15 - 8 = 7 \in S$ , and the same can be verified for all elements in  $\mathbb{Z} \setminus S$ . Thus  $S$  is symmetric.




Figure 1

**Definition 2.4.** A numerical semigroup  $S$  is said to be *pseudosymmetric* if and only if  $F(S)$  is even and if  $x \in \mathbb{Z} \setminus S$ , then either  $x = \frac{F(S)}{2}$  or  $F(S) - x \in S$ .

**Example 2.4.** Again consider  $S = \langle 5, 6, 13 \rangle = \{0, 5, 6, 10, 11, 12, 13, 15, \rightarrow\}$ . As previously discussed,  $F(S) = 14$  is even as required. As indicated in Figure 2, for every integer  $x \notin S$  we have either  $x = \frac{F(S)}{2}$  or  $F(S) - x \in S$ , and so  $S$  is pseudosymmetric.




Figure 2

The reason for our emphasis on symmetric and pseudosymmetric numerical semigroups will be made clear in the remaining sections of the paper.

## 2.1 Irreducible Numerical Semigroups

**Definition 2.5.** A numerical semigroup  $S$  is *irreducible* if it cannot be expressed as an intersection of two numerical semigroups that properly contain  $S$ . For a given Frobenius number  $F$ , the set of all irreducible numerical semigroups with Frobenius number  $F$  is denoted  $I(F)$ .

In [2], Branco and Rosales showed that a numerical semigroup is irreducible if and only if it is symmetric or pseudosymmetric. The following theorem [1] defines a certain numerical semigroup unique to every Frobenius number.

**Theorem 2.1.** *For every positive integer  $F$ , there exists a unique irreducible numerical semigroup  $C(F)$  whose generators are all larger than  $\frac{F}{2}$ . Moreover,*

$$\begin{aligned} C(F) &= \begin{cases} \{0, \frac{F+1}{2}, \rightarrow\} \setminus \{F\} & \text{if } F \text{ is odd,} \\ \{0, \frac{F}{2} + 1, \rightarrow\} \setminus \{F\} & \text{if } F \text{ is even.} \end{cases} \\ &= \begin{cases} \langle \frac{F+1}{2}, \frac{F+3}{2}, \dots, F-1 \rangle & \text{if } F \text{ is odd,} \\ \langle \frac{F}{2} + 1, \frac{F}{2} + 2, \dots, F-1, F+1 \rangle & \text{if } F \text{ is even.} \end{cases} \end{aligned}$$

**Example 2.5.** Using Theorem 2.1 we will now construct  $C(15)$ .

$$\begin{aligned} C(15) &= \{0, \frac{15+1}{2}, \rightarrow\} \setminus \{15\} \\ &= \{0, 8, 9, 10, 11, 12, 13, 14, 15, 16, \rightarrow\} \setminus \{15\} \\ &= \{0, 8, 9, 10, 11, 12, 13, 14, 16, \rightarrow\} \\ &= \langle 8, 9, 10, 11, 12, 13, 14 \rangle \end{aligned}$$

Note that when  $F$  is odd,  $C(F)$  is generated by an interval of consecutive integers of  $\mathbb{N}$ . This type of numerical semigroup, first investigated by García-Sánchez and Rosales in [3], led to a new characterization of symmetry for interval-generated numerical semigroups. The interested reader can consult [3] for more information.

## 3 Graph Theory Necessities

The remainder of this paper will focus on graphs of families of specific numerical semigroups that share a common Frobenius number. Therefore, a brief review of basic graph theory terminology is appropriate.

**Definition 3.1.** A *graph* is a collection of points called *vertices* and lines called *edges* that connect pairs of vertices. Two vertices are *adjacent* if an edge connects them.

**Definition 3.2.** A *tree graph* is a graph in which there is a unique sequence of alternating edges and vertices “connecting” each pair of vertices. A *directed tree graph* is a tree graph in which there is a direction associated with each edge.

**Example 3.1.** In Figure 3 we see an example of a directed tree graph. We will reference this tree in order to clearly define the terms that we use to describe tree graphs.




Figure 3

The designated vertex from which the tree appears to “hang” - i.e., the vertex ( $S_1$ ) at “the top” of the tree - is called the *root* of the tree. The *level* of a vertex is the length of the path from the vertex to the root. For instance, in Figure 3 the level of  $S_2$  is 1, and we say that  $S_2$  is in the first level. The *height* of a tree is the maximum level in the tree. Note that the height of the tree in Figure 3 is 3. Given any pair of adjacent vertices, the vertex in the greater level is called the *child* of the vertex in the smaller level. For example, in Figure 3,  $S_3$  is a child of  $S_1$ , and  $S_5$ ,  $S_6$ , and  $S_7$  are the children of  $S_2$ . A vertex with no children is called a *leaf*. A *branch* is the shortest “path” from the root to a leaf. As a result, the longest branch in a tree is equivalent to the height of the tree. Note that in Figure 3 the longest branch is the path from  $S_1$  to  $S_8$  passing through  $S_2$  and  $S_6$ .

## 4 Directed Tree Graphs

In an attempt to catalogue all symmetric and pseudosymmetric numerical semigroups with a fixed Frobenius number, Blanco and Rosales [1] developed an algorithm for systematically finding all such irreducible numerical semigroups.

**Theorem 4.1.** *Let  $F$  be a positive integer. Then the elements of  $I(F)$  comprise a directed tree graph, denoted  $G(I(F))$ , with root  $C(F)$ . If  $S$  is an element of  $I(F)$ , then the children of  $S$  are  $(S \setminus \{x_1\} \cup \{F - x_1\})$ ,  $(S \setminus \{x_2\} \cup \{F - x_2\})$ , ...,  $(S \setminus \{x_r\} \cup \{F - x_r\})$ , where  $\{x_1, \dots, x_r\}$  is the set of minimal generators of  $S$  such that for each  $x \in \{x_1, \dots, x_r\}$  the following conditions are satisfied:*

1.  $\frac{F}{2} < x < F$
2.  $2x - F \notin S$



$$3. 3x \neq 2F$$

$$4. 4x \neq 3F$$

$$5. F - x < m(s).$$

When a minimal generator,  $x$ , of  $S \in (G(I(F)))$  satisfies the five conditions, we say that  $x$  *spawns* a child of  $S$ , that child being  $(S \setminus \{x\} \cup \{F - x\})$ . A minimal generator of a given vertex that does not spawn a child is said to be *barren*.

**Example 4.1.** In order to build  $G(I(11))$ , the directed tree graph associated with Frobenius number 11, we begin by constructing the root,  $C(11)$ :

$$\begin{aligned} C(11) &= \left\{0, \frac{F(S) + 1}{2}, \rightarrow\right\} \setminus \{F\} = \{0, 6, 7, 8, 9, 10, 12, \rightarrow\} \\ &= \langle 6, 7, 8, 9, 10 \rangle. \end{aligned}$$

Using Theorem 4.1 and checking each minimal generator we find that the following generators spawn the given irreducible numerical semigroups in  $G(I(11))$ .

$$x_1 = 8 : (\langle 6, 7, 8, 9, 10 \rangle \setminus \{8\}) \cup \{3\} = \{0, 3, 6, 7, 9, 10, 12, \rightarrow\} = \langle 3, 7 \rangle$$

$$x_2 = 7 : (\langle 6, 7, 8, 9, 10 \rangle \setminus \{7\}) \cup \{4\} = \{0, 4, 6, 8, 9, 10, 12, \rightarrow\} = \langle 4, 6, 9 \rangle$$

$$x_3 = 6 : (\langle 6, 7, 8, 9, 10 \rangle \setminus \{6\}) \cup \{5\} = \{0, 5, 7, 8, 9, 10, 12, \rightarrow\} = \langle 5, 7, 8, 9 \rangle$$

These irreducible numerical semigroups are the children of the root  $C(11)$  in  $G(I(11))$ . By applying the same algorithm we find that, in  $G(I(11))$ ,  $\langle 3, 7 \rangle$  is barren, whereas  $\langle 4, 6, 9 \rangle$  has one child,  $\langle 2, 13 \rangle$ , and  $\langle 5, 7, 8, 9 \rangle$  has one child,  $\langle 4, 5 \rangle$  spawned by 9 and 7, respectively. The entire tree  $G(I(11))$  is shown in Figure 4 below.




Figure 4

The following images of specific directed tree graphs were generated by a program written in Mathematica and compiled in QTikZ. For simplicity, the tree vertices are labels of the form  $S_1, S_2, S_3$ , etc. (corresponding to the order in which they were found by the program) and the minimal generator from which each child spawned is not shown.

**Example 4.2.** Figure 5 below is an image of  $G(I(16))$ . It should be noted that this tree is isomorphic to  $G(I(18))$ , and in all likelihood they are the only such pair of isomorphic trees in the entire family  $G(I(F))$  for  $F > 11$ .




Figure 5

**Example 4.3.** Figure 6 is an image of  $G(I(30))$ .




Figure 6

The tree in Figure 6 shows many of the obvious characteristics of the trees for  $F > 12$ , as the left side contains longer branches and more children. Moreover, the “width” and the number of vertices vary dramatically from tree to tree. Further exploration of the structure of the trees can be found in the next section.

**Example 4.4.** Figure 7 is an image of  $G(I(27))$ .




Figure 7

## 5 New Results

In our investigation we have been especially interested in finding explicit formulas for the number of vertices, the width and the height of any given tree. This has in general been difficult, so we have resorted to examining less complicated properties of the trees which may lead to finding of formulas. For instance, we have investigated which minimal generators will spawn children in  $G(I(F))$  and how often, and we have examined the longest branches in the trees. To this end, the trees have been split into two categories, the odd Frobenius number trees and the even Frobenius number trees. The following properties of  $G(I(F))$  have been found:

**Proposition 5.1.** *Let  $F = 2k + 1$ . Then:*

1. *If  $3|F$ , then the minimal generator  $k + \frac{k+2}{3}$  does not spawn children in  $G(I(F))$ .*
2. *If  $k$  is even, then the minimal generators of  $C(F)$  greater than or equal to  $k + \frac{k}{2} + 1$  will not spawn children of  $C(F)$ .*
3. *If  $k$  is odd, then the minimal generators of  $C(F)$  greater than or equal to  $k + \frac{k+1}{2} + 1$  will not spawn children of  $C(F)$ .*

*Proof.* To prove part (1), note that  $3(k + \frac{k+2}{3}) = 4k + 2 = 2F$ . This violates condition 3 in Theorem 4.1, and thus the minimal generator  $k + \frac{k+2}{3}$  is barren.

For (2), note that  $2(k + \frac{k}{2} + 1) - (2k + 1) = k + 1$ . Since  $k + 1$  is a minimal generator of  $C(F)$ ,  $2(k + \frac{k}{2} + 1) - F \in C(F)$ , which violates condition 2 in Theorem 4.1. Also, if  $n > \frac{k}{2} + 1$ , then  $2(k + n) - F > k + 1$ , which means that  $2n - F \in C(F)$ . Therefore,  $k + n$  will not spawn children of  $C(F)$ .

Finally, for part (3) note that  $2(k + \frac{k+1}{2} + 1) - (2k + 1) = k + 2$ , and since  $k + 2$  is a minimal generator of  $C(F)$ ,  $k + \frac{k+1}{2} + 1 - F \in C(F)$ , which violates condition 2. Likewise for  $n \geq \frac{k+1}{2} + 2$ , note that  $2(k + n) - F > k + 2$ , so  $2n - F \in C(F)$ . Thus,  $k + n$  will not spawn children of  $C(F)$ .  $\square$

The following is another small, yet useful result we have found for studying the odd Frobenius number trees.

**Proposition 5.2.** *Let  $F = 2k + 1, k > 1$ . The smallest odd minimal generator of  $C(F)$  will always spawn a child of  $C(F)$ .*

*Proof.* For this proof we must show that the smallest odd minimal generator of  $C(F)$  satisfies all conditions of Theorem 4.1. For the first condition, suppose  $k$  is odd. The smallest odd minimal generator of  $C(F)$  is then  $k + 2$ . Note that  $\frac{F}{2} = k + \frac{1}{2}$ , hence  $\frac{F}{2} < k + 2 < F$ . For the second condition, note that  $2(k + 2) - F = 3$ , and since  $F > 3, k > 2$ . Thus  $k + 1 > 3$ , so  $2(k + 1) - F \notin S$ . For the third condition, suppose  $3(k + 2) = 2F$ . Then  $3k + 6 = 4k + 2$  and  $k = 4$ , which contradicts the assumption that  $k$  is odd. Therefore,  $3(k + 2) \neq 2F$ . To show that the fourth condition is satisfied, suppose that  $4(k + 2) = 3F$ . Then  $4k + 8 = 6k + 3$ ,

which implies that  $2k = 5$ , which is false. Therefore,  $4(k+2) \neq 3F$ . For the fifth condition, note that  $F - (k+2) = k-1$  and  $m(C(F)) = k+1$ , thus  $F - (k+2) < m(C(F))$ . Therefore,  $k+2$  will spawn a child of  $C(F)$ .

Next, suppose  $k$  is even. Then the smallest odd minimal generator of  $C(F)$  is  $k+1$ . For the first condition, note that  $\frac{F}{2} < k+1 < F$ . For the second condition, note  $2(k+1) - F = 1 < k+1$ , hence  $2(k+1) - F \notin C(F)$ . For the third condition, note that  $3(k+1) = 2F$  implies that  $k = 1$ , which is false because  $k > 1$ . Therefore,  $3(k+1) \neq 2F$ . For the fourth condition,  $4(k+1) = 3F$  implies that  $2k = 1$ , which is false. Therefore,  $4(k+1) \neq 3F$ . Finally, for the fifth condition, note that  $F - (k+1) = k$  and  $m(C(F)) = k+1$ , so  $F - (k+1) < m(C(F))$ . Therefore,  $k+1$  will spawn a child of  $C(F)$ .  $\square$

We have also derived analogous results for even Frobenius number trees:

**Proposition 5.3.** *Let  $F = 2k$ . Then:*

1. *If  $4|F$ , then the minimal generator  $k + \frac{F}{4}$  does not spawn children in  $G(I(F))$ .*
2. *If  $6|F$ , then the minimal generator  $k + \frac{F}{6}$  does not spawn children in  $G(I(F))$ .*
3. *If  $k$  is even,  $4 \nmid F$  and  $6 \nmid F$ , then the minimal generators of  $C(F)$ ,  $k+n$ , where  $1 \leq n \leq \frac{k}{2}$  will each spawn a child of  $C(F)$ .*

*Proof.* For the proof of part (1), note that  $4(k + \frac{F}{4}) = 4(k + \frac{2k}{4}) = 6k$ , and  $3F = 6k$ . Thus,  $4(k + \frac{F}{4}) = 3F$ , which violates condition 4 in Theorem 4.1. Therefore,  $k + \frac{F}{4}$  is barren in  $G(I(F))$ . For part (2), note that  $3(k + \frac{F}{6}) = 3(k + \frac{2k}{6}) = 4k$ , and  $2F = 4k$ . Thus,  $3(k + \frac{F}{6}) = 2F$ , which violates condition 3.

Finally, for part (3), note that to prove that a minimal generator spawns a child, it must be shown that the 5 conditions in Theorem 4.1 are satisfied. Let  $1 \leq n \leq \frac{k}{2}$ . For condition one, note that  $\frac{F}{2} = k$ , so  $\frac{F}{2} < k+n < F$ . For condition 2, note that  $2(k+n) - F = 2n < m(C(F))$ . For condition 3, note that if  $3(k+n) = 2F$ , then  $3k+3n = 4k$ , and  $3n = k$ , which contradicts the assumption  $6 \nmid F$ . Also, for condition 4, if  $4(k+n) = 3F$ , then  $4k+4n = 6k$  and  $4n = 2k$ , which contradicts the assumption  $4 \nmid F$ . Finally, for condition 5, note that  $F - k - n$ , so  $F - (k+n) \leq m(C(F))$  since  $n \geq 1$ . Therefore  $k+n$ ,  $1 \leq n \leq \frac{k}{2}$  will each spawn a child of  $C(F)$ .  $\square$

We have devoted a great deal of effort toward determining the heights of these trees. We have been especially interested in the longest branches of trees with odd Frobenius number greater than 11. For this study, the following lemma has been found to be useful.

**Lemma 5.1.** *Let  $S$  be an irreducible numerical semigroup with Frobenius number  $F = 2k+1$  such that  $S$  is spawned by  $k+n$ , where  $k+n$  is an odd minimal generator of  $C(F)$  for some positive integer  $n$ . Stated differently:  $S = S^* \setminus \{k+n\} \cup \{F - (k+n)\}$ , where  $S$  is a child of  $S^* \in G(I(F))$ . Then:*

1. *If  $k$  is even and  $w$  is the smallest odd such that  $k+n < k+w < F$  and  $k+w$  is the smallest odd minimal generator of  $S$ , then  $k+w$  will spawn a child of  $S$ .*

2. If  $k$  is odd and  $h$  is the smallest even such that  $k + n < k + h < F$  and  $k + h$  is the smallest odd minimal generator of  $S$ , then  $k + h$  will spawn a child of  $S$ .

*Proof.* First, suppose  $k$  is even. We must show that the minimal generators  $k + n$  satisfy conditions 1-5. For the first condition, note that  $\frac{F}{2} = k + \frac{1}{2} < k + w$ , so  $\frac{F}{2} < k + w < F$ . For the second condition, note that  $2(k + w) - F = 2w - 1$  and since  $w < k + 1$ , then  $2w < k + w + 1$  and  $2w - 1 < k + w$ , and since  $k + w$  is the smallest odd in  $S$ , then  $2w - 1 \notin S$ . For the third condition, note that if  $3(k + w) = 2F$ , then  $k = 3w - 2$ . But  $k$  is even and  $3w - 2$  is odd. Thus,  $3(k + w) \neq 2F$ . To show that the fourth condition is satisfied, note that  $4(k + w) = 3F$  implies that  $4w = 2k + 3$ . But  $2k + 3$  is odd and  $4w$  is even. Hence,  $4(k + w) \neq 3F$ . Finally, for the fifth condition, note that  $F - (k + w) = k + 1 - w$ , and since  $w > n$ , then  $k + 1 - w < k + 1 - n$ . Thus,  $F - (k + w) < m(S)$ . Therefore,  $k + w$  will spawn a child of  $S$ .

For the second part, suppose that  $k$  is odd. For the first condition, note that  $\frac{F}{2} = k + \frac{1}{2}$ , so  $\frac{F}{2} < k + h < F$ . For the second condition, note that  $2(k + h) - F = 2h - 1$  and recall that  $h < k + 1$ , so  $2h - 1 < k + h$ . Since  $k + h$  is the smallest odd in  $S$ , then  $2h - 1 \notin S$ . For the third condition, note that  $3(k + h) = 2F$  implies that  $k = 3h - 2$ . But  $k$  is odd and  $3h - 2$  is even. Thus,  $3(k + h) \neq 2F$ . For the fourth condition, note that if  $4(k + h) = 3F$ , then  $4h - 3 = 2k$ , but  $4h - 3$  is odd, a contradiction. Thus,  $4(k + h) \neq 3F$ . Finally, for the fifth condition, note that  $F - (k + h) = k + 1 - h$ . Since  $m < h$ , then  $k + 1 - h < k + 1 - m$ . Thus,  $k + 1 - h < m(S)$ . Therefore,  $k + h$  will spawn a child of  $S$ .  $\square$

To illustrate what Lemma 5.1 says, we present the following example:

**Example 5.1.** Let  $F = 2k + 1 = 11$ . Then  $k = 5$ . Consider the child of  $C(11)$   $S$  spawned by  $k + 2 = 7$ :

$$\begin{aligned} S &= C(11) \setminus \{7\} \cup \{11 - 7\} \\ &= \langle 4, 6, 9 \rangle. \end{aligned}$$

Note that  $k$  is odd, and 4 is the smallest even integer such that  $k + 2 < k + 4 < 11$  and  $k + 4 = 9$  is the smallest odd minimal generator of  $S$ . Thus, by Lemma 5.1, the minimal generator 9 will spawn a child of  $S$ .

The study of the odd Frobenius number trees led to the following:

**Theorem 5.1.** Let  $F = 2k + 1, k > 5$ . Note that  $C(F) = \langle k + 1, k + 2, k + 3, \dots, 2k \rangle$ . Then:

1. If  $k$  is even,  $G(I(F))$  contains a branch whose vertices are spawned by  $k + 1, k + 3, \dots, k + k - 1$  in this order.
2. If  $k$  is odd,  $G(I(F))$  contains a branch whose vertices are spawned by  $k + 2, k + 4, \dots, k + k - 1$  in this order.

*Proof.* Let  $k$  be even. Note that  $C(F) = \langle k+1, k+2, \dots, k+k \rangle$ . By Proposition 5.2,  $k+1$  will spawn a child of  $C(F)$ . Suppose  $k+n$  is the smallest odd minimal generator of a numerical semigroup  $S \in G(I(F))$ , where  $S$  is spawned by  $k+n-2$  and  $1 \leq k+n-2 \leq 2k-1$ . By Lemma 5.1,  $k+n$  will spawn a child,  $S'$ , of  $S$ .

Assuming the above, we must show that  $k+n+2$  will spawn a child of  $S'$ . Note that  $F - (k+n) = 2k+1 - (k+n)$  is even and all minimal generators of  $S'$  less than  $k+n$  are even. Thus  $k+n+2$  is the smallest odd minimal generator of  $S'$ . By Lemma 5.1,  $k+n+2$  will spawn a child of  $S'$ , as long as  $n+2 < 2k+3$ , as desired. A similar argument is used to prove part 2 and the result follows.  $\square$

**Example 5.2.** Figure 8 is an image of  $G(I(19))$ . Note that the longest branch is the one described in Theorem 5.1 .




Figure 8

We have generated images for all odd Frobenius number trees between  $F = 7$  and  $F = 67$ , and in all of these trees, where  $F > 11$ , the longest branch is the one described in Theorem 5.1 above. This leads us to the following conjecture.

**Conjecture 5.1.** *If  $F = 2k + 1$ , then the branch described in Theorem 5.1 is the unique longest branch in  $G(I(F))$ .*

## 6 Open Questions

We have yet to find a formula for the number of vertices in a given tree. This is an especially mysterious and difficult problem to explore since the number of vertices in the trees is not strictly increasing as the Frobenius numbers increase, as can be seen in Figure 9 below which shows the number of levels and vertices in  $G(I(F))$ .

$F(S)$	Levels	Vertices	$F(S)$	Levels	Vertices	$F(S)$	Levels	Vertices
14	2	6	27	6	45	40	6	196
15	3	7	28	4	37	41	10	420
16	2	7	29	7	83	42	6	203
17	4	15	30	4	36	43	10	546
18	2	7	31	7	109	44	7	342
19	4	20	32	5	70	45	11	498
20	3	11	33	8	101	46	7	527
21	5	18	34	5	106	47	11	926
22	3	20	35	8	174	48	7	411
23	5	36	36	5	77	49	12	1182
24	3	14	37	9	246	50	8	844
25	6	44	38	6	182	51	12	1121
26	4	35	39	9	227	52	8	981

Figure 9

Note that the Frobenius numbers which appear to be responsible for the non-increasing behavior of the numbers of vertices appear to always be divisible by 4 or 6 in the even case, and divisible by 3 in the odd case, which correspond to Propositions 5.1 and 5.3, respectively. Interesting as they are, these observations have not led to a conjecture for the number of vertices in a tree. As shown in the table, the number of vertices grows quite rapidly, making it very difficult to generate images of the trees with larger Frobenius numbers. The largest tree for which we have an image of is  $G(I(67))$ , which has 11972 vertices.

Determining the height of the trees is slightly less mysterious, and has led to the following conjecture.

**Conjecture 6.1.** *The height of  $G(I(F))$  is  $\lfloor \frac{k}{2} \rfloor$  for  $F = 2k + 1$  (corresponding to a unique branch of this length) and  $\lfloor \frac{k-1}{3} \rfloor$  for  $F = 2k$  (corresponding to non-unique branches of this length) for  $k > 6$ .*

Note that the conjectured height of the odd trees corresponds to the length of the branch described in Theorem 5.1. In our future work we hope to prove this conjecture. We also hope to find a formula for, or relatively sharp bounds on, the number of vertices in a given tree.

## References

- [1] Blanco, V., Rosales, J. C. *The Tree Of Irreducible Numerical Semigroups With Fixed Frobenius Number*. <http://arxiv.org/abs/1105.2147> (2011).

- [2] Branco, M. B., Rosales, J. C. *Irreducible Numerical Semigroups*. Pacific Journal of Mathematics. Vol. 209, No.1 (2003).
- [3] Garcíá-Sánchez, P. A., Rosales, J. C. *Numerical Semigroups Generated By Intervals*. Pacific Journal of Mathematics. Vol. 191, No. 1 (1999).
- [4] Rushall, J. *A Primer on Numerical Semigroups* (unpublished, 2010).