

Rose-Hulman Undergraduate Mathematics Journal

Volume 20
Issue 1

Article 5

Asymptotically Optimal Bounds for $(t,2)$ Broadcast Domination on Finite Grids

Timothy W. Randolph
Williams College, twr2@williams.edu

Follow this and additional works at: <https://scholar.rose-hulman.edu/rhumj>



Part of the [Discrete Mathematics and Combinatorics Commons](#)

Recommended Citation

Randolph, Timothy W. (2019) "Asymptotically Optimal Bounds for $(t,2)$ Broadcast Domination on Finite Grids," *Rose-Hulman Undergraduate Mathematics Journal*: Vol. 20 : Iss. 1 , Article 5.
Available at: <https://scholar.rose-hulman.edu/rhumj/vol20/iss1/5>

Asymptotically Optimal Bounds for $(t,2)$ Broadcast Domination on Finite Grids

Cover Page Footnote

Special thanks to Pamela E. Harris for her feedback on earlier versions of this draft.

Asymptotically Optimal Bounds for $(t, 2)$ broadcast Domination on Finite Grids

By Timothy W. Randolph

Abstract. Let $G = (V, E)$ be a graph and t, r be positive integers. The *signal* that a tower vertex T of signal strength t supplies to a vertex v is defined as $sig(T, v) = \max(t - dist(T, v), 0)$, where $dist(T, v)$ denotes the distance between the vertices v and T . In 2015 Blessing, Insko, Johnson, and Mauretour defined a (t, r) *broadcast dominating set*, or simply a (t, r) *broadcast*, on G as a set $\mathbb{T} \subseteq V$ such that the sum of all signals received at each vertex $v \in V$ from the set of towers \mathbb{T} is at least r . The (t, r) broadcast domination number of a finite graph G , denoted $\gamma_{t,r}(G)$, is the minimum cardinality over all (t, r) broadcasts for G .

1 Introduction

Let u and v be two vertices of the connected graph $G = (V, E)$. By $dist(u, v)$ we denote the distance between u and v , which is defined as the length of the shortest path between u and v or 0 if $u = v$. A dominating set for G is a subset $S \subseteq V$ such that for every $v \in V$, there exists a vertex $s \in S$ with $dist(v, s) \leq 1$. The domination number $\gamma(G)$ of a graph G is the cardinality of the smallest dominating set on G . In 1992, Chang [2] established the following bound on the domination number of the $m \times n$ grid graph $G_{m,n}$ with $m, n > 8$

$$\gamma(G_{m,n}) \leq \left\lfloor \frac{(n+2)(m+2)}{5} \right\rfloor - 4. \quad (1)$$

Goncalves, Pinlou, Rao and Thomasse [5] proved in 2011 that equality holds in Equation 1 when $n \geq m \geq 16$. For a historical perspective on the development of domination theory and related problems, the reader is referred to the work of Haynes, Hedetniemi, and Slater [6].

As the field of domination theory grows, questions arise concerning generalizations of the domination number. One generalization considers the k -distance dominating set for a graph $G = (V, E)$, defined as a subset $S \subseteq V$ such that for every $v \in V$, there exists a vertex $s \in S$ such that $dist(v, s) \leq k$. We denote the k -distance domination number as

Mathematics Subject Classification. 05C69

Keywords. Domination, Broadcasts, Grid graphs

$\gamma_k(G)$. In 2014, Grez and Farina [4] succeeded in bounding the k -distance domination number of the $m \times n$ grid graph as follows

$$\gamma_k(G_{m,n}) \leq \left\lfloor \frac{(m+2k)(n+2k)}{(2k^2+2k+1)} \right\rfloor - 4. \quad (2)$$

In 2014, Blessing, Insko, Johnson and Mauretour [1] defined (t, r) broadcast domination for positive integers t, r . In this setting, given a graph $G = (V, E)$ and a positive integer t , we say that a *tower vertex* T supplies a signal equal to $\text{sig}(T, v) = \max\{t - \text{dist}(T, v), 0\}$ to each vertex $v \in V$. A (t, r) broadcast is a set $S \subseteq V$ of tower vertices such that the sum of all signals supplied to each vertex is at least r . The (t, r) broadcast domination number of a graph G , denoted $\gamma_{t,r}(G)$, is defined as the minimal cardinality among all (t, r) broadcasts on G . We note that (t, r) broadcast domination is a natural generalization of domination and k -domination, as $\gamma(G) = \gamma_{2,1}(G)$ and $\gamma_k(G) = \gamma_{k+1,1}(G)$.

In their work, Blessing et al. studied the (t, r) broadcast domination number of grid graphs for the (t, r) pairs in the set $\{(2, 2), (3, 1), (3, 2), (3, 3)\}$. In particular, they established the following bounds on $(t, 2)$ broadcast domination numbers for grid graphs

$$\gamma_{2,2}(G_{m,n}) \leq \left\lfloor \frac{(m+2)(n+2)}{3} \right\rfloor - 5. \quad (3)$$

$$\gamma_{3,2}(G_{m,n}) \leq \left\lfloor \frac{(m+2)(n+2)}{8} \right\rfloor - 1. \quad (4)$$

In this paper, we establish an asymptotically optimal upper bound for $\gamma_{t,2}(G_{m,n})$ when $t > 2$.

If $G_{m,n}$ is the grid graph with dimensions $m \times n$, and $t > 2$, then

$$\gamma_{t,2}(G_{m,n}) \leq \left\lfloor \frac{(m+2(t-2))(n+2(t-2))}{2(t-1)^2} \right\rfloor.$$

In addition, we prove the following lower bound on $\gamma_{t,2}(G_{m,n})$, which is a corollary of a result of Drews, Harris, and Randolph [3].

If $G_{m,n}$ is the grid graph with dimensions $m \times n$, and $t > 2$, then

$$\gamma_{t,2}(G_{m,n}) \geq \frac{mn}{2(t-1)^2}.$$

The upper and lower bounds we establish are separated by a gap linear in m and n , and thus converge as m and n increase. In addition, Theorem 1 confirms the conjecture of Blessing et al. that Equation 4 is asymptotically optimal.

2 Bounding $\gamma_{t,2}(G_{m,n})$

In 2017, Drews, Harris, and Randolph [3] established the optimal $(t, 2)$ broadcast on G_∞ , the infinite graph

$$(\{v_{(i,j)} \mid i, j \in \mathbb{Z}\}, \{(v_{(i,j)}, v_{(i+1,j)}), (v_{(i,j)}, v_{(i,j+1)}) \mid i, j \in \mathbb{Z}\}).$$

Because no finite set is a (t, r) broadcast on G_∞ , the authors define an optimal (t, r) broadcast as a (t, r) broadcast with minimal broadcast density. The *broadcast density* of an infinite (t, r) broadcast S is defined as

$$\lim_{n \rightarrow \infty} \frac{|S \cap V_{n \times n}|}{|V_{n \times n}|},$$

where $V_{n \times n}$ is an $n \times n$ grid of vertices anchored by a fixed vertex in G_∞ . Intuitively, the broadcast density captures the proportion of the vertices of G_∞ contained in an infinite (t, r) broadcast.

In this section, we construct a $(t, 2)$ broadcast on the grid graph $G_{m,n}$ by transforming a subset of an optimal $(t, 2)$ broadcast on G_∞ . We then demonstrate that our upper bound on $\gamma_{t,2}(G_{m,n})$ and the upper bound of Blessing et al. on $\gamma_{3,2}(G_{m,n})$ converge to the optimal value as m and n increase, a result that follows from the optimality of the original $(t, 2)$ broadcast on G_∞ . To begin our analysis we recall the following theorem of Drews et al. [3].

Theorem 2.1. *If $t > 2$, the optimal broadcast density of a $(t, 2)$ broadcast on G_∞ is*

$$\gamma_{t,2}(G_\infty) = \frac{1}{2(t-1)^2}.$$

The *broadcast outline* of a tower vertex T is defined as the diamond shape formed by connecting the vertices of the set $\{v : \text{dist}(T, v) = t - 1\}$ when a grid graph $G_{m,n}$ is embedded in $\mathbb{Z} \times \mathbb{Z}$. Drews et al. proved that any set of vertices whose broadcast outlines form a tiling of $\mathbb{Z} \times \mathbb{Z}$ is an optimal $(t, 2)$ broadcast.

Figure 1 illustrates two tilings created by the broadcast outlines of optimal $(3, 2)$ broadcasts embedded in $\mathbb{Z} \times \mathbb{Z}$. Figure 1b illustrates an optimal $(t, 2)$ broadcast in which the broadcast outlines are aligned to form a diagonal lattice, hereafter referred to as a *rectilinear broadcast*.

To establish our bound on $\gamma_{t,2}(G_{m,n})$, we describe how a subset of the rectilinear $(t, 2)$ broadcast on G_∞ can be transformed into a $(t, 2)$ broadcast on $G_{m,n}$. Counting the number of elements in this subset gives an upper bound on $\gamma_{t,2}(G_{m,n})$.

We first prove the special case when $G_{m,n}$ is a path, in which case $m = 1$ or $n = 1$.

Lemma 2.2. *Let G be a path of length m , and t an integer greater than 2. Then*

$$\gamma_{t,2}(G) \leq \left\lfloor \frac{(m + 2(t - 2))(1 + 2(t - 2))}{2(t - 1)^2} \right\rfloor.$$




Figure 1: Optimal $(3, 2)$ broadcasts on the infinite grid.

Proof. We first observe that paths of length up to $2(t - 1) - 1$ can be dominated by a single vertex, as illustrated in Figure 2a. Likewise, two vertices at a distance of $2(t - 1)$ suffice to dominate a path of length up to $4(t - 1) - 1$. In general, k vertices spaced at intervals of $2(t - 1)$ constitute a $(t, 2)$ broadcast for paths of length up to $2k(t - 1) - 1$. Figure 2b illustrates such a broadcast in the $k = 3, t = 4$ case.




Figure 2: $(4, 2)$ broadcasts on $G_{m,1}$.

It follows that

$$\gamma_{t,2}(G_{m,1}) \leq \left\lfloor \frac{m + 2(t - 1)}{2(t - 1)} \right\rfloor = \left\lfloor \frac{mt - m + 2t^2 - 4t + 2}{2(t - 1)^2} \right\rfloor. \tag{5}$$

To establish this result, we must show that

$$\gamma_{t,2}(G_{m,1}) \leq \left\lfloor \frac{(m + 2(t - 2))(1 + 2(t - 2))}{2(t - 1)^2} \right\rfloor = \left\lfloor \frac{2mt - 3m + 4t^2 - 14t + 12}{2(t - 1)^2} \right\rfloor. \tag{6}$$

Comparing the numerators of Equations 5 and 6, we find that when $m > 1$ and $t > 2$, and when $m \geq 1$ and $t > 3$,

$$mt - m + 2t^2 - 4t + 2 \leq 2mt - 3m + 4t^2 - 14t + 12.$$

Finally, in the $m = 1, t = 3$ case, we have that

$$\left\lfloor \frac{mt - m + 2t^2 - 4t + 2}{2(t-1)^2} \right\rfloor = \left\lfloor \frac{2mt - 3m + 4t^2 - 14t + 12}{2(t-1)^2} \right\rfloor = 1. \tag{7}$$

Thus, the result holds for all $m \geq 1, t > 2$. □

Our next result establishes that when $m, n > 1$, $G_{m,n}$ can be dominated by a subset of a rectilinear $(t, 2)$ broadcast covering an area only slightly larger than $m \times n$.

Let $G_{m,n} = (V_G, E_G)$ be the grid graph with dimensions $m \times n$ induced on G_∞ by the vertices $\{v_{(i,j)} \mid 0 \leq i < m, 0 \leq j < n\}$. Similarly, let $H_{m,n} = (V_H, E_H)$ be the grid graph with dimensions $(m + 2(t - 2)) \times (n + 2(t - 2))$ induced on G_∞ by the vertices $\{v_{(i,j)} \mid -(t - 2) \leq i < m + (t - 2), -(t - 2) \leq j < n + (t - 2)\}$. Thus $G_{m,n}$ is centered inside $H_{m,n}$. Let \mathbb{T} be a rectilinear $(t, 2)$ broadcast on G_∞ . For an example see Figure 3, which illustrates $G_{12,6}$, $H_{12,6}$, and \mathbb{T} in the $t = 3$ case.




Figure 3: $G_{12,6}$, $H_{12,6}$, and rectilinear $(3, 2)$ broadcast \mathbb{T} .

We now establish that the subset of \mathbb{T} within $H_{m,n}$ is sufficient to dominate $G_{m,n}$.

Lemma 2.3. *Let m, n , and t be integers such that $m, n > 1$ and $t > 2$. Let \mathbb{T} be a rectilinear broadcast on G_∞ , and let $G_{m,n}$ and $H_{m,n}$ be grid graphs embedded in $\mathbb{Z} \times \mathbb{Z}$ as described previously. Then the set of tower vertices $V_H \cap \mathbb{T}$ supplies at least 2 signal to each vertex $v_{(i,j)} \in V_G$.*

Proof. Let $v_{(i,j)}$ be a vertex in V_G . We consider each possible location of $v_{(i,j)}$ with respect to the broadcast outlines of tower vertices in \mathbb{T} .

Suppose first that $v_{(i,j)}$ is contained inside the broadcast outline of a tower vertex $T \in \mathbb{T}$, in which case $\text{dist}(T, v_{(i,j)}) \leq t - 2$ and $\text{sig}(T, v) \geq 2$. As every vertex u with $\text{dist}(u, v_{(i,j)}) \leq t - 2$ is contained in V_H , $T \in V_H$ and $V_H \cap \mathbb{T}$ supplies at least 2 signal to $v_{(i,j)}$.

Suppose that $v_{(i,j)}$ is located on the broadcast outlines of exactly two towers $T_{(x_0,y_0)}$ and $T_{(x_1,y_1)}$ in \mathbb{T} . As \mathbb{T} is rectilinear, this condition implies that $v_{(i,j)}$ is located along a straight edge of each broadcast outline, not at a corner. As $|i - x_0| + |j - y_0| = |i - x_1| + |j - y_1| = t - 1$, and $v_{(i,j)}$ is not located at the corner of a broadcast outline, we have that

$$|i - x_0|, |i - x_1|, |j - y_0|, |j - y_1| < t - 1.$$

Thus, we have that $T_{(x_0,y_0)}, T_{(x_1,y_1)} \in V_H$ and $V_H \cap \mathbb{T}$ supplies at least 2 signal to $v_{(i,j)}$.

Finally, suppose that $v_{(i,j)}$ is located at the corner of four broadcast outlines. As \mathbb{T} is rectilinear, the towers corresponding to these broadcast outlines are located at the points $(i - (t - 1), j)$, $(i + (t - 1), j)$, $(i, j - (t - 1))$, and $(i, j + (t - 1))$. Because $m, n > 1$ by assumption, we have that at least two of these towers must be contained within H_V , which ensures that $V_H \cap \mathbb{T}$ supplies at least 2 signal to $v_{(i,j)}$. \square

With Lemmas 2.2 and 2.3 in hand, we are now prepared to prove our main result.

If $G_{m,n}$ is the grid graph with dimensions $m \times n$, and $t > 2$, then

$$\Upsilon_{t,2}(G_{m,n}) \leq \left\lfloor \frac{(m + 2(t - 2))(n + 2(t - 2))}{2(t - 1)^2} \right\rfloor.$$

Proof. Lemma 2.2 establishes Theorem 1 in the case where $m = 1$ or $n = 1$. We proceed with the assumption that $m, n > 1$.

Let \mathbb{T} be a rectilinear broadcast on G_∞ , and let $G_{m,n}$ and $H_{m,n}$ be grid graphs embedded in $\mathbb{Z} \times \mathbb{Z}$ as described previously. By Lemma 2.3, the vertex set $V_H \cap \mathbb{T}$ supplies at least 2 signal to each vertex in V_G .

We transform $V_H \cap \mathbb{T}$ into a $(t, 2)$ broadcast on $G_{m,n}$ by replacing each vertex in $(V_H \setminus V_G) \cap \mathbb{T}$ with a corresponding vertex in V_G . Let B denote the set that we will transform into our $(t, 2)$ broadcast. To begin, set B equal to $(V_H \setminus V_G) \cap \mathbb{T}$. For each vertex $v \in (V_H \setminus V_G) \cap \mathbb{T}$, there exists a unique vertex $v' \in V_G$ that minimizes $\text{dist}(v, v')$. For each vertex $v \in (V_H \setminus V_G) \cap \mathbb{T}$, remove v from B and replace it with v' . We claim that this operation leaves the cardinality of B unchanged and preserves the property that the vertices in B supply at least 2 signal to each vertex in V_G . Figure 4 illustrates the replacement operation for $G_{12,6}$ in the $t = 4$ case.

To prove our claim, we establish that the replacement operation maps each vertex $v \in (V_H \setminus V_G) \cap \mathbb{T}$ to a vertex $v' \notin V_G \cap \mathbb{T}$ and that for any two vertices u and v in $(V_H \setminus V_G) \cap \mathbb{T}$, the replacement operation maps u and v to distinct vertices.




Figure 4: Vertices in $(V_H \setminus V_G) \cap \mathbb{T}$ are replaced with counterpart vertices in V_G to create a $(4, 2)$ broadcast on $G_{12,6}$.

First, let v be a vertex in $(V_H \setminus V_G) \cap \mathbb{T}$, and v' be the vertex in V_G that minimizes $dist(v, v')$. Because \mathbb{T} is a rectilinear broadcast, for any pair of vertices $s, t \in \mathbb{T}$, we have $dist(s, t) \geq 2(t - 1)$. However, $dist(v, v')$ is at most $2(t - 2)$, which occurs when v is located at a corner of $H_{m,n}$. Because $dist(v, v') < 2(t - 1)$ and $v \in \mathbb{T}$, $v' \notin \mathbb{T}$ and thus $v' \notin V_G \cap \mathbb{T}$.

Second, let u and v be two vertices in $(V_H \setminus V_G)$ such that the same vertex minimizes $dist(u, v')$ and $dist(v, v')$ over all vertices $v' \in V_G$. In this case, $dist(u, v)$ is at most $2(t - 2)$, which occurs when v' is located at a corner of $G_{m,n}$ and u and v are separated from v' by horizontal and vertical paths of length $(t - 2)$. Because $dist(u, v) < 2(t - 1)$, it is not possible that both u and v are in \mathbb{T} .

Thus, each time we replace a vertex $v \in (V_H \setminus V_G) \cap \mathbb{T}$ with the vertex $v' \in V_G$ that minimizes $dist(v, v')$, we are guaranteed that v' is not already in B . Furthermore, for every vertex $v \in (V_H \setminus V_G) \cap \mathbb{T}$, the vertex $v' \in V_G$ that minimizes $dist(v, v')$ satisfies $dist(v, w) > dist(v', w)$ for every vertex $w \in V_G$. Thus, B supplies at least as much signal to each vertex $v \in V_G$ as $V_H \cap \mathbb{T}$. B is thus a $(t, 2)$ dominating set for $G_{m,n}$.

We employ the probabilistic method to minimize $|B| = |V_H \cap \mathbb{T}|$ over all possible grid subgraphs H with dimensions $(m + 2(t - 2)) \times (n + 2(t - 2))$. Pick a coordinate (x, y) at random and determine a grid subgraph $H(x, y)$ by setting $v_{(x,y)}$ as its lower left corner. Because \mathbb{T} is an optimal $(t, 2)$ broadcast on the infinite grid, its broadcast density is

$\frac{1}{2(t-1)^2}$ by Theorem 2.1. The expected value of $|V_{H(x,y)} \cap \mathbb{T}|$ is thus $\frac{(m+2(t-2))(n+2(t-2))}{2(t-1)^2}$, which implies the existence of a point (x, y) for which

$$|V_{H(x,y)} \cap \mathbb{T}| = \left\lfloor \frac{(m+2(t-2))(n+2(t-2))}{2(t-1)^2} \right\rfloor = |B|.$$

Thus for $t > 2$, and any grid graph $G_{m,n}$, there exists a $(t, 2)$ broadcast of size at most $\left\lfloor \frac{(m+2(t-2))(n+2(t-2))}{2(t-1)^2} \right\rfloor$. \square

Our result is nearly optimal. To demonstrate this, we derive the following lower bound as a corollary to Theorem 2.1.

If $G_{m,n}$ is the grid graph with dimensions $m \times n$, and $t > 2$, then

$$\Upsilon_{t,2}(G_{m,n}) \geq \frac{mn}{2(t-1)^2}.$$

Proof. Suppose for contradiction that for some positive integers m, n , and t ,

$$\Upsilon_{t,2}(G_{m,n}) < \frac{mn}{2(t-1)^2}.$$

By assumption, there exists a set S of vertices with $|S| < \frac{mn}{2(t-1)^2}$ that dominates $G_{m,n}$. Thus, we can generate a $(t, 2)$ broadcast on G_∞ by dividing G_∞ into grids with dimensions $m \times n$ and selecting vertices that dominate each grid according to S .

\mathbb{T} is thus an infinite $(t, 2)$ broadcast with broadcast density less than $\frac{mn}{2(t-1)^2}$, which is contradictory by Theorem 2.1. \square

The upper bound of Theorem 1 and the lower bound of Theorem 1 grow at a rate quadratic in m and n , while the difference between them grows at a rate linear in m and n . Thus as m and n increase, the ratio of the two bounds approaches 1. When $t = 3$, Theorem 1 states that

$$\Upsilon_{3,2}(G_{m,n}) \leq \left\lfloor \frac{(m+2)(n+2)}{8} \right\rfloor. \quad (8)$$

This bound differs by a small constant factor from the upper bound on $\Upsilon_{3,2}(G_{m,n})$ provided by Blessing et al. [1]. We thus confirm the conjecture of the authors that their bound is asymptotically optimal.

3 Open Problems

We conclude by providing several open problems and directions for future work.

- In [1] and [4], the authors systematically improve their bounds by a constant value of 4 by adjusting vertices at the corners of $G_{m,n}$. Are similar constant improvements possible for our bounds on $(t, 2)$ broadcast domination numbers?

- In [3], the authors provide an upper bound on the density of optimal $(t, 3)$ broadcasts on G_∞ . By a letterboxing method similar to that employed in this paper, this bound may be translated to an upper bound on $\gamma_{t,3}(G_{m,n})$. However, the original bound is not proven to be optimal, and the resulting bound for finite grids may be off by an amount proportional to the size of the grid. Does this bound approach optimality? Can it be improved?

Acknowledgements

The author expresses his thanks to Pamela E. Harris for her feedback on earlier drafts of this manuscript, and to an anonymous reviewer for careful reading and helpful suggestions.

References

- [1] David Blessing, Katie Johnson, Christie Mauretour, and Erik Insko. “On (t, r) broadcast domination numbers of grids”. In: *Discrete Appl. Math.* 187 (2015), pp. 19–40. ISSN: 0166-218X. URL: <https://doi.org/10.1016/j.dam.2015.02.005>.
- [2] Tony Yu Chang. *Domination numbers of grid graphs*. Thesis (Ph.D.)—University of South Florida. ProQuest LLC, Ann Arbor, MI, 1992, p. 120. URL: http://gateway.proquest.com/openurl?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:dissertation&res_dat=xri:pqdiss&rft_dat=xri:pqdiss:9235073.
- [3] Benjamin F. Drews, Pamela E. Harris, and Timothy W. Randolph. “Optimal (t,r) Broadcasts on the Infinite Grid”. In: *Discrete Applied Mathematics* (2018).
- [4] Michael Farina and Armando Grez. “New Upper Bounds on the Distance Domination Numbers of Grids”. In: *Rose-Hulman Undergraduate Mathematics Journal* 17.2 (2016), p. 7.
- [5] Daniel Gonçalves, Alexandre Pinlou, Michaël Rao, and Stéphan Thomassé. “The domination number of grids”. In: *SIAM J. Discrete Math.* 25.3 (2011), pp. 1443–1453. ISSN: 0895-4801. URL: <https://doi.org/10.1137/11082574>.
- [6] Teresa W. Haynes, Stephen T. Hedetniemi, and Peter J. Slater. *Fundamentals of domination in graphs*. Vol. 208. Monographs and Textbooks in Pure and Applied Mathematics. Marcel Dekker, Inc., New York, 1998, pp. xii+446. ISBN: 0-8247-0033-3.

Timothy W. Randolph

Williams College

twr2@williams.edu