

# 1 The dynamics of root cap sloughing in 2 *Arabidopsis* is regulated by peptide signalling

3 Chun-Lin Shi, Daniel von Wangenheim<sup>1</sup>, Ullrich Herrmann<sup>1</sup>, Mari Wildhagen<sup>1</sup>, Ivan  
4 Kulik, Andreas Kopf, Takashi Ishida, Vilde Olsson, Mari Kristine Anker, Markus Albert,  
5 Melinka A. Butenko, Georg Felix, Shinichiro Sawa, Manfred Claassen, Jiří Friml &  
6 Reidunn B. Aalen\*

---

7 <sup>1</sup> These authors have contributed equally to this work.

8 \*Corresponding author: Reidunn B. Aalen ([reidunn.aalen@ibv.uio.no](mailto:reidunn.aalen@ibv.uio.no))

9 Affiliations:

10 Chun-Lin Shi, Ullrich Herrmann, Mari Wildhagen, Vilde Olsson, Mari Kristine Anker,  
11 Melinka A. Butenko, & Reidunn B. Aalen are or have been at the Section of Genetics  
12 and Evolutionary Biology, Department of Biosciences, University of Oslo, 0316 Oslo,  
13 Norway.

14 Ullrich Herrmann's present address is Plant Developmental Biology and Plant  
15 Physiology, University of Kiel, Am Botanischen Garten 5, 24118 Kiel, Germany

16 Markus Albert and Georg Felix are at the Zentrum für Molekularbiologie der Pflanzen,  
17 University Tübingen, 72076 Tübingen, Germany

18 Ivan Kulik, Daniel von Wangenheim and Jiří Friml are from Institute of Science and  
19 Technology Austria, Am Campus 1, 3400 Klosterneuburg, Austria

20 Daniel von Wangenheim's current address is Centre for Plant Integrative Biology,  
21 University of Nottingham, Loughborough, LE12 5RD, UK

22 Takashi Ishida is at the International Research Organization for Advanced Science and  
23 Technology (IROAST), and Shinichiro Sawa is at the Graduate School of Science and  
24 Technology, of Kumamoto University, Kumamoto 860-8555, Japan Kumamoto  
25 University, Kumamoto 860-8555, Japan

26 Andreas Kopf and Manfred Claassen are from ETH Zurich, HPT E 73, Auguste-Piccard-Hof  
27 1, 8093 Zürich, Switzerland

28

29 **ABSTRACT**

30

31 The root cap protects the stem cell niche of Angiosperm roots from damage. In *Arabidopsis*  
32 lateral root cap (LRC) cell files covering the meristematic zone are regularly abolished  
33 through programmed cell death, while the outermost layer of the root cap covering the tip is  
34 regularly sloughed. Efficient coordination with stem cells producing new layers is needed to  
35 maintain a constant size of the cap. We present a signalling pair, the peptide IDA-LIKE1  
36 (IDL1) and its receptor HAESA-LIKE2 (HSL2), mediating such communication. Live  
37 imaging over several days characterized the sloughing process from initial fractures in LRC  
38 cell files to full separation of a layer. Enhanced expression of IDL1 in the separating root cap  
39 layers resulted in increased frequency of sloughing, balanced with generation of new layers in  
40 a HSL2-dependent manner. Mutations in either *IDL1* or *HSL2* slowed down cell division,  
41 maturation and separation, and suggested involvement of programmed cell death in the  
42 sloughing event. Transcriptome analyses linked IDL1-HSL2 signalling to the transcription  
43 factors BEARSKIN1/2.

44

45 The root cap provides protection for the root apical meristem (RAM) and senses  
46 environmental conditions <sup>1</sup>. The RAM encompasses stem cells, surrounding the Quiescent  
47 Centre (Fig. 1a), both for the tissues of the root proper, and for the root cap. Although both  
48 forms of stem cells continuously produce new cells, the root cap has a defined size, due to  
49 homeostatic balance between generation and loss of root cap cells <sup>2</sup>. Many species shed living,  
50 single cells, so called border cells, but in *Brassicaceae*, including *Arabidopsis thaliana*, the  
51 outermost layer of the root tip is released intact from the underlying cell layer <sup>3</sup>. The  
52 *Arabidopsis* root cap distal to (below) the QC, consists of a core of 5-6 tiers of gravity-sensing  
53 columella cells (COL) surrounded by layers of lateral root cap (LRC) cells (Fig. 1a). The LRC  
54 extends to cover the meristematic zone (MZ), and there the outermost layer regularly  
55 undergoes programmed cell death (PCD) whereby it is replaced by a new LRC layer  
56 developing underneath the old one <sup>4</sup>. Less is known regarding the genetic and molecular  
57 mechanisms governing the detachment of the outer layers of the distal root cap, how the  
58 frequency of sloughing is regulated and how the homeostatic balance with stem cells is  
59 secured <sup>2</sup>.

60 Root cap sloughing requires breakdown of cell walls and is in that respect similar to  
61 other cell separation processes such as flower or fruit abscission <sup>5</sup>, however, sloughing is a  
62 recurrent event in the same organ. Hence, the challenge is to compensate for each lost root  
63 cap layer by generation of a new, and to detach the oldest layer when a new layer has been  
64 formed. In the distal root cap, this requires communication between old separating layers and  
65 the stem cells of COL and LRC, as well as a sequential coordination of generation, maturation  
66 and separation of root cap cells.

67 Mutations in four genes encoding closely related NAC (NAM, ATAF, CUC)  
68 transcription factors (TFs) – FEZ, SOMBRERO (SMB), BEARSKIN1 (BRN1) and BRN2 –  
69 have provided insights into the development and differentiation of the root cap <sup>4,6-8</sup>. FEZ is

70 implicated in the formative division of the common stem cells for the LRC and the epidermal  
71 layer (EPI). SMB regulates differentiation and maturation<sup>6-8</sup>, while the double mutant *brn1*  
72 *brn2* is impaired in cell wall degradation and accumulates additional undetached root cap  
73 layers<sup>8</sup>.

74 Downstream of these TFs, genes have been identified that encode enzymes involved  
75 cell wall remodelling and synthesis<sup>8</sup>. Similar genes have been discovered in other cell  
76 separation processes in flowering plants<sup>9,10</sup>. Floral organ abscission and emergence of the  
77 lateral root through the overlying tissues in *Arabidopsis*, are regulated by the small peptide  
78 INFLORESCENCE DEFICIENT IN ABSCISSION (IDA) that signals through two closely  
79 related leucine-rich repeat receptor-like kinases (LRR-RLKs), HAESA (HAE) and HAESA-  
80 LIKE2 (HSL2)<sup>11-15</sup>. *IDA* belongs to a family of *IDA-LIKE* (*IDL*) genes, encoding  
81 preproteins with a common C-terminal motif (Supplementary Fig. 1), hypothesized to have  
82 a function similar to *IDA*. Here, we address the function in root cap sloughing of *IDL1*,  
83 expressed in the root tip, and identify HSL2 as its receptor. Rather than specifically  
84 controlling the cell separation step of the sloughing process, this signalling pair regulates the  
85 dynamics of root cap detachment and generation of new root cap layers.

86

## 87 **RESULTS**

### 88 **Identification of HSL2 as the receptor of IDL1**

89 *IDL1* is prominently expressed in COL cells at the centre of the two adjacent root cap layers  
90 of the primary root that will undergo separation (Fig. 1a-b). *IDL1* can rescue the abscission  
91 deficiency of the *ida* mutant<sup>16</sup>, indicating that *IDL1* peptide can interact with the receptors of  
92 *IDA*. Using promoter- $\beta$ -glucuronidase (*GUS*) reporter lines for *HAE* and *HSL2* we found that  
93 only *pHSL2:GUS* had an expression pattern in the primary root tip, with the strongest  
94 expression in the youngest LRC cell files (Supplementary Fig. 1). This was confirmed when

95 using a construct with the *HSL2* promoter coupled to a nuclear targeted fluorescent protein  
96 (pHSL2::Venus-H2B) (Fig. 1c). The N-terminal signal sequence of the IDL1 prepropeptide  
97 (Supplementary Fig. 1) indicates that the mature peptide is exported and can move in the  
98 apoplastic space to interact with the extracellular domain of the preferred receptor<sup>17,18</sup>. Thus,  
99 the expression patterns were compatible with a hypothesized function of IDL1 and HSL2 as a  
100 ligand-receptor pair involved in communication between cells in the root cap.

101 To investigate whether IDL1 could activate and bind the receptor, we took advantage  
102 of methods previously used for IDA and HSL2<sup>12</sup>. Addition of a synthetic IDA peptide of 12  
103 amino acids, with hydroxylation of the central proline residue (mIDA), to *Nicotiana*  
104 *benthamiana* leaf pieces expressing the HSL2, results in an immediate receptor activation  
105 measurable as oxidative burst. The mIDA peptide (Fig. 1d) activates HSL2 down to a half-  
106 maximal efficient concentration (EC50) of 1 nM<sup>12</sup>. The corresponding hydroxylated dodeca  
107 IDL1 peptide (mIDL1, Fig. 1d), tested in the same assay, functioned also in a dose-dependent  
108 manner with a similar efficiency (EC50 ~3 nM, Fig. 1e).

109 Competition assays with acridinium-labelled and unlabelled IDA-derived peptides  
110 have demonstrated specific and reversible binding of the ectodomain of HSL2 expressed in *N.*  
111 *benthamiana* leaves<sup>12</sup>. The same experimental set-up showed that unlabelled mIDL1  
112 competed for binding of the HSL2 ectodomain with an efficiency similar to the unlabelled  
113 IDA peptide, with a half-maximal inhibitory concentration (IC50) of ~30 nM (Fig. 1f).

114 Root cap expression, together with binding and activation of HSL2 by IDL1,  
115 suggested that this peptide and receptor might serve as a root cap signalling module.

116

### 117 ***IDL1* expression increases the frequency of sloughing in a HSL2-dependent manner**

118 Since a number of LRR-RLKs with high similarity in their ectodomains are expressed in the  
119 root<sup>19</sup>, ectopic ligand expression or exogenously applied synthetic peptides might lead to

120 interaction with non-native receptors<sup>20</sup>. With the aim of testing whether IDL1-HSL2 was  
121 specifically involved in root cap sloughing, we therefore used the promoter of *IDL1* itself to  
122 generate Enhanced IDL1 (EnhIDL1) plants with estradiol (ES)-inducible enhanced expression  
123 of both *IDL1* and *GUS* (Fig. 2a-b)<sup>21</sup>. EnhIDL1 plants were crossed to an *hsl2* knockout  
124 mutant<sup>16</sup> to test the importance of HSL2 for IDL1 function. Cell-specific and efficient ES-  
125 induction was confirmed by strongly enhanced GUS expression in EnhIDL1 roots both in  
126 wild type Col-0 and *hsl2* mutant background as compared to the *pIDL1:GUS* expression  
127 pattern and by qRT-PCR (Fig. 1b, 2b and Supplemental Fig. 1). These lines did not deviate in  
128 root length (Supplementary Fig. 1).

129 Cell elongation, cell division and differentiation take place in the root tip during the  
130 first days after germination. To ensure that homeostatic balance between gain and loss of root  
131 cap layers had been established, our experiments were performed on roots with full-sized  
132 RAM and root cap (Supplementary Fig. 2). Seven days after germination (DAG) roots of Col-  
133 0, EnhIDL1 and EnhIDL1 *hsl2* seedlings grown vertically on agar with 10  $\mu$ M ES, covered by  
134 a cover slip, were tracked, using a laser scanning microscope with a vertical stage<sup>22</sup>. Images  
135 taken in two experiments every 15 minutes for 64 and 70 hours (h) revealed a stepwise  
136 sloughing process (Fig. 2c-e). The first signs of a forthcoming sloughing event were  
137 characteristic fractures or gaps in LRC cell files in a region just above the Quiescent Centre  
138 (QC) and about 5h later at the level of the youngest starch-filled COL tiers (Fig. 2c, red and  
139 yellow arrows). These gaps initiated a gradual loosening of the outermost LRC layer towards  
140 the tip. Finally, the COL cells at the centre of the cap separated from the layer underneath  
141 followed by detachment of the whole layer as it slid to the side and was lost (Fig. 2c, white  
142 arrow and Supplementary Video 1). On average, 0.6 root cap layers were shed per root per  
143 day in Col-0 (Fig. 2f), with an average of ~18h for a completed sloughing process, and

144 thereafter another ~18h before a new event of root cap layer separation would be initiated (Fig.  
145 2g and Supplementary Table 1).

146 The sloughing frequency in EnhIDL1 was 50 % higher than for Col-0, i.e. 0.9 events  
147 per root per day (Fig. 2f). This increase seemed to be caused by a shorter interval between the  
148 completion of one event and the initiation of the next (Fig. 2g and Supplementary Table 1),  
149 and overlapping events with fractures in the second outermost layer, before the previous event  
150 was fully completed (Fig. 2d, green arrow, 2h, and Supplementary Video 2).

151 In *hsl2* background the increased sloughing frequency of EnhIDL1 was rescued and  
152 did not deviate from that of Col-0 (Fig. 2f). However, the sloughing process developed most  
153 often differently in that the outermost layer was split open close to the tip at the border  
154 between the COL and LRC cells (Figure 2e and h). The split cap layer, sometimes leaving the  
155 central COL region behind, seemed to be still attached to the LRC sheet surrounding the  
156 meristematic region (Fig. 2e, white arrows, and Supplementary Video 3).

157 Live imaging was supplemented with data for roots grown on vertical agar plates. In  
158 this set-up, shed layers of the distal cap, recognizable by detached LRC files, did not slide  
159 away immediately, but accumulated in front of the growing roots, most likely due to less  
160 friction when the roots were not covered by a cover slip. Caps were, however, easily lost  
161 when removed from the agar, and sloughed caps were therefore counted directly on the plates  
162 at 6, 8 and 10 DAG, with and without 10  $\mu$ M ES (Fig. 3a-b). Col-0, EnhIDL1 *hsl2*, and  
163 EnhIDL1 on non-inducing medium sloughed about 0.6 caps per day (Fig. 3a-b). On ES  
164 medium, on the other hand, EnhIDL1, accumulated up to eight sloughed root cap layers  
165 (Supplementary Fig. 3) and differed significantly with an average loss of one root cap layer  
166 per day (Fig. 3a). Thus, these results were in full accordance with the live imaging results.

167

168

169 **Higher frequency of IDL1-induced sloughing is compensated by more stem cell divisions**

170 We questioned whether a higher frequency of sloughing reduced the number of attached  
171 layers, however, all three lines maintained stably ~6 attached COL tiers (including the stem  
172 cell layer) (Supplementary Fig. 3), and did not differ significantly from each other (Student's  
173 t-test  $P \geq 0.1$ ), irrespective of ES in the growth medium (Fig. 3c).

174 More frequent shedding, but at the same time constant cell number and size of the root  
175 cap, imply an increased frequency of new initiations of COL tiers and LRC cell files to  
176 compensate for the lost caps. Recent divisions of COL stem cells, positioned just below the  
177 QC (Fig. 1a), can be recognized by the smaller cell size in the new 1st COL tier and their  
178 weaker cell wall staining with propidium iodide (PI) compared to older COL cells<sup>6</sup>. The COL  
179 stem cells are surrounded by the shared stem cells for LRC and epidermal cell files, and in  
180 medial optical sections of the root tip, they are seen as four COL stem cells with one EPI/LRC  
181 stem cell on each side (Fig. 1a). New epidermal cells are generated by anticlinal division,  
182 while periclinal divisions represent initiation of a new LRC cell file. On inductive medium,  
183 cell wall patterns indicating recent cell divisions were found for COL in 80% and for LRC in  
184 60% of EnhIDL1 PI-stained root tips (Fig. 3d-e). This was in contrast to Col-0 and even more  
185 so for EnhIDL1 *hsl2*, where < 15 % of the root tips showed signs of recent stem cell divisions  
186 (Fig. 3d-e).

187 These findings imply that increased expression level of *IDL1* not only leads to  
188 increased sloughing, but also an increased frequency of initiation of new root cap layers.

189

190 **Mutations in *IDL1* and *HSL2* slow down the sloughing process**

191 To further explore the function of the IDL1-HSL2 signalling in the root cap, we investigated  
192 the sloughing behaviour of *hsl2*<sup>16</sup> and an *idll* CRISPR/Cas9 mutant line (Supplementary Fig.  
193 4) compared to Col-0. While the number of attached layers remained close to 6 for both Col-0



194 and the mutants, a subtle, but significant difference was observed regarding the number of  
195 sloughed cap layers at 10 DAG with on average 3.0 caps for Col-0 and 2.4 for the mutants  
196 (Fig. 4a). Inspection of the youngest LRC cell files furthermore suggested that the frequencies  
197 of generation of new LRC cell layers had slowed down in both mutants. After the initial  
198 periclinal division of a EPI/LRC stem cell, the new LRC cell file is extended by anticlinal  
199 divisions. The new file will be neighbouring an older and longer LRC cell file on the lower  
200 side and an epidermal cell file on the upper side until the next new LRC layer is initiated (Fig.  
201 4b-d). In the wild type, one cell of the two youngest cell files neighboured the epidermal cell  
202 file, in contrast to 2-3 cells in both the *hsl2* and *idl1* mutants (Fig. 4b-c). This is an expected  
203 result if the periclinal divisions initiating a new LRC file take place less frequently relative to  
204 the anticlinal divisions of already initiated LRC cell files (Fig. 4d).

205

#### 206 **Mutation in *HSL2* affects genes in both the LRC and the COL cell layers**

207 To gain a better understanding of the function of the signalling initiated by activation of *HSL2*,  
208 RNA sequencing was performed on RNA isolated from the RAM of root tips of Col-0 and  
209 *hsl2*. In line with the subtle phenotype of *hsl2*, few genes were grossly affected by the  
210 mutation. However, using a P adjusted (P<sub>adj</sub>) cut-off value of 0.1, 156 genes were identified  
211 with an expression level < 70 % of the Col-0 level (Supplementary Table 2). In an attempt to  
212 identify where the genes affected by HSL2 signalling were expressed, we took advantage of a  
213 data set for tissue-specific gene expression in the Arabidopsis RAM<sup>23</sup>. This dataset provides  
214 information on differential gene expression in the different root tissue, QC, LRC, COL, EPI,  
215 cortex (COR) and endodermis (END), based on cell sorting using marker genes for these  
216 tissues<sup>23</sup>. Eighty-one of the *hsl2* down-regulated genes were identified as differentially  
217 expressed according to this data set. We defined a gene as characteristic for a specific tissue if  
218 it had the highest expression level for this tissue compared all the other tissues. Evaluation

219 according to a binomial test ( $p=0.1$ , see Methods for details) found significant enrichment of  
220 genes expressed in LRC and COL (Table 1). Genes encoding extracellular proteins were well  
221 represented among the LRC expressed genes, while many genes most highly expressed in  
222 COL encoded proteins involved in catalytic and hydrolytic activity including starch and cell  
223 wall degradation.

224         Activation of RLK receptors is transduced via phosphorylation cascades and  
225 activation of transcription factors, which at present are unknown for HSL2 in the root tip.  
226 However, we found an overlap between the *hsl2* down-regulated genes and genes identified in  
227 a recent transcriptome analysis of NAC TFs <sup>8</sup>. Of sixty genes reported to be positively  
228 regulated by overexpression of SMB and downregulated in the triple mutant *smb brn1 brn2*,  
229 twenty-five genes, a highly significant number (Fischer's exact test  $P < 10^{-10}$ ), was also down  
230 regulated in *hsl2* mutant root tips (Supplementary Table 2). There was an overrepresentation  
231 of genes encoding extracellular proteins and enzymes, hereunder enzymes involved in cell  
232 wall remodelling and starch metabolism (Supplementary Table 2) <sup>8,24,25</sup>.

233         The three NAC TFs were not significantly down regulated in *hsl2* according to our  
234 RNAseq analyses, however, to investigate whether IDL1 signalling could affect expression  
235 patterns, lines with translational fusion constructs for BRN1 and BRN2 were crossed into the  
236 EnhIDL1 line and the *idl1* mutant. Both *BRN1* and *BRN2* are expressed in the outer LRC cell  
237 files <sup>8</sup>. In EnhIDL1, the expression pattern of *pBRN1:BRN1-GFP* was very similar to Col-0  
238 grown with ES in the medium, while *pBRN2:BRN2-GFP* consistently showed an additional  
239 layer of expressing cells at the tip (Supplementary Fig. 5). In the *idl1* mutant, on the other  
240 hand, *pBRN2:BRN2-GFP* expression was similar to Col-0, but fewer cells expressed  
241 *pBRN1:BRN1-GFP* in the LRC cells (Supplementary Fig. 5).

242         The *ROOT CAP POLYGALACTURONASE (RCPG)*, a direct target of BRN1 <sup>8</sup>, was of  
243 the most strongly downregulated genes both in the triple mutant *smb brn1 brn2* and *hsl2*

244 (Supplementary Table 2). In loosened Col-0 root cap layers the construct pRCPG:nYG, with  
245 nuclear targeted YFP driven by the *RCPG* promoter<sup>8</sup>, was expressed in more cells and with  
246 higher intensity than in *hsl2* and *idl1* background (Fig. 5a). Moreover, pRCPG:nYG was  
247 expressed in undetached LRC cells in the mutants (Fig. 5a), suggesting delayed cell  
248 separation. pRCPG:nYG expression was also seen in undetached root cap layers in the  
249 EnhIDL1 line, in this case suggesting premature expression (Supplementary Fig. 5).

250 This influence on *RCPG* expression suggests that IDL1-HSL2, like the BRN TFs, are  
251 involved in the final step of the sloughing process – the actual cell separation.

252

### 253 **IDL1-HSL2 regulates *BFN1* expression at PCD Site II**

254 We noted that in contrast to *hsl2* and *idl1*, most Col-0 cells expressing *RCPG* had initiated the  
255 sloughing process and their nuclei stained with PI (Fig. 5a and Supplementary Fig. 5), which  
256 is indicative of plasma-membrane permeabilization. Such PI entrance has been used as a  
257 hallmark of initiation of developmental programmed cell death (dPCD) associated with the  
258 recurrent turnover of the LRC covering the MZ<sup>4</sup>. Interestingly, in the *hsl2* transcriptome five  
259 out of eight genes strongly associated with dPCD were down-regulated; the *BIFUNCTIONAL*  
260 *NUCLEASE1 (BFN1)*, *METACAPASE9 (MC9)*, *SAPOSIN-LIKE ASPARTYL PROTEASE3*  
261 (*PASPA3*), *EXITUS (EXI)* and *RIBONUCLEASE3 (RNS3)*<sup>4,26</sup> (Supplementary Table 2). To  
262 investigate the potential involvement of IDL1-HSL2 in dPCD in the root cap, a transcriptional  
263 fusion construct with the *BFN1* promoter and a nuclear-targeted GFP (*BFN1<sub>pro</sub>:nGFP*)<sup>27</sup> was  
264 crossed into EnhIDL1 and the mutants. In Col-0, two known sites of dPCD, at the upper end  
265 of the MZ (Site I) and close to the QC (Site II), were identified as expected (Fig. 5b).

266 Interestingly, in addition to the strong expression of pHSL2::Venus-H2B in young LRC cell  
267 files, HSL2-driven nuclear-targeted Venus was found in LRC cell files surrounding the MZ  
268 and stretching past Site II (Fig. 5c). A third PCD site was found in loosened LRC cell files

269 consistent with the high number of nuclei there staining with PI (Fig. 5a-b), and suggestive of  
270 PCD in the sloughed cells. The pattern of BFN1<sub>pro</sub>:nGFP expression in EnhIDL1 at these sites  
271 was similar to Col-0. In contrast, *idl1* and *hsl2* root tips had significantly fewer cells  
272 expressing the BFN1<sub>pro</sub>:nGFP construct, in particular at Site II (Fig. 5b and Supplementary  
273 Fig. 6). Interestingly, Site II coincides with the position of the fractures initiating the  
274 detachment of the distal root cap (Fig. 5b), which suggestively may provide a molecular link  
275 between the IDL1-HSL2 signalling pair and initiation of the sloughing process.

276

## 277 **DISCUSSION**

278 Dynamics of root cap sloughing and mechanisms securing homeostatic balance between loss  
279 and gain of layers of the distal root cap have been difficult to study because of the duration of  
280 the process, the easily lost, small cap layers, and few available mutants<sup>2</sup>. Continuous live  
281 imaging over three days, analyses of transcriptome data and cellular expression patterns have  
282 enabled us to detail the different phases of root cap sloughing and the involvement of IDA-  
283 HSL2 in this process (Fig. 6). Under our conditions, 0.6 distal cap layers were shed per root  
284 per day in Col-0. Enhanced expression of *IDL1* increased this frequency by shortening the  
285 interval from one event to the next, or by detaching two layers at the same time. We have  
286 identified IDL1 as a peptide ligand that in nanomolar concentrations binds and activates the  
287 HSL2 receptor. Both IDL1-increased sloughing and the balancing increase in stem cell  
288 divisions were dependent on HSL2, i.e. IDA-HSL2 signalling controls the dynamics of  
289 sloughing at the root tip.

290 We had expected cell separation phenotypes of *idl1* and *hsl2* similar to the total  
291 deficiency in floral organ abscission seen in the *ida* mutant and the double mutant *hae hsl2*<sup>16</sup>.  
292 However, homeostasis appears maintained in the two mutants, albeit at a much lower  
293 frequency. Cell division, maturation and separation were slowed down, but not abolished. It is

294 likely that other factors, including other peptides and receptors, are involved in the separation  
295 step. A few mutants have been identified that release single cells instead of entire root cap  
296 layers, like *nin-like protein 7 (nlp7)*, *quasimodo1-1 (qua1-1)* and *qua2-1*<sup>25,28</sup>. In contrast, the  
297 *brn1 brn2* double mutant, and *active quiescent center1-1 (aqc1-1)*, are delayed in separation  
298 of root cap layers<sup>8,29</sup>. The latter mutant is defect in the gene encoding tyrosylprotein  
299 sulftransferase (TPST), an enzyme responsible for sulfation of a number of peptide ligands.  
300 The disorganized root cap and stunted growth in this mutant is likely to be a consequence of  
301 several dysfunctional peptides lacking sulfation.

302         Live imaging of roots and marker-gene expression in the EnhIDL1 line and the *idl1*  
303 and *hsl2* mutants suggest that IDL1-HSL2 signalling is controlling initiation of the sloughing  
304 process. Intriguingly, the region where the first formative fractures or gaps in the LRC cell  
305 files are found, seems to coincide with dPCD Site II, and a speculative suggestion is that these  
306 fractures are created by PCD. dPCD starting from Site I is involved in the recurrent removal  
307 of the LRC surrounding the MZ<sup>4</sup>. Differential control of dPCD Sites I and II would make it  
308 possible for the plant to operate with different frequencies of renewal of the LRC surrounding  
309 the MZ and of sloughed layers at the tip.

310         IDL1-HSL2 control of the position of initiation may also explain the split caps still  
311 connected to the LRC surrounding the MZ observed for EnhIDL1 in *hsl2* background. With  
312 fewer fractures, MZ cell division and COL cell elongation is likely to exert mechanical stress  
313 that tears apart cells at other, weaker sites. Similarly, in *ida* and *hae* mutants, separation of  
314 cells in the cortex and epidermis overlying developing lateral root primordia (LRPs) is  
315 impaired, but the LRPs eventually emerge forcibly through these tissues<sup>10</sup>. These two  
316 examples suggest that regulation of growth of the LRP and the primary root is not tightly  
317 coupled to the respective cell separation events.

318 BRN1, BRN2 and SMB are implicated in root cap maturation and root cap detachment  
319 <sup>7,8</sup>. pHSL2::Venus-H2B was found to have overlapping expression pattern with the BRN TFs  
320 in the LRC, and a subset of genes regulated by these TFs was also found downstream in the  
321 HSL2 signalling pathway. Differences in spatial expression patterns compared to Col-0 of  
322 transcriptional and translational reporter construct crossed into our lines, indicated faster  
323 maturation in EnhIDL1 and slower maturation in the mutants. Together this suggests HSL2 to  
324 play a role in maturation and cell layer separation in addition to the role in initiation. *BFNI<sub>pro</sub>*-  
325 controlled marker-gene expression and PI-stained nuclei in LRC cells of root caps undergoing  
326 sloughing, indicate that root cap sloughing involves both cell separation and cell death.

327 RNAseq data for *hsl2* combined with publicly available information on differential  
328 gene expression in root tissues were used in an attempt to identify the cell types that primarily  
329 respond to HSL2 activation. This analysis identified overrepresentation of genes preferably  
330 expressed in LRC and COL cells. The major effect of receptor activation will occur in the  
331 cells where the receptor is expressed, indicating that HSL2 is present and responsive for  
332 activation both in LRC and COL cells. Additionally, this analysis suggests that activation of  
333 HSL2 gives different outcomes in different tissues.

334 IDL1-HSL2 communication facilitates dynamic regulation of the homeostatic balance  
335 between stem cell division and sloughing activity. Since the root cap is guiding root growth  
336 by sensing and responding to external cues, like gravity, water potential or the presence of  
337 obstacles in the soil <sup>1,2</sup>, it is highly likely that the frequency of root cap shedding is triggered  
338 by environmental factors, possibly by influencing the expression level of IDL1 and/or HSL2.  
339 At present, it is not known whether or how external condition might feed into the circle of  
340 generation and loss of root cap layers. Factors involved in transcriptional regulation of *IDL1*  
341 and *HSL2*, peptide processing and export, and interacting partners have not been identified.  
342 Additionally, several parallel pathways partake in regulation of stem cell activity, the NAC

343 transcription factors FEZ and SMB, the RETINOBLASTOMA-RELATED protein, the  
344 WOX5 transcription factor, and the AUXIN RESPONSE FACTORS ARF10 and ARF16<sup>30,32</sup>,  
345 but their relationship to the IDL1-HSL2 signalling pathway have not yet been unravelled.

346 It will be exciting to take advantage of root tip tracking by live imaging and to explore  
347 these relations in the future.

348

349

## 350 **Methods**

351 **Accession numbers** of the genes studied in this work: *IDL1* – At3g25655, *HSL2* –  
352 At5g65710; *HAE* – At4g28490; *BRN1* – At1g33280, *BRN2* – At4g10350, *BFN1* – AT1G11190  
353 and *RCPG*– At1g65570.

354 **Constructs and plant lines.** The *pIDL1:GUS*, *pHAE:GUS*, *pHSL2:GUS*, *BFN1<sub>pro</sub>:nGFP*,  
355 *pRCPG:nYG*, *pBRN1:BRN1-GFP* and *pBRN2:BRN2-GFP* lines have been described  
356 previously<sup>8,16,27</sup>. A 2300 bp long fragment upstream of the start codon of *HSL2*<sup>31</sup> was cloned  
357 into destination vector *promoter::Venus-H2B*<sup>32</sup>, using the Invitrogen Gateway<sup>TM</sup> system, and  
358 introduced into *Arabidopsis* Col-0 plants by floral dipping. A single-locus homozygote  
359 *pHS2::Venus-H2B* line was identified.

360 Gateway cloning and floral dipping was also used generate Enhanced IDL1  
361 (EnhIDL1) plants; the *IDL1* coding region was inserted after the estradiol inducible promoter  
362 in the vector pMDC221, and a 1555 bp *IDL1* promoter fragment was cloned in front of the  
363 XVE transcription factor in the pLB12 vector<sup>21</sup>. Of six *pIDL1 pLB12* lines checked for GUS  
364 expression pattern (after induction) comparable to that of the published *pIDL1:GUS*<sup>10,30</sup>, a  
365 single-locus line (checked on 25 mg/l Hygromycin (Hyg)), was crossed to a single-locus *IDL1*

366 *pMDC221* line (checked on 50 mg/l Kanamycin (Km)). EnhIDL1 plants homozygous for both  
367 constructs were identified by selection on Hyg and Km and checked non-segregating GUS  
368 expression. Of two lines with comparable sloughing behaviour, GUS expression level and  
369 pattern, one was chosen for the experiments shown here. This line was furthermore used for  
370 all crosses to mutants and marker-lines.

371 Two *idll* mutants were made with CRISPR/Cas9 technology using an IDL1-targeting  
372 vector construct based on the pDe-Cas9 systems provided by Holger Puchta<sup>33</sup>. A 20 base pair  
373 gRNA target site, unique for IDL1 including BslI recognition motif, was selected for the  
374 mutagenesis. Vector construction and generation of mutants were performed as described  
375 previously<sup>34</sup>. Through selections with BASTA and CAPS analyses, mutants containing 2 bps  
376 deletion (*idll-cr1*) and 4 bps deletion (*idll-cr2*) in the open reading frame of *IDL1* were  
377 isolated. Both mutations result in frame shifts generating premature stop codons in the central  
378 variable region of IDL1 prepropeptide and the C-terminal peptide motif is thereby deleted  
379 (Supplementary Fig. 4). Phenotypic differences between the alleles were not observed  
380 (Supplementary Fig. 4). Data for *idll-cr1* (named *idll* here) is presented.

381 Primers for genotyping and generation of constructs are listed in Supplementary Table  
382 3.

383 **Growth conditions and registration of number of root cap layers.** Sterilized seeds were  
384 stratified at 4 °C for 72 h. Seedlings were grown at 22 °C, 8 h dark and 16 h light (100 µE/m<sup>2</sup>  
385 light intensity) on square vertical plates with sucrose-free 0.5 x MS medium adjusted to pH  
386 6.0, with 10µM 17-β-estradiol for induction of enhanced expression of *IDL1* in EnhIDL1  
387 lines<sup>21</sup>. Sloughed root cap layers were counted and imaged directly on the agar plates (using a  
388 Zeiss Axioplan2 imaging microscope) as they easily fall off when roots are moved. Attached  
389 caps were imaged and counted after immersion in Lugol solution (Sigma 6265) for 1 minute  
390 for detection of the COL cell tiers with amyloplasts (Supplementary Fig. 3). No significant



391 difference in length was observed between genotypes or with ES in the medium  
392 (Supplementary Fig. 1).

393 **Oxidative burst measurements and receptor-ligand binding assay.** *N. benthamiana* leaves  
394 transiently expressing HSL2 constructs were used for oxidative burst measurements and  
395 peptide binding assays as previously described<sup>12</sup>. EC50 and IC50 curves are representative  
396 for N=3 independent experiments, each with three technical replicates.

397 **Microscopy.** Histochemical GUS assays were performed as previously described<sup>16</sup>. Confocal  
398 laser scanning microscopy (CLSM) was carried out with Olympus FV1000 and ZEISS  
399 LSM880 confocal microscopes at the Oslo NorMIC imaging node. Roots were stained with  
400 10  $\mu$ M propidium iodide (PI).

401 **For live imaging** seeds were surface sterilized by chlorine gas, sown on 1% agar AM-  
402 medium (0.5 x MS, pH 5.8, 1% agar), stratified for 3 days at 4°C and cultivated in a growth  
403 incubator at 22°C in a 16/8 h day/night cycle with 120–140  $\mu$ mol/m<sup>2</sup>/s light.

404 Eight day-old plants were put on blocks (rectangular parallelepipeds) of 1.5% agar AM-  
405 medium (0.5 x MS, pH 5.8) supplemented with 10 $\mu$ M 17- $\beta$ -estradiol. Roots and shoots were  
406 put on the blocks' basis and sides, respectively<sup>22</sup>. These samples were put into the chambered  
407 cover-glass (Thermo Scientific Nunc, catalogue no: 15536), which were then mounted onto a  
408 laser scanning microscope with a vertical stage (roots down, shoots up). Photographs of root  
409 tips were taken every 15 minutes, in two experiments over 64 and 70 h, with in total 9 roots  
410 for Col-0 and 10 roots for EnhIDL1 and EnhIDL1 *hsl2*.

411 **RNA extraction and RNAseq analysis.** Total RNA was extracted in three replicates from 0.5  
412 mm segments of primary root tips of 7-day-old Col-0 and *hsl2* seedlings using  
413 Trizol/chloroform extraction follow by purification using the RNeasy Plant Mini Kit  
414 (Qiagen). The quality of the RNA was ascertained using the Agilent 2100 Bioanalyzer.

415 RNAseq was performed at the Functional Genomics Center Zurich, Switzerland. Deseq2 with  
416 standard parameters was used to analyse expression levels  
417 (<https://bioconductor.org/packages/3.7/bioc/vignettes/DESeq2inst/doc/DESeq2.html>).

418 **Statistical methods** *Student's t-test* (2-tailed) was used to analyse whether root cap  
419 phenotype and marker-gene expression in different lines deviated significantly from Col-0 cf.  
420 Fig. 2, 3 and 4, and Supplementary Fig. 1, and 6. If not otherwise stated in the figure legends  
421  $\geq 10$  roots were used for each genotype, treatment and age. Two to five biological independent  
422 repeats were used. In all graphs, standard deviations are shown as bars.

423 *Binomial test* was applied to test for enrichment of maximal expression level for a specific  
424 tissue across all other tissues. The probability to find  $k$  maximal expressions out of  $N$  genes in  
425 total is given by

$$P(X \geq k) = \sum_{i=k}^N \binom{N}{i} p^i (1-p)^{N-i}$$

426 whereby  $p$  is the per gene probability to be maximal and  $\binom{N}{i}$  is the binomial coefficient.

427 *Fisher's exact test* was used to calculate the likelihood of overlap of regulated genes by  
428 *SMBox /smb brn1 brn2* (NAC TFs) and *hsl2*.

429

430 **References**

- 431 1 Arnaud, C., Bonnot, C. m., Desnos, T. & Nussaume, L. The root cap at the forefront.  
432 *Comp. Rendus Biol.* **333**, 335-343 (2010).
- 433 2 Kumpf, R. P. & Nowack, M. K. The root cap: a short story of life and death. *J. Exp.*  
434 *Bot.* **66**, 5651-5662 (2015).
- 435 3 Driouich, A., Durand, C., Cannesan, M. A., Percoco, G. & Vire-Gibouin, M. Border  
436 cells versus border-like cells: are they alike? *J. Exp. Bot.* **61**, 3827-3831 (2010).
- 437 4 Fendrych, M. *et al.* Programmed cell death controlled by ANAC033/SOMBRERO  
438 determines root cap organ size in *Arabidopsis*. *Curr. Biol.* **24**, 931-940 (2014).
- 439 5 Lewis, M. W., Leslie, M. E. & Liljegren, S. J. Plant separation: 50 ways to leave your  
440 mother. *Curr. Opin. Plant Biol.* **9**, 59-65 (2006).
- 441 6 Willemsen, V. *et al.* The NAC domain transcription factors FEZ and SOMBRERO  
442 control the orientation of cell division plane in arabidopsis root stem cells. *Devel. Cell*  
443 **15**, 913-922 (2008).
- 444 7 Bennett, T. *et al.* SOMBRERO, BEARSKIN1, and BEARSKIN2 regulate root cap  
445 maturation in *Arabidopsis*. *Plant Cell* **22**, 640-654 (2010).
- 446 8 Kamiya, M. *et al.* Control of root cap maturation and cell detachment by BEARSKIN  
447 transcription factors in *Arabidopsis*. *Develop.* **143**, 4063-4072 (2016).
- 448 9 Sundaresan, S. *et al.* De novo transcriptome sequencing and development of  
449 abscission zone-specific microarray as a new molecular tool for analysis of tomato  
450 organ abscission. *Front. Plant Sci.* **6**, 1258 (2015).
- 451 10 Kumpf, R. P. *et al.* Floral organ abscission peptide IDA and its HAE/HSL2 receptors  
452 control cell separation during lateral root emergence. *Proc. Natl. Acad. Sci. USA* **110**,  
453 5235-5240 (2013).

- 454 11 Aalen, R. B., Wildhagen, M., Stø, I. M. & Butenko, M. A. IDA: a peptide ligand  
455 regulating cell separation processes in *Arabidopsis*. *J. Exp. Bot.* **64**, 5253-5261 (2013).
- 456 12 Butenko, M. A. *et al.* Tools and strategies to match peptide-ligand receptor pairs.  
457 *Plant Cell* **26**, 1838-1847 (2014).
- 458 13 Estornell, L. H. *et al.* The IDA peptide controls abscission in arabidopsis and citrus.  
459 *Front. Plant Sci.* **6**, 1003 (2015).
- 460 14 Santiago, J. *et al.* Mechanistic insight into a peptide hormone signaling complex  
461 mediating floral organ abscission. *Elife* **5**, doi:10.7554/eLife.15075 (2016).
- 462 15 Patharkar, O. R. & Walker, J. C. Core Mechanisms regulating developmentally timed  
463 and environmentally triggered abscission. *Plant Physiol.* **172**, 510-520 (2016).
- 464 16 Stenvik, G. E. *et al.* The EPIP peptide of INFLORESCENCE DEFICIENT IN  
465 ABSCISSION is sufficient to induce abscission in *Arabidopsis* through the receptor-  
466 like kinases HAESA and HAESA-LIKE2. *Plant Cell* **20**, 1805-1817 (2008).
- 467 17 Matsubayashi, Y. Post-Translational modifications in secreted peptide hormones in  
468 plants. *Plant Cell Physiol.* **52**, 5-13 (2011).
- 469 18 Butenko, M. A., Vie, A. K., Brembu, T., Aalen, R. B. & Bones, A. M. Plant peptides  
470 in signalling: looking for new partners. *Trends Plant Sci.* **14**, 255-263 (2009).
- 471 19 ten Hove, C. A. *et al.* Probing the roles of LRR RLK genes in *Arabidopsis thaliana*  
472 roots using a custom T-DNA insertion set. *Plant Mol. Biol.* **76**, 69-83 (2011).
- 473 20 Yamada, M. & Sawa, S. The roles of peptide hormones during plant root development.  
474 *Curr. Opin. Plant Biol.* **16**, 56-6 (2013).
- 475 21 Brand, L. *et al.* A versatile and reliable two-component system for tissue-specific gene  
476 induction in *Arabidopsis*. *Plant Physiol.* **141**, 1194-1204 (2006).
- 477 22 von Wangenheim, D. *et al.* Live tracking of moving samples in confocal microscopy  
478 for vertically grown roots. *Elife* **6**, doi:10.7554/eLife.26792 (2017).

- 479 23 Nawy, T. *et al.* Transcriptional profile of the arabidopsis root quiescent center. *Plant*  
480 *Cell* **17**, 1908-1925 (2005).
- 481 24 del Campillo, E., Abdel-Aziz, A., Crawford, D. & Patterson, S. E. Root cap specific  
482 expression of an endo-beta-1,4-D-glucanase (cellulase): a new marker to study root  
483 development in *Arabidopsis*. *Plant Mol. Biol.* **56**, 309-323. (2004).
- 484 25 Karve, R., Suarez-Roman, F. & Iyer-Pascuzzi, A. S. The transcription factor NIN-  
485 LIKE PROTEIN7 controls border-like cell release. *Plant Physiol.* **171**, 2101-2111,  
486 doi:10.1104/pp.16.00453 (2016).
- 487 26 Olvera-Carrillo, Y. *et al.* A conserved core of programmed cell death indicator genes  
488 discriminates developmentally and environmentally induced programmed cell death in  
489 plants. *Plant Physiol.* **169**, 2684-2699 (2015).
- 490 27 Bollhöner, B. *et al.* Post mortem function of AtMC9 in xylem vessel elements. *New*  
491 *Phytol.* **200**, 498-510 (2013).
- 492 28 Durand, C. *et al.* The organization pattern of root border-like cells of *Arabidopsis* is  
493 dependent on cell wall homogalacturonan. *Plant Physiol.* **150**, 1411-1421 (2009).
- 494 29 Zhou, W. *et al.* *Arabidopsis* tyrosylprotein sulfotransferase acts in the  
495 auxin/PLETHORA pathway in regulating postembryonic maintenance of the root stem  
496 cell niche. *Plant Cell* **22**, 3692-3709 (2010).
- 497 30 Bennett, T., den Toorn, A., Willemsen, V. & Scheres, B. Precise control of plant stem  
498 cell activity through parallel regulatory inputs. *Develop.* **141**, 4055-4064 (2014).
- 499 31 Stø, I. M. *et al.* Conservation of the abscission signaling peptide IDA during  
500 Angiosperm evolution: withstanding genome duplications and gain and loss of the  
501 receptors HAE/HSL2. *Front. Plant Sci.* **6** (2015).

- 502 32 Somssich, M., Bleckmann, A. & Simon, R. Shared and distinct functions of the  
503 pseudokinase CORYNE (CRN) in shoot and root stem cell maintenance of  
504 *Arabidopsis*. *J. Exp. Bot.* **67**, 4901-4915 (2016).
- 505 33 Fauser, F., Schiml, S. & Puchta, H. Both CRISPR/Cas-based nucleases and nickases  
506 can be used efficiently for genome engineering in *Arabidopsis thaliana*. *Plant J.* **79**,  
507 348-359, (2014).
- 508 34 Yamaguchi, Y. L. *et al.* A Collection of Mutants for CLE-Peptide-Encoding Genes in  
509 *Arabidopsis* Generated by CRISPR/Cas9-Mediated Gene Targeting. *Plant Cell*  
510 *Physiol.* **58**, 1848-1856 (2017).
- 511
- 512
- 513
- 514
- 515
- 516

517 **Acknowledgements**

518 We want to thank Keiji Nakajima for pRCPG:nYG, pBRN1:BRN1-GFP and pBRN2:BRN2-  
519 GFP seeds and Malcolm Bennett for BFN1<sub>pro</sub>:nGFP; Roy Falleth, Solveig Engebretsen and  
520 Vegard Iversen for technical support in the lab and phytotron, Jonfinn Blix Knutsen for  
521 preparation of RNA for sequencing, and Mike Koomey for critical reading of the manuscript.  
522 This work has been supported by the Research Council of Norway (grant 312785) to the  
523 Aalen lab, and by the ERASysApp project “Rootbook” to Aalen and Claassen.

524

525 **Material requests**

526 Constructs or seeds, as well as the raw data underlying the RNAseq data in Supplementary  
527 Table 2 are available from R.B.A. upon request, e-mail: reidunn.aalen@ibv.uio.no.

528

529 **Contributions**

530 C.-L.S., T.I., S.S., U.H., M.A.B, M.W., M.K.A. and V.O. generated *Arabidopsis* lines; J.F.  
531 and D.v.W. designed, D.v.W. and I.K. performed, and D.v.W., I.K. and R.B.A. analysed live  
532 imaging experiments; G.F., M.A., M.A.B designed and M.W. performed IDL1-HSL2  
533 interaction studies; M.C. and R.B.A. designed and A.K., M.C. and R.B.A. analysed RNAseq  
534 data; the rest of the experiments were designed by C.-L.S. and R.B.A., performed by C.-L.S.  
535 together with U.H., M.W., V.O. and analyzed by R.B.A., C.-L.S., M.W. and U.H.; C.-L.S.  
536 drafted the manuscript; R.B.A. wrote the paper with input from other authors.

537 **Competing interests**

538 The authors declare no competing interests.

539 12

540 **Figure legends:**

541 **Fig. 1 | IDL1 peptide interacts with and activates HSL2.** **a**, The tissues of the *Arabidopsis*  
542 root tip. LRC – lateral root cap. **b**, pIDL1:GUS after overnight incubation with X-gluc. **c**,  
543 Confocal medial optical section of 7 DAG PI-stained root tip expressing pHSL2::Venus-H2B.  
544 The QC is indicated by an ellipse. **d**, Alignment of mIDA and mIDL1 peptides. \* -  
545 hydroxylated proline. **e**, mIDL1 activates HSL2 transiently expressed in *N. benthamiana* in a  
546 dose-dependent manner. Activation resulted in oxidative burst which was monitored as  
547 relative light units (RLU) using a luminol-based assay<sup>8</sup>. EC50 - half-maximal efficient  
548 concentration. **f**, Unlabelled mIDL1 compete efficiently for binding of HSL2. The mIDA  
549 peptide with an N-terminal Valine residue labelled with acridinium (acri-V-IDA) bound to *N.*  
550 *benthamiana*-expressed HSL2, was treated with increasing concentrations of unlabelled  
551 mIDL1 or V-IDA. 100% binding – assay without competitors; negative control (background  
552 signal) – leaf tissue not expressing HSL2 incubated with 10 nM acri-V-IDA peptide and  
553 increasing concentrations of unlabelled V-IDA. IC50 - half-maximal inhibitory concentration.

554

555 **Fig. 2 | Live imaging details the sloughing process in Col-0, the effect of enhanced**  
556 **expression of IDL1 and dependency on HSL2.** **a**, The two-component constructs for  
557 inducible enhanced expression of IDL1 and GUS. The IDL1 promoter controls expression of  
558 XVE, which in the presence of estradiol (ES) binds the OlexA-TATA box. **b**, ES-induced  
559 GUS expression in Col-0 and *hsl2* mutant background after 1 hour (h) incubation with X-Gluc.  
560 Controls were grown without ES. **c**, Representative stages of the standard ~ 18h sloughing  
561 process of Col-0, captured by live imaging, i.e. initial fracture in LRC cell files, first above,  
562 after several h just below the QC (red and yellow arrows), and finally separation of a whole  
563 cap layer (white arrow). **d**, Overlapping sloughing events in EnhIDL1 root tips, i.e. a new



564 initiating gap (green arrow) occurring before completion of the ongoing event. **e**, Split root  
565 caps of *EnhIDL1hsl2*, i.e. gaps at the LRC-COL border (blue arrow) followed by detachment  
566 of a split cap (white arrows) still attached to the LRC covering the meristematic zone. **f**,  
567 Number of initiations and detachments per root per 24h. Bars indicate standard deviation. \* -  
568 significantly different from Col-0 (Student's t-test  $P < 0.05$ ). Number of roots - Col-0 9,  
569 *EnhIDL1* and *EnhIDL1hsl2*<sup>10</sup>. **g**, Average time from initiation to detachment and from  
570 detachment to initiation of a new event. N – number of events per genotype. Standard  
571 deviations are given as bars. \* - significantly different from Col-0 (Student's t-test  $P < 0.04$ ). **h**,  
572 Total number of sloughed caps and their phenotypes (cf. c-e) documented by live imaging  
573 over 64 to 70 hours.

574

575 **Fig. 3 | Higher frequency of sloughing is compensated with more stem cell divisions. a**,  
576 Number of detached root cap layers 6, 8 and 10 days after germination (DAG). \* -  
577 significantly different from other lines (grey), and without ES (blue) (Student's t-test  $P <$   
578  $0.005$ ). N=10-12 roots per genotype per treatment and per time point. Standard deviations are  
579 given as bars. **b**, Root tips after 10 days on medium with 10  $\mu$ M ES, and control without ES. **c**,  
580 Number of attached COL tiers at 8 and 10 DAG. **d**, Percentage of root caps with new 1st tier  
581 COL cells, and new LRC cell generated after division of the respective stem cells. \* -  
582 significantly different from Col-0 (Student's t-test  $P < 0.01$ ). **e**, Medial optical sections of 7  
583 DAG PI-stained stem cell niches of roots grown with ES, with schematic drawing below.  
584 Dots mark novel cell walls in *EnhIDL1*.

585

586 **Fig. 4 | *idl1* and *hsl2* mutants influence sloughing frequency and LRC division patterns.**  
587 **a**, Number of sloughed and attached layers. \* - significantly different from Col-0 (Student's t-

588 test  $P_{idl1} < 10^{-3}$ ,  $P_{hsl2} < 10^{-6}$ ,  $N = 10$ ). **b**, Stem cell niche of the indicated genotypes. Asterisks  
589 indicate LRC/Epidermis stem cells (pink), and LRC cells directly neighbouring the epidermal  
590 cell file (light and dark blue). **c**, Schematic presentation of cellular patterning shown in b.  
591 Arrows show the directions of cell divisions. **d**, Number of cells in the first and second LRC  
592 cell file neighbouring the epidermal cell file. \* - significantly different from Col-0 (Student's  
593 t-test  $P < 10^{-5}$ ).  $N_{Col-0}=22$ ,  $N_{hsl2}=17$  and  $N_{idl1}=18$ . Standard deviations are given as bars.

594

595 **Fig. 5 | IDL1-HSL2 regulates genes involved in cell wall degradation and programmed**  
596 **cell death.** Confocal images of PI stained 7 DAG root tips of the indicated genotypes. The  
597 white ellipses mark the QCs. **a**, Medial optical sections of roots expressing RCPG, lower  
598 panel only the channel showing the PI stain. Note many PI-stained nuclei in Col-0 (cf. also  
599 Supplementary Fig. 6). **b**, 3D reconstructions from z-stacks of root tips expressing the PCD  
600 marker  $BFN1_{pro}:nGFP$ . Note reduced expression in the *idl1* and *hsl2* mutants, especially at  
601 Site II (see also Supplementary Fig. 6). **c**, Medial optical section of PI stained 7 DAG root  
602 expressing  $pHSL2::Venus-H2B$ , with enlargement (blue rectangle) of LRC cell file in the MZ.  
603 Arrow heads point to PI stained nuclei.

604

605 **Fig. 6 | IDL1-HSL2 signalling affects the dynamics of root cap sloughing.** IDL1-activated  
606 HSL2 regulates the dynamics of sloughing events, through regulation of PCD and cell wall  
607 remodelling genes, and triggering of COL and LRC stem cell divisions.

608

609

610 **Table 1. Tissue specificity of HSL2 root tip regulated genes.**

611

| Tissue                        | Marker <sup>1</sup> | # genes <sup>2</sup><br>with highest<br>expression <sup>3</sup> | P-values | FDR<br>adjusted <sup>612</sup><br>p-values |
|-------------------------------|---------------------|-----------------------------------------------------------------|----------|--------------------------------------------|
| LRC                           | LRC                 | 20                                                              | 9.0 E-04 | 3.1E-02                                    |
| COL                           | pet111              | 29                                                              | 1.16E-06 | 8.14E-06                                   |
| QC                            | AGL42               | 8                                                               | 0.908    | 0.999 <sup>614</sup>                       |
| Epidermis<br>(hair cells)     | COBL9               | 12                                                              | 0.494    | 0.999                                      |
| Epidermis<br>(non-hair cells) | gl2                 | 1                                                               | 0.999    | 0.999                                      |
| Cortex                        | CORTEX              | 10                                                              | 0.737    | 0.999 <sup>616</sup>                       |
| Endodermis                    | scr5                | 1                                                               | 0.999    | 0.999                                      |

617

618 <sup>1</sup>Makers used for sorting cells of a particular tissue prior to RNA isolation (Nawy et al. 2005)

619 <sup>2</sup>Genes down regulated in the *hsl2* mutant relative to Col-0, cf. Supplementary Table 2.

620 <sup>3</sup>Based on expression data in radial tissue types (Nawy et al. 2005). Genes were assigned  
621 to the tissue with the highest expression level across all tissues.

622