

REVIEW ARTICLE

Mechanisms of radial glia progenitor cell lineage progression

Robert Beattie and Simon Hippenmeyer

Institute of Science and Technology Austria, Klosterneuburg, Austria

Correspondence

S. Hippenmeyer, Institute of Science and Technology Austria, Am Campus 1, 3400 Klosterneuburg, Austria
Fax: +43 (0) 2243 9000 2000
Tel: +43 (0) 2243 9000 5101
E-mail: simon.hippenmeyer@ist.ac.at

(Received 10 October 2017, revised 31 October 2017, accepted 6 November 2017, available online 22 November 2017)

doi:10.1002/1873-3468.12906

Edited by Wilhelm Just

The mammalian cerebral cortex is responsible for higher cognitive functions such as perception, consciousness, and acquiring and processing information. The neocortex is organized into six distinct laminae, each composed of a rich diversity of cell types which assemble into highly complex cortical circuits. Radial glia progenitors (RGPs) are responsible for producing all neocortical neurons and certain glia lineages. Here, we discuss recent discoveries emerging from clonal lineage analysis at the single RGP cell level that provide us with an inaugural quantitative framework of RGP lineage progression. We further discuss the importance of the relative contribution of intrinsic gene functions and non-cell-autonomous or community effects in regulating RGP proliferation behavior and lineage progression.

Keywords: cerebral cortex; clonal analysis; gliogenesis; mosaic analysis with double markers; neurodevelopment; neurogenesis; neuroscience; radial glia progenitor; single-cell labeling

What makes the human cortex unique and how did the regulatory cell proliferation programs in neural stem cells (NSCs) evolve to accommodate the generation of bigger and more complex brains during evolution? These are key questions that require a clear understanding of the cellular and molecular processes controlling the development of the cortical entity from a relatively simple neuroepithelium. Across most mammals the overall cortical architecture is remarkably well conserved, however, the relative size and neuropil density in human has expanded significantly [1–5]. The cortex is composed of six distinct layers with a diverse array of cell types including excitatory projection neurons, inhibitory interneurons, and astrocyte- and oligodendrocyte glial cells [5–7]. The cellular and molecular mechanisms of generating cell type diversity and regulating NSC lineage progression in the dorsal

telencephalon *in vivo* are mostly unknown. Key questions include: What is the quantitative and qualitative output of a single stem cell and how is the output/stem cell potential modulated over time? Which genetic and epigenetic factors regulate the temporal progression of a stem cell along its lineage? What is the relative contribution of cell-intrinsic vs. environmental and/or niche factors? Here, we focus on the above questions and discuss recent progress contributing to our conceptual understanding of cortical radial glia progenitor (RGP) cell lineage progression. For this review we mainly discuss advances that contribute to our quantitative understanding of the production of cortical projection neurons which are generated from dorsal telencephalic RGPs. We refer the reader to excellent recent reviews for the discussion of interneuron and glia genesis, and diversity [8–12].

Abbreviations

GW, gestation week; hESCs, human embryonic stem cells; IKNM, interkinetic nuclear migration; IP, intermediate progenitor; iPSCs, induced pluripotent stem cells; MADM, mosaic analysis with double markers; MST, mitotic somal translocation; NESC, neural epithelial stem cell; NSC, neural stem cell; oRG, outer radial glia cells; oSVZ, outer SVZ; RGP, radial glia progenitor; scRNA-seq, single-cell RNA sequencing; TAPs, transit amplifying progenitors; V-SVZ, ventricular-subventricular zone.

Radial glia progenitors generating cell type diversity in the neocortex

In the neocortex, projection neurons are derived from a common progenitor cell known as the RGP cell [13–16]. RGP themselves are derived from neural epithelial stem cells (NESCs) that compose the early embryonic neuroepithelium. NESCs were first identified by His over 100 years ago and are defined morphologically by a long basal process and an apical process that remains in contact with both the ventricle and the pial surface [17]. Sauer later confirmed that these cells undergo mitosis at the lumen surface, and that they contain apical–basal cell polarity. Furthermore, Sauer introduced the first model for interkinetic nuclear migration (IKNM) [18], where mitosis occurs near the apical side of the neural tube and the two daughter cells migrate away postdivision [19,20]. At approximately embryonic day (E) 9 in mice and around gestation week (GW) 5–6 in humans, NESCs begin to transition into RGPs [3,5,21]. Nascent RGPs initially undergo symmetric proliferative (aka amplification) divisions resulting in the expansion of the progenitor pool [5,22,23]. At around E12, RGPs transition into a neurogenic state and divide asymmetrically thereby producing cortical projection neurons [24,25]. The earliest born neurons (destined to become layer 6 projection neurons) split the preplate into the superficial marginal zone and the deeper subplate [26,27]. Through consecutive waves of neurogenesis, nascent neurons migrate radially along the RGP cell process into the most superficial layer of the developing cortex where they mature and differentiate. This process continues with each new wave of neurons migrating past the previous, resulting in the formation of distinct cortical laminae in an ‘inside-out’ fashion [28–37]. Early born, deep layer neurons (layers 5–6) are largely composed of corticofugal neurons that innervate brain regions beyond the neocortex including the thalamus, brainstem, and spinal cord [38,39]. Later born superficial neurons (layers 2–4) consist of intracortical neurons that project locally, ipsilaterally, or to the contralateral cortical hemisphere. The neurogenic expansion occurs in a waxing, surging, and waning output pattern of neurons, finishing at E17 in mice and approximately GW20 in humans [40–42]. While the laminar position allows a rough classification of projection neurons it also dictates the ultimate connectivity of cortical projection neurons. Based on physiological connectivity patterns, the concept of a canonical microcircuit has been established [43,44]. In recent years many other criteria have been employed to enable the classification of cortical cell types ranging from morphological, physiological to transcriptomic

fingerprints and myelination patterns [45]. In particular, single-cell RNA sequencing (scRNAseq) has ushered in a revolution in our understanding of the dynamic gene expression patterns and states; and their correlation with cellular fate and cortical cell type diversity [46,47]. Many of the technological advances related to the current state of the art scRNAseq methods, and how these advances have expanded our knowledge of cortical projection neuron heterogeneity have been recently reviewed [48–52]. Single-cell transcriptomes and methylomes [53] represent a robust measure to classify distinct cell types and predict lineage trees based on hierarchical clustering algorithms although the mechanistic principles responsible for their generation by RGPs *in vivo* remain to a large extent enigmatic.

Radial glia progenitors can also produce glia cells [54], including astrocytes and oligodendrocytes, which play important roles in the development, maintenance, and function of neuronal circuits [55,56]. Although gliogenesis has been shown to follow neurogenesis in the developing brain [57–63], the mechanisms of lineage progression from neurogenesis to gliogenesis, especially at the individual RGP cell level remain essentially unexplored [9,11,54,64].

Shortly after birth, the embryonic neuroepithelium transforms into the postnatal NSC niche in the ventricular–subventricular zone (V-SVZ) within the lateral ventricle [54,65,66]. While discrete subpopulations of RGPs give rise to ependymal cells [67], other RGPs transform into V-SVZ type B1 cells [68,69]. Type B1 cells represent the principal stem cell progenitors in adult neurogenesis [70]. Type B1 cells generate type C cells which represent transit amplifying progenitors (TAPs). Type C TAPs significantly expand the lineage and produce neuroblasts destined to populate the olfactory bulb [54,65,66].

The developmental programs regulating the successive generation of postmitotic neurons and glia cells, followed by progressive generation of postnatal progenitor cells by telencephalic RGPs need to be precisely implemented and regulated. Impairments in RGP lineage progression lead to alterations in the cortical cytoarchitecture which is thought to represent the major underlying cause for several neurological disorders including microcephaly or megalencephaly; and more subtle neurodevelopmental diseases including schizophrenia, autism, and epilepsy [3,71–74].

RGP lineage diversity and cortex size

The relative increase in size and complexity of the mammalian cerebral cortex during evolution correlates

with the acquisition of more sophisticated traits. A particular intriguing example is the skilled hand use where the transition from power grip toward precision grip correlates with the physical addition of new cortical fields involved in proprioception [75]. The culmination of brain growth (both in size and complexity) eventually led to the emergence of higher and unique cognitive traits that are characteristic to the human brain. In order to begin to understand how the brain can progressively increase in size, it is essential to obtain the complete RGP lineage trees in different species across evolution. This is a daunting task and currently the RGP-derived lineage of cortical projection neurons is best characterized in mice. However, recent efforts to recapitulate some of the earlier stages of human brain development, including regional patterning *in vitro* using human embryonic stem cells (hESCs) [76] may promise new insight in the future; at least on the general cell biological level by inferring lineage tree branches from cultured cells. Although hESC cultures have great potential, recent studies have employed induced pluripotent stem cells (iPSCs) derived from humans, chimpanzees, and macaques and directed these cell lines to a dorsal telencephalic fate. While these systems do not fully recapitulate the *in vivo* cellular niche (including environmental factors that may have a direct role in regulating cellular fate) the cell-intrinsic potential and putative differences among species-specific iPSC lines is interesting. It has been observed that human iPSCs display more prolonged symmetric proliferative divisions while in macaque they transition to an asymmetric neurogenic division relatively quickly [77]. Once human progenitors switched to asymmetric neurogenic divisions, they continued producing neurons for a longer period of time than macaque. Therefore, it is evident that at the cellular level fundamental differences and species-specific progenitor traits may indicate distinct regulatory mechanisms which may contribute to the evolution of adapted progenitor proliferation potential and thus brain size.

Radial glia progenitors have been shown to not only give rise to neurons directly but also generate a diverse range of distinct transient progenitor cell types with varying degrees of potency. These include intermediate progenitors (IPs), short neural precursors, TAPs and outer SVZ progenitors (oSVZ) or also referred to as outer radial glia cells (oRGs), and have been described [78–81]. The cellular and molecular features of oRGs have been discussed at length in recent reviews [82,83], and here we will only focus on the oRG proliferation potential with regard to the increase in cortex size during evolution. Given the emerging diversity of

progenitor cells originating from RGPs it is important to analyze not only the total output of RGPs but also the proliferation potential of every class of IP. What types of neurons and/or glia cells are produced by each type of IP and how does the output change during overall lineage progression? In other words, what is the precise contribution of each progenitor and how do they contribute to quantity and cell type diversity during development and in different species? These are challenging questions for the field and in order to approach them concretely it is critical to observe and quantify neurogenesis at the single progenitor level. At the qualitative level, it will be revealing to characterize and classify the progenitors and their output by using morphological parameters paired with single-cell transcriptomes.

Elegant work from the Kriegstein lab and others has shown that human RGPs transit into distinct morphotypes during development and lineage progression [83–85]. For instance, after GW 17, ventrally located RGPs lose pia-contacting basal processes and transition into a ‘truncated’ RGP [85] which contribute to the expansion of the human cortical plate. oRGs themselves have been shown to have a huge proliferation and thus output potential since they are capable of producing several hundred neurons and thereby amplify the overall output of individual RGPs along their lineage [86,87]. Based on morphological analyses oRGs have been suggested to divide into several distinct classes which may individually exhibit additional distinctions in their output potential [84]. oRGs exhibit a unique dynamic mode of locomotion called mitotic somal translocation (MST) [79]. Immediately before cytokinesis, the soma rapidly translocates toward the cortical plate, which is independent of mitosis and working through an alternate mechanism to that of IKNM and saltatory nuclear migration [79]. While MST requires the activation of the Rho effector ROCK and nonmuscle myosin II the molecular mechanisms underlying MST are not well understood [88]. It is tempting to speculate that oRG-specific MST may be regulated precisely to optimize and/or tune their proliferation potential along their migration path. oRGs not only amplify the overall RGP output but they also constitute a critical scaffold for radially migrating neurons and their presence and proliferation properties have been implicated in gyrification [78,89–93].

Since oRGs are largely absent in the mouse brain, many studies have used variations of cerebral organoid systems which to varying degrees recapitulate some of the cortical structures seen in the human. In order to rigorously study the cellular properties of oRGs, a

number of experimental protocols [90,94–97], including the use of induced LIF/STAT3 signaling [98], have been continuously optimized. Although culture systems present certain caveats when it comes to the study of neurogenesis in human [99], these organoid systems still enable the study of human neural stem, RGP, and oRG cells in a short temporal window of human brain development [100]. Furthermore, they allow the introduction of perturbations into human-derived RGPs and follow the downstream repercussions. This has been particularly beneficial in addressing the role of Zika virus-induced cell death and proliferation deficits in RGPs which results in microcephaly in prenatal infants of infected mothers [98,101–103]. *In vitro* human iPSC model systems have also been surprisingly robust in recapitulating the temporal order of cortical neurogenesis in certain disease contexts including lissencephaly [95,104].

Outer radial glia cells are also found in other non-primates, including ferrets. However, the mechanisms governing their proliferation and differentiation properties seem to differ from the ones in primates [78,80]. In mouse there is a small number of oSVZ radial glia-like cells (also referred to as basal RG or bRG), however, these differ significantly from oRGs in humans [81,105]. Under normal physiological conditions, mouse bRGs are incapable of symmetric proliferative divisions and express *Tbr2* at midneurogenesis, a marker characteristically expressed by IPs [89]. While expression of specific genes including *TBC1D3* in ventricular cortical progenitors can promote the generation and expansion of bRGs in mice, the underlying mechanisms controlling the generation of these cells is still not entirely clear [106].

Quantitative analysis of RGP lineage progression at population level

In order to decipher the precise path of RGP lineage progression the qualitative and quantitative RGP output has to be assessed [23,51,107]. Traditionally, a key approach to study neuronal output of proliferating RGPs in the developing cortex was to analyze (by pulse-chase labeling analysis) the behavior of a population of cells and infer the patterns of division down to the single RGP level [108,109]. These analyses helped to shape our current model of RGP lineage progression in the neocortex which is based on the concept that multipotent RGPs first undergo symmetric proliferative (amplification) divisions followed by sequential asymmetric divisions generating neurons for distinct laminae in a defined temporal order [5]. Transplantation studies [110,111] showed that cortical RGPs

progressively restrict their potential. In other words, ‘late’ RGPs cannot revert back to produce ‘early’ lower layer neurons, whereas ‘early’ RGPs keep their full potential regardless of the age of the host. Indeed, the neurons which were produced by the ‘early’ progenitors were those expected based on the recipient’s stage [110,111]. Interestingly, cortical RGPs retain their ability to sequentially produce distinct cell types in culture conditions [61,112,113].

In order to directly measure the dynamics of RGP proliferation behavior during lineage progression, criteria such as cell cycle length at different developmental time periods and the proportion of cells that exit the cell cycle after each division represent two key measurements. Pioneering experiments from Nowakowski and colleagues predicted that the cardinal RGP undergoes 11 rounds of division beginning at E11 and ending at E17 which marks the end of neurogenesis in the mouse [109]. There are four variables which are commonly used when determining the total RGP output of the neocortex. First, the growth fraction or the proportion of RGP and IP cells that are actively proliferating [109]. Second, the number of RGP cell cycles during the cortical neurogenic time period. Third and fourth, the number of cells exiting the cell cycle and the number of cells remaining in a proliferative state after each round of division, respectively [114]. With these variables, Nowakowski and colleagues were able to very accurately model the growth of the cortex. For example, the absolute number of cells undergoing proliferation will change from one cell cycle to the next. When neurogenesis begins, the number of proliferating (P) RGP would be close to a maximum P value of 1, with 1 implying that every cell in the developing cortex is proliferating. With each cell cycle, a certain fraction of these daughter cells would exit the cell cycle and either become quiescent (Q) or terminally differentiate. The value of Q will gradually increase from 0 until it eventually reaches 1 at the end of neurogenesis. P and Q are inversely correlated and together can be used to calculate the absolute number of cells proliferating during each round of cell division [114]. Correlating these findings with the limited human data at the time was challenging. However, in chimpanzees it appears that similar to mouse the cell cycle length increases, and fewer proliferating cells can be observed, in later time windows [40]. These data led to two major conclusions. First, RGPs gradually become committed, stop proliferating, and eventually differentiate. Second, the key factor determining cortex size across evolution is not only reflected in the added diversity of progenitors such as oRGs (see above) but also in the time period of active RGP

proliferation and the length of the cell cycle at progressive later times [78–80,84,115]. While this model accurately accounts for the growth and absolute number of neocortical neurons, it does not take into consideration more subtle differences such as adjusting for the proportion of different progenitor cell types. It also does not allow for small changes in the ratio of RGP to IP or bRG and how this would affect overall output as illustrated also above in the previous section.

In order to move from population analyses to single-cell approaches, many studies used retroviral labeling of individual cells. The lineage of single proliferating RGPs were traced using mainly unicolor marking to then retroactively infer division patterns [116–121]. This method was also combined with live imaging to visualize individual cell divisions in real time, however, due to the technical limitations of the explant culture system, this was limited to shorter time periods of 24–48 h and did not allow for them to follow these cells into their final mature state [16,60]. Nowakowski and colleagues also combined various approaches and compared cell cycle behavior at the population level with the lineage analysis of the single retro-labeled RGP [20,22,40,109]. Their findings were remarkably robust and through mathematical modeling they were able to deduce that multiple populations of progenitor cells must be proliferating in parallel albeit not in synchrony. While *in vivo* lineage analysis provides a snapshot of clone composition at specific timepoints in development, it does not always provide complete information about the birth order of all clonally related cells. By culturing cortical RGPs *in vitro*, the temporal order of neurogenesis could be tracked reliably [61,113,122]. The fate of all progeny was followed through long-term live imaging, allowing for the generation of intricate cell lineage tree models to represent the birth order of each cell [113]. From these data, mathematical modeling was used to generate theoretical predictive models of cortical neurogenesis and identify division patterns of progenitor cells.

***In vivo* model of RGP lineage progression at single-cell level**

While population-based approaches provided a robust frame work of RGP progenitor proliferation patterns and properties, high-resolution single-cell approaches are necessary to establish a definite model of RGP lineage progression *in vivo*. In order to pursue high-resolution single-cell lineage tracing, progenitor stem cells should be marked in a sparse but permanent manner. In the most optimal case, the marker will be

transferred to the whole lineal progeny and even after numerous rounds of cell division still robustly mark all daughter cells. To this end, a large number of methods and approaches have been developed in the last decades which afford lineage tracing and clonal studies [51,123–125]. In particular, a variety of combinatorial fluorescent systems have been developed for performing high-resolution *in vivo* lineage analysis. Very prominent is the ‘Brainbow’ approach which is widely used in a variety of systems and which has recently been used to trace cortical RGP clones [126,127]. Another system, ‘CLoNe’, utilizes transposition vectors for cortical neuron lineage tracing [128,129] and the astrocyte-specific ‘Star Track’ specifically labels astrocyte progenitors and progeny [130]. The advantage of all the above systems is that the reporter constructs stably integrate into the genome, and thereby reliably label lineally related daughter cells. Combinatorial labeling with multiple markers also allows for the distinction between closely localized or spatially overlapping clones [127,128].

The monitoring, however, of precise progenitor division patterns *in situ*, and mapping entire lineage trees originating from an individual progenitor still represents a substantial challenge in the field. To this end, we have recently advanced the mosaic analysis with double markers (MADM) technology which provides an unprecedented genetic approach for *in vivo* lineage tracing in the mouse [131–133]. For MADM, two reciprocally chimeric marker genes are targeted separately to identical loci on homologous chromosomes. The chimeric marker genes (*GT* and *TG* alleles) consist of partial coding sequences for green (eGFP[G]) and red (tdT[T], tandem dimer Tomato) fluorescent proteins separated by an intron containing the loxP site. Following Cre recombinase-mediated interchromosomal recombination during mitosis, functional green and red fluorescent proteins are reconstituted resulting in two daughter cells each expressing one of the two fluorescent proteins (upon G2-X events: recombination in G2 of the cell cycle followed by X segregation, for technical details refer to Refs [131–133]). Analysis of MADM-based G2-X events in conjunction with temporally controlled tamoxifen (TM)-inducible CreER can provide exact information on birth dates of RGP (and other stem cell) clones and their cell division patterns (i.e., symmetric vs. asymmetric) [23,132,134]. An added MADM feature is the possible introduction of gene mutations allowing clonal two-color labeling with concomitant genetic manipulation. As such, these MADM applications permit the tracing of stem cell lineage progression in genetic mosaics with wild-type daughter cells labeled with one color (e.g., red) and

homozygous mutant siblings with the other (e.g., green) in an unlabeled heterozygous environment. In summary, MADM can provide an unambiguous quantitative optical readout of the proliferation mode (symmetric vs. asymmetric) of progenitors at the single-cell level and thus permit the determination of the developmental progenitor potential *in situ*.

In order to gain insight into the precise patterns of RGP division patterns and proliferation behavior, during neuron and glia production we have recently performed MADM-based quantitative clonal analysis [135]. Our systematic clonal analysis suggests that the behavior of individual RGPs is remarkably coherent and predictable across all developmental stages. RGPs initially undergo symmetric division with a predictable proliferation potential before transitioning to asymmetric neurogenic division. Importantly, the explicit identification of asymmetric neurogenic MADM clones enabled a quantitative assessment of the neurogenic potential of individual RGPs as they switch from symmetric proliferative division to asymmetric neurogenic division. We found that RGPs in the neurogenic phase do not undergo terminal differentiation in a stochastic manner but rather follow a defined nonrandom program of cell cycle exit resulting in a unitary output of about eight to nine neurons per individual RGP. Perhaps interestingly, the size of asymmetric neurogenic

clones was similar across neocortical areas with distinct functions, suggesting that the unitary neuronal output is a general property of cortical RGPs. Upon completion of neurogenesis, a defined fraction of individual RGPs proceed to gliogenesis, whereby about one in six neurogenic RGPs proceed to produce glia—astrocytes and/or oligodendrocytes—indicating a coupling between gliogenesis and neurogenesis at a predictable rate. Altogether, these MADM-based clonal analyses revealed definitive ontogeny of neocortical excitatory neurons and glia [135] (Fig. 1). While the MADM analysis detailed above provides a quantitative framework of lineage progression at the individual RGP cell level, the quality of distinct clones with unitary output remains to be determined. In other words, while the canonical RGP output is approximately eight to nine neurons, their distribution in the cortical plate may be fixed or display heterogeneity to various degrees. Furthermore, the clonal distribution pattern could differ in distinct functional areas. In order to address these questions it will be important in the future to monitor potential clonal heterogeneity at single-cell resolution and correlate the neuronal distribution with the functional areas in the neocortex. While we focus here mainly on neuronal output from RGPs, the predictable rate of glia production based on MADM analysis suggests a specific inherent gliogenic




Fig. 1. Deterministic RGP behavior and unitary production of projection neurons in the neocortex. Systematic clonal analysis suggests that the behavior of RGPs is coherent and predictable across all developmental stages. RGPs initially undergo symmetric division with a predictable proliferation potential before transitioning to asymmetric neurogenic divisions. RGPs in the neurogenic phase do not undergo terminal differentiation in a stochastic manner but rather follow a defined nonrandom program of cell cycle exit resulting in a unitary output of about eight to nine neurons per individual RGP. Roman numerals VI to II refer to the serial production of neurons destined to cortical layers which are numbered accordingly. Upon completion of neurogenesis, a defined fraction of individual RGPs proceed to gliogenesis whereby about one in six neurogenic RGPs proceed to produce glia—astrocytes and/or oligodendrocytes—indicating a coupling between gliogenesis and neurogenesis at a predictable rate (Adapted from Ref. [135]).

potential which requires further analysis. Future studies should also integrate the quantitative concept of unitary neuron production at the cortical neuronal circuit level and evaluate the functional implications with respect to the canonical cortical wiring diagram.

Molecular and cellular mechanisms of RGP lineage progression

The MADM-based clonal analysis provides an inaugural quantitative framework of RGP lineage progression (Fig. 1), but the cellular and molecular mechanisms are not well understood. Key questions include: How is the switch from symmetric proliferative RGP divisions to asymmetric neurogenic RGP divisions controlled? Which mechanisms determine the neurogenic and gliogenic RGP potential? How is the deterministic mode of cortical neuron production regulated? In order to address the above questions the MADM system provides a platform for directed candidate gene approaches [132,136–141]. To this end we recently commenced to functionally analyze the molecular requirements controlling the first critical step in RGP lineage progression: the switch from symmetric (expanding) to asymmetric (neurogenic) RGP progenitor division. One key regulator of the mode of cell division is the signaling protein LGL1 [aka Lgl1, lethal giant larvae homolog 1 (*Drosophila*)], which regulates intracellular polarity in a variety of cellular contexts, and likely plays an important role in RGPs in mouse *in vivo* [142–146]. Albeit being predicted to contribute to embryonic RGP lineage progression, how *Lgl1* controls this process is not entirely clear. Furthermore, the relationship between LGL1-mediated cell polarity, ventricular zone architecture, and cortical RGP behavior has not been extensively studied *in vivo*. The functional requirement of *Lgl1* at later stages during NSC lineage progression, and including gliogenesis, is essentially unknown due to lethality of *Lgl1* knockout mice at birth. The analysis of RGP lineage progression in *Lgl1* mutant mice is somewhat compromised due to the severe and progressive disruption of the VZ resulting in disorganization and tumor-like growth of RGPs in the form of rosettes [143]. This, however, also raises the possibility that substantial aspects of the phenotype in whole tissue *Lgl1* knockout could be the result of a combination of cell-autonomous and non-autonomous and/or community effects. In order to address this issue, and to determine the relative contribution of cell-autonomous *Lgl1* signaling and non-cell-autonomous mechanisms in RGP lineage progression, we capitalized on the MADM system. We developed the following genetic strategy: subtractive

phenotypic RGP analysis in genetic *Lgl1* mosaics (heterozygous, normal background; *Lgl1*-MADM) and conditional *Lgl1* knockouts (mutant background; cKO-*Lgl1*-MADM). In other words, *Lgl1* mutant RGP cells are either surrounded by an environment with 'normal' heterozygous and wild-type cells (*Lgl1*-MADM), or by mutant cells (cKO-*Lgl1*-MADM) [147]. The above genetic strategy represents a unique experimental paradigm (Fig. 2) which can be applied in principle to any candidate gene of interest to determine the relative contribution of intrinsic gene function and the effect of non-cell-autonomous effect on the overall phenotype *in vivo*. Interestingly, in cKO-*Lgl1*-MADM (but not *Lgl1*-MADM) the formation of heterotopic masses or subcortical band heterotopias was a predominant phenotype correlating with the downregulation of basolateral adherens junctional components similar like in the full knockout of *Lgl1* [143]. The cKO-*Lgl1*-MADM appeared to phenocopy *Numb/Numbl* double mutants [148–150]. NUMB localizes to the basolateral cadherin–catenin adhesion complex and is thought to control the trafficking of components such as N-cadherin (CDH2) [150]. Intriguingly, the loss of CDH2 [151] or α E-catenin [152], both resulted in the formation of heterotopias. A recent study observed that nonphosphorylated LGL1 strongly bound CDH2, whereas LGL1 with amino acid substitutions that mimicked phosphorylation did not interact with CDH2 [153]. These data suggest that LGL1 plays a critical role in adherens junction formation by regulating junctional CDH2 integrity presumably by regulating its internalization and/or intracellular trafficking [153]. Since *Lgl1* has also been suggested to play a role in polarized secretion and exocytosis regulation [154] it will be interesting to determine any putative functional relationship of *Lgl1* and *Numb* in regulating adherens junctional integrity and/or in controlling RGP proliferation dynamics. In mosaic *Lgl1*-MADM mice, where *Lgl1* was deleted only sparsely and/or removed from single RGPs, the mutant progenitors did proliferate normally with a unitary neuron output. These data indicate that wild-type progenitors surrounding mutant ones, maintain their integrity in the VZ in a non-cell-autonomous manner and that the exuberant RGP proliferation (and thus disturbed RGP lineage progression) in cKO-*Lgl1*-MADM and heterotopia formation is the result of community effects rather than the consequence of cell-autonomous *Lgl1* deficit.

One striking observation in individual *Lgl1*-MADM clones was that loss of *Lgl1* did not change the unitary neuron output but led to a massive increase in clonally related parenchymal astrocytes. However, it is currently not clear whether the increased proliferation of




Fig. 2. Mosaic analysis with double markers-based genetic dissection of cell-autonomous gene function and non-cell-autonomous effects regulating RGP lineage progression. The genetic assay relies on comparative analysis of multiple MADM paradigms to distinguish cell-autonomous genetic functions from non-cell-autonomous effects. MADM-based genetic dissection of a gene of interest (*Gene X*) requires mutant alleles to be introduced distal to the MADM cassettes via meiotic recombination (for details how to introduce mutant alleles into the MADM system see also Ref. [132]). (A–C) Schematic illustration of experimental paradigm in control-MADM (A, wild-type), *Gene X*-MADM (B, genetic mosaic), and cKO-*Gene X*-MADM (C, conditional/full knockout). In control-MADM, GFP⁺ (green), tdT⁺ (red), and unlabeled (vast majority) cells are all WT. In *Gene X*-MADM, GFP⁺ (green) cells are *Gene X*^{-/-}, tdT⁺ (red) cells are *Gene X*^{+/+}, and unlabeled cells are *Gene X*^{+/-}. In cKO-*Gene X*-MADM, GFP⁺ (green), tdT⁺ (red), and the vast majority of unlabeled cortical projection neurons are all *Gene X*^{-/-}. By phenotypically comparing the GFP⁺ *Gene X*^{-/-} cells in *Gene X*-MADM (B, D) to the genotypically identical GFP⁺ *Gene X*^{-/-} cells in cKO-*Gene X*-MADM (C, E) the cell-autonomous gene functions and relative contribution of non-cell-autonomous effects can be identified and quantified at single-cell resolution (Adapted from Ref. [147]).

astrocyte intermediate progenitor cells [54] solely is responsible for the increased number of *Lgll*^{-/-} cortical astrocytes. The loss of *Lgll* could in addition also lead to changes of the inherent gliogenic potential in RGPs. It will thus be informative to determine whether the fraction of gliogenic RGPs (1/6 in wild-type) is increased in *Lgll*-MADM. Furthermore, the increased astrocyte production in RGPs lacking *Lgll* was dependent on *Egfr* suggesting a functional relationship. In this regard, it is tempting to speculate that astrocyte overproduction could reflect the loss of a specific *Lgll*-dependent function in polarized secretion and/or exocytosis, in order to regulate cell surface abundance of astrocyte production-stimulating (such as for instance EGFR) and/or -inhibiting factors. It is intriguing to speculate that the control of polarized secretion, exocytosis [154,155], and possibly further intracellular trafficking events, could actually represent one unifying function of *Lgll* in the control of proliferating RGPs during several sequential stages in their

lineage progression (Fig. 3). Indeed, the loss of *Lgll* also compromises postnatal neurogenesis in a cell-autonomous manner although the underlying mechanisms remain to be clarified [147].

Outlook and perspectives

A rough framework of RGP lineage progression has been defined based on clonal analyses but a number of outstanding questions need to be addressed in the future. First, what is the degree of progenitor cell type diversity in the developing neuroepithelium and later in the VZ and SVZ? Do all RGPs harbor the same quantitative and qualitative potential for neuron/glia output? Recent lineage tracing and fate-mapping experiments employing distinct Cre/CreER-based approaches in combination with defined promoter elements suggest a significant level of progenitor cell type diversity [156–160]. While many promoters driving Cre/CreER recombinases lead to clones spanning all


Lgl1* function required for VZ integrity (control of embryonic neurogenesis)**Lgl1* regulates cortical astrocyte production*****Lgl1* controls postnatal neurogenesis in V-SVZ**

Fig. 3. Discrete sequential functions of *Lgl1* in regulating RGP behavior in the developing neocortex. Schematic model of RGP lineage progression and *Lgl1* functions at distinct stages of cortex development. See text for details (Adapted from Ref. [147]).

cortical layers, others appear to mark clones with more restricted laminar distribution. In most of the above studies, however, the analysis was not carried out at the single progenitor cell level. It thus remains an open question how diverse the RGP population really is with regard to the neurogenic and gliogenic inherent potential. The intrinsic RGP output potential could also be adjusted along the path of lineage progression and distinct RGP populations could respond differently to such regulation. It is interesting to note in this regard that once RGPs switch from symmetric to asymmetric neurogenic division about five of six RGPs lose the capacity to produce glia [135]. The advancement of scRNA-seq technologies holds great promise that in the near future the transcriptional profiles of large fractions of RGPs can be mapped in more detail, and the level of RGP cell diversity determined, at least at the level of gene expression. Building upon such data it will be important to correlate the gene expression profile with neuro- and gliogenic potential at the individual RGP level to evaluate the full spectrum of RGP cell type diversity.

What are the cellular and molecular mechanisms in cortical RGPs regulating the fine balance between proliferation and differentiation into neurons and/or glia cells, to specify the cerebral cortex of the correct size and cellular composition? While previous efforts

greatly contributed to our current framework of neocortical genesis, experimental paradigms addressing the function of specific genes were mostly based upon whole population approaches (e.g., full and/or conditional knockout studies). However, the lack of true single-cell resolution of progeny fate vital for dissecting progenitor division patterns has previously often precluded a definitive understanding. MADM offers a promising solution and permits quantitative clonal analysis, concurrent with genetic manipulation, of precise division patterns and lineage progression at unprecedented individual progenitor cell resolution. With MADM it is also possible to define and quantify the relative contributions of molecular genetic cell-autonomous and non-cell-autonomous mechanisms controlling lineage progression in RGPs at single-cell resolution [147]. Future MADM analyses hold the potential to systematically analyze lineage progression in any stem cell and tissue, and probe the relative contributions of the intrinsic and extrinsic components of any gene function to the overall phenotype.

While lineage analysis in higher order mammals remains technically challenging, recent studies have cleverly utilized naturally occurring endogenous retroelements, to create lineage maps in the human brain [51,161]. However, it is still unclear whether the deterministic and/or unitary mode of neuron production

in RGP represents a general principle. Thus, it will be important to establish models or RGP lineage progression at high single-cell resolution in other species than the mouse and especially in human. Recent studies have begun using cerebral organoids to approach the above issue [90,162–164] but complementary approaches beyond culture systems will be needed to obtain a more realistic model reflecting the *in vivo* condition. Ultimately, the combination of multidisciplinary approaches in cell culture and *in vivo*, and involving distinct species including human may promise a deeper understanding of the molecular mechanisms controlling (a) RGP lineage progression; (b) regulation of brain size in general; and (c) why human brain development is so sensitive to disruption of particular signaling pathways in pathological neurodevelopmental microcephaly, megalencephaly, or psychiatric disorders. In a broader context, the anticipated results will likely contribute to our knowledge of cortical neuron and/or glia specification and may potentially reveal a logic that can generate neuronal/glia diversity, thus providing a possible foundation for prospective future embryonic stem cell-based approaches in the context of directed brain repair [165–168].

Acknowledgements

We thank all members of the Hippenmeyer laboratory for discussion and comments on the manuscript. This work was supported by IST Austria Institutional Funds; a program grant from the Human Frontiers Science Program (RGP0053/2014) to SH; and the People Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme (FP7/2007–2013) under REA grant agreement no 618444 to SH.

References

- 1 Florio M, Borrell V and Huttner WB (2017) Human-specific genomic signatures of neocortical expansion. *Curr Opin Neurobiol* **42**, 33–44.
- 2 Lui JH, Hansen DV and Kriegstein AR (2011) Development and evolution of the human neocortex. *Cell* **146**, 18–36.
- 3 Silbereis JC, Pochareddy S, Zhu Y, Li M and Sestan N (2016) The cellular and molecular landscapes of the developing human central nervous system. *Neuron* **89**, 248–268.
- 4 Sousa AMM, Meyer KA, Santpere G, Gulden FO and Sestan N (2017) Evolution of the human nervous system function, structure, and development. *Cell* **170**, 226–247.
- 5 Taverna E, Götz M and Huttner WB (2014) The cell biology of neurogenesis: toward an understanding of the development and evolution of the neocortex. *Annu Rev Cell Dev Biol* **30**, 465–502.
- 6 Bartolini G, Ciceri G and Marín O (2013) Integration of GABAergic interneurons into cortical cell assemblies: lessons from embryos and adults. *Neuron* **79**, 849–864.
- 7 Lodato S and Arlotta P (2015) Generating neuronal diversity in the mammalian cerebral cortex. *Annu Rev Cell Dev Biol* **31**, 699–720.
- 8 Bandler RC, Mayer C and Fishell G (2017) Cortical interneuron specification: the juncture of genes, time and geometry. *Curr Opin Neurobiol* **42**, 17–24.
- 9 Bayraktar OA, Fuentealba LC, Alvarez-Buylla A and Rowitch DH (2015) Astrocyte development and heterogeneity. *Cold Spring Harb Perspect Biol* **7**, a020362.
- 10 Haim LB and Rowitch DH (2017) Functional diversity of astrocytes in neural circuit regulation. *Nat Rev Neurosci* **18**, 31–41.
- 11 Molofsky AV and Deneen B (2015) Astrocyte development: a guide for the perplexed. *Glia* **63**, 1320–1329.
- 12 Wamsley B and Fishell G (2017) Genetic and activity-dependent mechanisms underlying interneuron diversity. *Nat Rev Neurosci* **18**, 299–309.
- 13 Malatesta P, Hack MA, Hartfuss E, Kettenmann H, Klinkert W, Kirchhoff F and Gotz M (2003) Neuronal or glial progeny: regional differences in radial glia fate. *Neuron* **37**, 751–764.
- 14 Malatesta P, Hartfuss E and Gotz M (2000) Isolation of radial glial cells by fluorescent-activated cell sorting reveals a neuronal lineage. *Development* **127**, 5253–5263.
- 15 Miyata T, Kawaguchi A, Okano H and Ogawa M (2001) Asymmetric inheritance of radial glial fibers by cortical neurons. *Neuron* **31**, 727–741.
- 16 Noctor SC, Flint AC, Weissman TA, Dammerman RS and Kriegstein AR (2001) Neurons derived from radial glial cells establish radial units in neocortex. *Nature* **409**, 714–720.
- 17 His W (1889) Die Neuroblasten und Deren Entstehung im Embryonalen Mark. Vol 15. S. Hirzel, Leipzig.
- 18 Sauer FC (1935) Mitosis in the neural tube. *J Comp Neurol* **62**, 377–405.
- 19 Lee HO and Norden C (2013) Mechanisms controlling arrangements and movements of nuclei in pseudostratified epithelia. *Trends Cell Biol* **23**, 141–150.
- 20 Takahashi T, Nowakowski R and Caviness V (1993) Cell cycle parameters and patterns of nuclear movement in the neocortical proliferative zone of the fetal mouse. *J Neurosci* **13**, 820–833.

- 21 Howard BM, Mo Z, Filipovic R, Moore AR, Antic SD and Zecevic N (2008) Radial glia cells in the developing human brain. *Neuroscientist* **14**, 459–473.
- 22 Cai L, Hayes NL, Takahashi T, Caviness VS and Nowakowski RS (2002) Size distribution of retrovirally marked lineages matches prediction from population measurements of cell cycle behavior. *J Neurosci* **69**, 731–744.
- 23 Postiglione MP and Hippenmeyer S (2014) Monitoring neurogenesis in the cerebral cortex: an update. *Future Neurol* **9**, 323–340.
- 24 Alvarez-Buylla A, Garcia-Verdugo JM, Mateo AS and Merchant-Larios H (1998) Primary neural precursors and intermitotic nuclear migration in the ventricular zone of adult canaries. *J Neurosci* **18**, 1020–1037.
- 25 Garcia-Verdugo JM, Ferron S, Flames N, Collado L, Desfilis E and Font E (2002) The proliferative ventricular zone in adult vertebrates: a comparative study using reptiles, birds, and mammals. *Brain Res Bull* **57**, 765–775.
- 26 Hoerder-Suabedissen A and Molnar Z (2015) Development, evolution and pathology of neocortical subplate neurons. *Nat Rev Neurosci* **16**, 133–146.
- 27 Luskin MB and Shatz CJ (1985) Studies of the earliest generated cells of the cat's visual cortex: cogeneration of subplate and marginal zones. *J Neurosci* **5**, 1062.
- 28 Alvarez-Buylla A and Nottebohm F (1988) Migration of young neurons in adult avian brain. *Nature* **335**, 353–354.
- 29 Angevine JB Jr and Sidman RL (1961) Autoradiographic study of cell migration during histogenesis of cerebral cortex in the mouse. *Nature* **192**, 766–768.
- 30 Evsyukova I, Plestant C and Anton ES (2013) Integrative mechanisms of oriented neuronal migration in the developing brain. *Annu Rev Cell Dev Biol* **29**, 299–353.
- 31 Hansen AH, Duellberg C, Mieck C, Loose M and Hippenmeyer S (2017) Cell polarity in cerebral cortex development – cellular architecture shaped by biochemical networks. *Front Cell Neurosci* **11**, 176.
- 32 Hippenmeyer S (2014) Molecular pathways controlling the sequential steps of cortical projection neuron migration. *Adv Exp Med Biol* **800**, 1–24.
- 33 Komuro H and Rakic P (1998) Distinct modes of neuronal migration in different domains of developing cerebellar cortex. *J Neurosci* **18**, 1478–1490.
- 34 Marín O, Valiente M, Ge X and Tsai L-H (2010) Guiding neuronal cell migrations. *Cold Spring Harb Perspect Biol* **2**, a001834.
- 35 McConnell SK (1995) Constructing the cerebral cortex: neurogenesis and fate determination. *Neuron* **15**, 761–768.
- 36 Rakic P (1974) Neurons in rhesus monkey visual cortex: systematic relation between time of origin and eventual disposition. *Science* **183**, 425–427.
- 37 Rakic P (2009) Evolution of the neocortex: perspective from developmental biology. *Nat Rev Neurosci* **10**, 724–735.
- 38 Greig LFC, Woodworth MB, Galazo MJ, Padmanabhan H and Macklis JD (2013) Molecular logic of neocortical projection neuron specification, development and diversity. *Nat Rev Neurosci* **14**, 755–769.
- 39 Lodato S, Shetty AS and Arlotta P (2015) Cerebral cortex assembly: generating and reprogramming projection neuron diversity. *Trends Neurosci* **38**, 117–125.
- 40 Caviness V, Takahashi T and Nowakowski R (1995) Numbers, time and neocortical neuronogenesis: a general developmental and evolutionary model. *Trends Neurosci* **18**, 379–383.
- 41 deAzevedo LC, Fallet C, Moura-Neto V, Dumas-Duport C, Hedin-Pereira C and Lent R (2003) Cortical radial glial cells in human fetuses: depth-correlated transformation into astrocytes. *J Neurobiol* **55**, 288–298.
- 42 Zecevic N (2004) Specific characteristic of radial glia in the human fetal telencephalon. *Glia* **48**, 27–35.
- 43 Douglas RJ and Martin KA (1991) A functional microcircuit for cat visual cortex. *J Physiol* **440**, 735–769.
- 44 Steinmetz MA, Motter BC, Duffy CJ and Mountcastle VB (1987) Functional properties of parietal visual neurons: radial organization of directionalities within the visual field. *J Neurosci* **7**, 177–191.
- 45 Tomassy GS, Berger DR, Chen H-H, Kasthuri N, Hayworth KJ, Vercelli A, Seung HS, Lichtman JW and Arlotta P (2014) Distinct profiles of myelin distribution along single axons of pyramidal neurons in the neocortex. *Science* **344**, 319–324.
- 46 Tasic B, Menon V, Nguyen TN, Kim TK, Jarsky T, Yao Z, Levi B, Gray LT, Sorensen SA, Dolbeare T *et al.* (2016) Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. *Nat Neurosci* **19**, 335–346.
- 47 Zeisel A, Muñoz-Manchado AB, Codeluppi S, Lönnerberg P, La Manno G, Juréus A, Marques S, Munguba H, He L, Betsholtz C *et al.* (2015) Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-seq. *Science* **347**, 1138–1142.
- 48 Lein E, Borm LE and Linnarsson S (2017) The promise of spatial transcriptomics for neuroscience in the era of molecular cell typing. *Science* **358**, 64.
- 49 Lein ES, Belgard TG, Hawrylycz M and Molnár Z (2017) Transcriptomic perspectives on neocortical structure, development, evolution, and disease. *Annu Rev Neurosci* **40**, 629–652.

- 50 Poulin J-F, Tasic B, Hjerling-Leffler J, Trimarchi JM and Awatramani R (2016) Disentangling neural cell diversity using single-cell transcriptomics. *Nat Neurosci* **19**, 1131–1141.
- 51 Woodworth MB, Girsakis KM and Walsh CA (2017) Building a lineage from single cells: genetic techniques for cell lineage tracking. *Nat Rev Genet* **18**, 230–244.
- 52 Zeng H and Sanes JR (2017) Neuronal cell-type classification: challenges, opportunities and the path forward. *Nat Rev Neurosci* **18**, 530–546.
- 53 Luo C, Keown CL, Kurihara L, Zhou J, He Y, Li J, Castanon R, Lucero J, Nery JR, Sandoval JP *et al.* (2017) Single-cell methylomes identify neuronal subtypes and regulatory elements in mammalian cortex. *Science* **357**, 600.
- 54 Kriegstein A and Alvarez-Buylla A (2009) The glial nature of embryonic and adult neural stem cells. *Annu Rev Neurosci* **32**, 149–184.
- 55 Chung W-S, Allen NJ and Eroglu C (2015) Astrocytes control synapse formation, function, and elimination. *Cold Spring Harb Perspect Biol* **7**, a020370.
- 56 Freeman MR and Rowitch DH (2013) Evolving concepts of gliogenesis: a look way back and ahead to the next 25 years. *Neuron* **80**, 613–623.
- 57 Costa MR, Bucholz O, Schroeder T and Gotz M (2009) Late origin of glia-restricted progenitors in the developing mouse cerebral cortex. *Cereb Cortex* **19** (Suppl 1), i135–i143.
- 58 Ge W-P, Miyawaki A, Gage FH, Jan YN and Jan LY (2012) Local generation of glia is a major astrocyte source in postnatal cortex. *Nature* **484**, 376.
- 59 Magavi S, Friedmann D, Banks G, Stolfi A and Lois C (2012) Coincident generation of pyramidal neurons and protoplasmic astrocytes in neocortical columns. *J Neurosci* **32**, 4762–4772.
- 60 Noctor SC, Martinez-Cerdeno V, Ivic L and Kriegstein AR (2004) Cortical neurons arise in symmetric and asymmetric division zones and migrate through specific phases. *Nat Neurosci* **7**, 136–144.
- 61 Qian X, Shen Q, Goderie SK, He W, Capela A, Davis AA and Temple S (2000) Timing of CNS cell generation: a programmed sequence of neuron and glial cell production from isolated murine cortical stem cells. *Neuron* **28**, 69–80.
- 62 Schmechel DE and Rakic P (1979) A Golgi study of radial glial cells in developing monkey telencephalon: morphogenesis and transformation into astrocytes. *Anat Embryol (Berl)* **156**, 115–152.
- 63 Voigt T (1989) Development of glial cells in the cerebral wall of ferrets: direct tracing of their transformation from radial glia into astrocytes. *J Comp Neurol* **289**, 74–88.
- 64 Tabata H (2015) Diverse subtypes of astrocytes and their development during corticogenesis. *Front Neurosci* **9**, 114.
- 65 Bond AM, Ming G-L and Song H (2015) Adult mammalian neural stem cells and neurogenesis: five decades later. *Cell Stem Cell* **17**, 385–395.
- 66 Lim DA and Alvarez-Buylla A (2016) The adult ventricular–subventricular zone (V-SVZ) and olfactory bulb (OB) neurogenesis. *Cold Spring Harb Perspect Biol* **8**, a018820.
- 67 Spassky N, Merkle FT, Flames N, Tramontin AD, García-Verdugo JM and Alvarez-Buylla A (2005) Adult ependymal cells are postmitotic and are derived from radial glial cells during embryogenesis. *J Neurosci* **25**, 10–18.
- 68 Fuentealba LC, Rompani SB, Parraguez JI, Obernier K, Romero R, Cepko CL and Alvarez-Buylla A (2015) Embryonic origin of postnatal neural stem cells. *Cell* **161**, 1644–1655.
- 69 Merkle FT, Tramontin AD, García-Verdugo JM and Alvarez-Buylla A (2004) Radial glia give rise to adult neural stem cells in the subventricular zone. *Proc Natl Acad Sci USA* **101**, 17528–17532.
- 70 Doetsch F, Caille I, Lim DA, García-Verdugo JM and Alvarez-Buylla A (1999) Subventricular zone astrocytes are neural stem cells in the adult mammalian brain. *Cell* **97**, 703–716.
- 71 Barkovich AJ, Dobyns WB and Guerrini R (2015) Malformations of cortical development and epilepsy. *Cold Spring Harb Perspect Med* **5**, a022392.
- 72 Desikan RS and Barkovich AJ (2016) Malformations of cortical development. *Ann Neurol* **80**, 797–810.
- 73 Geschwind DH and Flint J (2015) Genetics and genomics of psychiatric disease. *Science* **349**, 1489–1494.
- 74 Hu WF, Chahrour MH and Walsh CA (2014) The diverse genetic landscape of neurodevelopmental disorders. *Annu Rev Genomics Hum Genet* **15**, 195–213.
- 75 Padberg J, Franca JG, Cooke DF, Soares JGM, Rosa MGP, Fiorani M, Gattass R and Krubitzer L (2007) Parallel evolution of cortical areas involved in skilled hand use. *J Neurosci* **27**, 10106–10115.
- 76 Yao Z, Mich JK, Ku S, Menon V, Krostag A-R, Martinez RA, Furchtgott L, Mulholland H, Bort S, Fuqua MA *et al.* (2017) A single-cell roadmap of lineage bifurcation in human ESC models of embryonic brain development. *Cell Stem Cell* **20**, 120–134.
- 77 Otani T, Marchetto MC, Gage FH, Simons BD and Livesey FJ (2016) 2D and 3D stem cell models of primate cortical development identify species-specific differences in progenitor behavior contributing to brain size. *Cell Stem Cell* **18**, 467–480.
- 78 Fietz SA, Kelava I, Vogt J, Wilsch-Brauninger M, Stenzel D, Fish JL, Corbeil D, Riehn A, Distler W, Nitsch R *et al.* (2010) OSVZ progenitors of human and ferret neocortex are epithelial-like and expand by integrin signaling. *Nat Neurosci* **13**, 690–699.

- 79 Hansen DV, Lui JH, Parker PR and Kriegstein AR (2010) Neurogenic radial glia in the outer subventricular zone of human neocortex. *Nature* **464**, 554–561.
- 80 Reillo I, de Juan Romero C, Garcia-Cabezas MA and Borrell V (2011) A role for intermediate radial glia in the tangential expansion of the mammalian cerebral cortex. *Cereb Cortex* **21**, 1674–1694.
- 81 Wang X, Tsai JW, LaMonica B and Kriegstein AR (2011) A new subtype of progenitor cell in the mouse embryonic neocortex. *Nat Neurosci* **14**, 555–561.
- 82 Florio M and Huttner WB (2014) Neural progenitors, neurogenesis and the evolution of the neocortex. *Development* **141**, 2182–2194.
- 83 Ostrem B, Di Lullo E and Kriegstein A (2017) oRGs and mitotic somal translocation – a role in development and disease. *Curr Opin Neurobiol* **42**, 61–67.
- 84 Betizeau M, Cortay V, Patti D, Pfister S, Gautier E, Bellemin-Menard A, Afanassieff M, Huissoud C, Douglas RJ, Kennedy H *et al.* (2013) Precursor diversity and complexity of lineage relationships in the outer subventricular zone of the primate. *Neuron* **80**, 442–457.
- 85 Nowakowski TJ, Pollen AA, Sandoval-Espinosa C and Kriegstein AR (2016) Transformation of the radial glia scaffold demarcates two stages of human cerebral cortex development. *Neuron* **91**, 1219–1227.
- 86 Lewitus E, Kelava I and Huttner WB (2013) Conical expansion of the outer subventricular zone and the role of neocortical folding in evolution and development. *Front Hum Neurosci* **7**, 424.
- 87 Pollen AA, Nowakowski TJ, Chen J, Retallack H, Sandoval-Espinosa C, Nicholas CR, Shuga J, Liu SJ, Oldham MC, Diaz A *et al.* (2015) Molecular identity of human outer radial glia during cortical development. *Cell* **163**, 55–67.
- 88 Ostrem BEL, Lui JH, Gertz CC and Kriegstein AR (2014) Control of outer radial glial stem cell mitosis in the human brain. *Cell Rep* **8**, 656–664.
- 89 Florio M, Albert M, Taverna E, Namba T, Brandl H, Lewitus E, Haffner C, Sykes A, Wong FK and Peters J (2015) Human-specific gene ARHGAP11B promotes basal progenitor amplification and neocortex expansion. *Science* **347**, 1465–1470.
- 90 Li Y, Muffat J, Omer A, Bosch I, Lancaster MA, Sur M, Gehrke L, Knoblich JA and Jaenisch R (2017) Induction of expansion and folding in human cerebral organoids. *Cell Stem Cell* **20**, 385–396.e383.
- 91 Nonaka-Kinoshita M, Reillo I, Artegiani B, Ángeles Martínez-Martínez M, Nelson M, Borrell V and Calegari F (2013) Regulation of cerebral cortex size and folding by expansion of basal progenitors. *EMBO J* **32**, 1817–1828.
- 92 Pilz G-A, Shitamukai A, Reillo I, Pacary E, Schwausch J, Stahl R, Ninkovic J, Snippert HJ, Clevers H, Godinho L *et al.* (2013) Amplification of progenitors in the mammalian telencephalon includes a new radial glial cell type. *Nat Commun* **4**, 2125.
- 93 Stahl R, Walcher T, De Juan Romero C, Pilz GA, Cappello S, Irmeler M, Sanz-Aquela JM, Beckers J, Blum R, Borrell V *et al.* (2013) Trnp1 regulates expansion and folding of the mammalian cerebral cortex by control of radial glial fate. *Cell* **153**, 535–549.
- 94 Birey F, Andersen J, Makinson CD, Islam S, Wei W, Huber N, Fan HC, Metzler KRC, Panagiotakos G, Thom N *et al.* (2017) Assembly of functionally integrated human forebrain spheroids. *Nature* **545**, 54–59.
- 95 Iefremova V, Manikakis G, Krefft O, Jabali A, Weynans K, Wilkens R, Marsoner F, Brändl B, Müller F-J, Koch P *et al.* (2017) An organoid-based model of cortical development identifies non-cell-autonomous defects in wnt signaling contributing to Miller-Dieker syndrome. *Cell Rep* **19**, 50–59.
- 96 Lancaster MA, Renner M, Martin C-A, Wenzel D, Bicknell LS, Hurles ME, Homfray T, Penninger JM, Jackson AP and Knoblich JA (2013) Cerebral organoids model human brain development and microcephaly. *Nature* **501**, 373–379.
- 97 Qian X, Nguyen HN, Song MM, Hadiono C, Ogden SC, Hammack C, Yao B, Hamersky GR, Jacob F, Zhong C *et al.* (2016) Brain-region-specific organoids using mini-bioreactors for modeling ZIKV exposure. *Cell* **165**, 1238–1254.
- 98 Watanabe M, Buth JE, Vishlaghi N, de la Torre-Ubieta L, Taxis J, Khakh BS, Coppola G, Pearson CA, Yamauchi K, Gong D *et al.* (2017) Self-organized cerebral organoids with human-specific features predict effective drugs to combat Zika virus infection. *Cell Rep* **21**, 517–532.
- 99 Suzuki IK and Vanderhaeghen P (2015) Is this a brain which I see before me? Modeling human neural development with pluripotent stem cells. *Development* **142**, 3138–3150.
- 100 Huch M, Knoblich JA, Lutolf MP and Martinez-Arias A (2017) The hope and the hype of organoid research. *Development* **144**, 938–941.
- 101 Gabriel E, Ramani A, Karow U, Gottardo M, Natarajan K, Gooi LM, Goranci-Buzhala G, Krut O, Peters F, Nikolic M *et al.* (2017) Recent Zika virus isolates induce premature differentiation of neural progenitors in human brain organoids. *Cell Stem Cell* **20**, 397–406.e395.
- 102 Onorati M, Li Z, Liu F, Sousa AMM, Nakagawa N, Li M, Dell’Anno MT, Gulden FO, Pochareddy S, Tebbenkamp ATN *et al.* (2016) Zika virus disrupts phospho-TBK1 localization and mitosis in human

- neuroepithelial stem cells and radial glia. *Cell Rep* **16**, 2576–2592.
- 103 Qian X, Nguyen HN, Jacob F, Song H and Ming G-L (2017) Using brain organoids to understand Zika virus-induced microcephaly. *Development* **144**, 952–957.
- 104 Bershteyn M, Nowakowski TJ, Pollen AA, Di Lullo E, Nene A, Wynshaw-Boris A and Kriegstein AR (2017) Human iPSC-derived cerebral organoids model cellular features of lissencephaly and reveal prolonged mitosis of outer radial glia. *Cell Stem Cell* **20**, 435–449.e434.
- 105 Shitamukai A, Konno D and Matsuzaki F (2011) Oblique radial glial divisions in the developing mouse neocortex induce self-renewing progenitors outside the germinal zone that resemble primate outer subventricular zone progenitors. *J Neurosci* **31**, 3683–3695.
- 106 Ju XC, Hou QQ, Sheng AL, Wu KY, Zhou Y, Jin Y, Wen T, Yang Z, Wang X and Luo ZG (2016) The hominoid-specific gene TBC1D3 promotes generation of basal neural progenitors and induces cortical folding in mice. *eLife* **5**, e18197.
- 107 Blanpain C and Simons BD (2013) Unravelling stem cell dynamics by lineage tracing. *Nat Rev Mol Cell Biol* **14**, 489.
- 108 Cai L, Hayes NL and Nowakowski RS (1997) Local homogeneity of cell cycle length in developing mouse cortex. *J Neurosci* **17**, 2079.
- 109 Takahashi T, Nowakowski RS and Caviness VS Jr (1995) The cell cycle of the pseudostratified ventricular epithelium of the embryonic murine cerebral wall. *J Neurosci* **15**, 6046–6057.
- 110 Desai AR and McConnell SK (2000) Progressive restriction in fate potential by neural progenitors during cerebral cortical development. *Development* **127**, 2863–2872.
- 111 Frantz GD and McConnell SK (1996) Restriction of late cerebral cortical progenitors to an upper-layer fate. *Neuron* **17**, 55–61.
- 112 Shen Q, Wang Y, Dimos JT, Fasano CA, Phoenix TN, Lemischka IR, Ivanova NB, Stifani S, Morrisey EE and Temple S (2006) The timing of cortical neurogenesis is encoded within lineages of individual progenitor cells. *Nat Neurosci* **9**, 743–751.
- 113 Slater JL, Landman KA, Hughes BD, Shen Q and Temple S (2009) Cell lineage tree models of neurogenesis. *J Theor Biol* **256**, 164–179.
- 114 Takahashi T, Nowakowski RS and Caviness VS (1996) The leaving or Q fraction of the murine cerebral proliferative epithelium: a general model of neocortical neuronogenesis. *J Neurosci* **16**, 6183–6196.
- 115 Rakic P (1988) Specification of cerebral cortical areas. *Science* **241**, 170–176.
- 116 Kornack DR and Rakic P (1995) Radial and horizontal deployment of clonally related cells in the primate neocortex: relationship to distinct mitotic lineages. *Neuron* **15**, 311–321.
- 117 Luskin MB, Pearlman AL and Sanes JR (1988) Cell lineage in the cerebral cortex of the mouse studied in vivo and in vitro with a recombinant retrovirus. *Neuron* **1**, 635–647.
- 118 McCarthy M, Turnbull DH, Walsh CA and Fishell G (2001) Telencephalic neural progenitors appear to be restricted to regional and glial fates before the onset of neurogenesis. *J Neurosci* **21**, 6772–6781.
- 119 Price J, Turner D and Cepko C (1987) Lineage analysis in the vertebrate nervous system by retrovirus-mediated gene transfer. *Proc Natl Acad Sci USA* **84**, 156–160.
- 120 Walsh C and Cepko CL (1988) Clonally related cortical cells show several migration patterns. *Science* **241**, 1342–1345.
- 121 Walsh C and Cepko CL (1992) Widespread dispersion of neuronal clones across functional regions of the cerebral cortex. *Science* **255**, 434–440.
- 122 Qian X, Goderie SK, Shen Q, Stern JH and Temple S (1998) Intrinsic programs of patterned cell lineages in isolated vertebrate CNS ventricular zone cells. *Development* **125**, 3143–3152.
- 123 Buckingham ME and Meilhac SM (2011) Tracing cells for tracking cell lineage and clonal behavior. *Dev Cell* **21**, 394–409.
- 124 Kretschmar K and Watt FM (2012) Lineage tracing. *Cell* **148**, 33–45.
- 125 Legue E and Joyner AL (2010) Chapter ten-genetic fate mapping using site-specific recombinases. *Methods Enzymol* **477**, 153–181.
- 126 Livet J, Weissman TA, Kang H, Draft RW, Lu J, Bennis RA, Sanes JR and Lichtman JW (2007) Transgenic strategies for combinatorial expression of fluorescent proteins in the nervous system. *Nature* **450**, 56–62.
- 127 Loulier K, Barry R, Mahou P, Le Franc Y, Supatto W, Matho KS, Ieng S, Fouquet S, Dupin E, Benosman R *et al.* (2014) Multiplex cell and lineage tracking with combinatorial labels. *Neuron* **81**, 505–520.
- 128 García-Moreno F, Vasistha NA, Begbie J and Molnár Z (2014) CLoNe is a new method to target single progenitors and study their progeny in mouse and chick. *Development* **141**, 1589–1598.
- 129 Vasistha NA, García-Moreno F, Arora S, Cheung AFP, Arnold SJ, Robertson EJ and Molnár Z (2015) Cortical and clonal contribution of Tbr2 expressing progenitors in the developing mouse brain. *Cereb Cortex* **25**, 3290–3302.
- 130 García-Marqués J and López-Mascaraque L (2013) Clonal identity determines astrocyte cortical heterogeneity. *Cereb Cortex* **23**, 1463–1472.
- 131 Hippenmeyer S (2013) Dissection of gene function at clonal level using mosaic analysis with double markers. *Front Biol* **8**, 557–568.

- 132 Hippenmeyer S, Youn YH, Moon HM, Miyamichi K, Zong H, Wynshaw-Boris A and Luo L (2010) Genetic mosaic dissection of *Lis1* and *Ndel1* in neuronal migration. *Neuron* **68**, 695–709.
- 133 Zong H, Espinosa JS, Su HH, Muzumdar MD and Luo L (2005) Mosaic analysis with double markers in mice. *Cell* **121**, 479–492.
- 134 Espinosa JS and Luo L (2008) Timing neurogenesis and differentiation: insights from quantitative clonal analyses of cerebellar granule cells. *J Neurosci* **28**, 2301–2312.
- 135 Gao P, Postiglione MP, Krieger TG, Hernandez L, Wang C, Han Z, Streicher C, Papusheva E, Insolera R, Chugh K *et al.* (2014) Deterministic progenitor behavior and unitary production of neurons in the neocortex. *Cell* **159**, 775–788.
- 136 Espinosa JS, Wheeler DG, Tsien RW and Luo L (2009) Uncoupling dendrite growth and patterning: single-cell knockout analysis of NMDA receptor 2B. *Neuron* **62**, 205–217.
- 137 Hippenmeyer S, Johnson RL and Luo L (2013) Mosaic analysis with double markers reveals cell-type-specific paternal growth dominance. *Cell Rep* **3**, 960–967.
- 138 Joo W, Hippenmeyer S and Luo L (2014) Dendrite morphogenesis depends on relative levels of NT-3/TrkC signaling. *Science* **346**, 626–629.
- 139 Liang H, Xiao G, Yin H, Hippenmeyer S, Horowitz JM and Ghashghaei HT (2013) Neural development is dependent on the function of specificity protein 2 in cell cycle progression. *Development* **140**, 552–561.
- 140 Liu C, Sage JC, Miller MR, Verhaak RG, Hippenmeyer S, Vogel H, Foreman O, Bronson RT, Nishiyama A, Luo L *et al.* (2011) Mosaic analysis with double markers reveals tumor cell of origin in glioma. *Cell* **146**, 209–221.
- 141 Riccio P, Cebrian C, Zong H, Hippenmeyer S and Costantini F (2016) Ret and Etv4 promote directed movements of progenitor cells during renal branching morphogenesis. *PLoS Biol* **14**, e1002382.
- 142 Betschinger J, Mechtler K and Knoblich JA (2003) The Par complex directs asymmetric cell division by phosphorylating the cytoskeletal protein Lgl. *Nature* **422**, 326.
- 143 Klezovitch O, Fernandez TE, Tapscott SJ and Vasioukhin V (2004) Loss of cell polarity causes severe brain dysplasia in Lgl1 knockout mice. *Genes Dev* **18**, 559–571.
- 144 Vasioukhin V (2006) Lethal giant puzzle of Lgl. *Dev Neurosci* **28**, 13–24.
- 145 Wirtz-Peitz F and Knoblich JA (2006) Lethal giant larvae take on a life of their own. *Trends Cell Biol* **16**, 234–241.
- 146 Yamanaka T, Horikoshi Y, Sugiyama Y, Ishiyama C, Suzuki A, Hirose T, Iwamatsu A, Shinohara A and Ohno S (2003) Mammalian Lgl forms a protein complex with PAR-6 and aPKC independently of PAR-3 to regulate epithelial cell polarity. *Curr Biol* **13**, 734–743.
- 147 Beattie R, Postiglione MP, Burnett LE, Laukoter S, Streicher C, Pauler FM, Xiao G, Klezovitch O, Vasioukhin V, Ghashghaei TH *et al.* (2017) Mosaic analysis with double markers reveals distinct sequential functions of Lgl1 in neural stem cells. *Neuron* **94**, 517–533.e513.
- 148 Li H-S, Wang D, Shen Q, Schonemann MD, Gorski JA, Jones KR, Temple S, Jan LY and Jan YN (2003) Inactivation of Numb and Numbl like in embryonic dorsal forebrain impairs neurogenesis and disrupts cortical morphogenesis. *Neuron* **40**, 1105–1118.
- 149 Petersen PH, Zou K, Hwang JK, Jan YN and Zhong W (2002) Progenitor cell maintenance requires numb and numbl like during mouse neurogenesis. *Nature* **419**, 929–934.
- 150 Rašin M-R, Gazula V-R, Breunig JJ, Kwan KY, Johnson MB, Liu-Chen S, Li H-S, Jan LY, Jan Y-N and Rakic P (2007) Numb and Numbl are required for maintenance of cadherin-based adhesion and polarity of neural progenitors. *Nat Neurosci* **10**, 819–827.
- 151 Gil-Sanz C, Landeira B, Ramos C, Costa MR and Müller U (2014) Proliferative defects and formation of a double cortex in mice lacking Mltt4 and Cdh2 in the dorsal telencephalon. *J Neurosci* **34**, 10475–10487.
- 152 Lien W-H, Klezovitch O, Fernandez TE, Delrow J and Vasioukhin V (2006) α E-catenin controls cerebral cortical size by regulating the hedgehog signaling pathway. *Science* **311**, 1609–1612.
- 153 Jossin Y, Lee M, Klezovitch O, Kon E, Cossard A, Lien W-H, Fernandez TE, Cooper JA and Vasioukhin V (2017) Lgl1 connects cell polarity with cell-cell adhesion in embryonic neural stem cells. *Dev Cell* **41**, 481–495.e485.
- 154 Musch A, Cohen D, Yeaman C, Nelson WJ, Rodriguez-Boulan E and Brennwald PJ (2002) Mammalian homolog of *Drosophila* tumor suppressor lethal (2) giant larvae interacts with basolateral exocytic machinery in Madin-Darby canine kidney cells. *Mol Biol Cell* **13**, 158–168.
- 155 Lehman K, Rossi G, Adamo JE and Brennwald P (1999) Yeast homologues of tomosyn and lethal giant larvae function in exocytosis and are associated with the plasma membrane SNARE, Sec9. *J Cell Biol* **146**, 125–140.
- 156 Eckler MJ, Nguyen TD, McKenna WL, Fastow BL, Guo C, Rubenstein JL and Chen B (2015) Cux2-positive radial glial cells generate diverse subtypes of neocortical projection neurons and macroglia. *Neuron* **86**, 1100–1108.
- 157 Franco SJ, Gil-Sanz C, Martinez-Garay I, Espinosa A, Harkins-Perry SR, Ramos C and Müller U (2012)

- Fate-restricted neural progenitors in the mammalian cerebral cortex. *Science* **337**, 746.
- 158 Gil-Sanz C, Espinosa A, Fregoso SP, Bluske KK, Cunningham CL, Martinez-Garay I, Zeng H, Franco SJ and Muller U (2015) Lineage tracing using Cux2-Cre and Cux2-CreERT2 mice. *Neuron* **86**, 1091–1099.
- 159 Guo C, Eckler MJ, McKenna WL, McKinsey GL, Rubenstein JLR and Chen B (2013) Fezf2 expression identifies a multipotent progenitor for neocortical projection neurons, astrocytes, and oligodendrocytes. *Neuron* **80**, 1167–1174.
- 160 Kaplan ES, Ramos-Laguna KA, Mihalas AB, Daza RAM and Hevner RF (2017) Neocortical Sox9+ radial glia generate glutamatergic neurons for all layers, but lack discernible evidence of early laminar fate restriction. *Neural Dev* **12**, 14.
- 161 Evrony GD, Lee E, Mehta BK, Benjamini Y, Johnson RM, Cai X, Yang L, Haseley P, Lehmann HS, Park PJ *et al.* (2015) Cell lineage analysis in human brain using endogenous retroelements. *Neuron* **85**, 49–59.
- 162 He Z, Han D, Efimova O, Guijarro P, Yu Q, Oleksiak A, Jiang S, Anokhin K, Velichkovsky B, Grunewald S *et al.* (2017) Comprehensive transcriptome analysis of neocortical layers in humans, chimpanzees and macaques. *Nat Neurosci* **20**, 886–895.
- 163 Mora-Bermúdez F, Badsha F, Kanton S, Camp JG, Vernot B, Köhler K, Voigt B, Okita K, Maricic T, He Z *et al.* (2016) Differences and similarities between human and chimpanzee neural progenitors during cerebral cortex development. *eLife* **5**, e18683.
- 164 Quadrato G, Nguyen T, Macosko EZ, Sherwood JL, Min Yang S, Berger DR, Maria N, Scholvin J, Goldman M, Kinney JP *et al.* (2017) Cell diversity and network dynamics in photosensitive human brain organoids. *Nature* **545**, 48–53.
- 165 Espuny-Camacho I, Michelsen KA, Gall D, Linaro D, Hasche A, Bonnefont J, Bali C, Orduz D, Bilheu A, Herpoel A *et al.* (2013) Pyramidal neurons derived from human pluripotent stem cells integrate efficiently into mouse brain circuits in vivo. *Neuron* **77**, 440–456.
- 166 Falkner S, Grade S, Dimou L, Conzelmann K-K, Bonhoeffer T, Götz M and Hübener M (2016) Transplanted embryonic neurons integrate into adult neocortical circuits. *Nature* **539**, 248–253.
- 167 Gaspard N, Bouschet T, Hourez R, Dimidschstein J, Naeije G, van den Ameele J, Espuny-Camacho I, Herpoel A, Passante L, Schiffmann SN *et al.* (2008) An intrinsic mechanism of corticogenesis from embryonic stem cells. *Nature* **455**, 351–357.
- 168 Michelsen KA, Acosta-Verdugo S, Benoit-Marand M, Espuny-Camacho I, Gaspard N, Saha B, Gaillard A and Vanderhaeghen P (2015) Area-specific reestablishment of damaged circuits in the adult cerebral cortex by cortical neurons derived from mouse embryonic stem cells. *Neuron* **85**, 982–997.