

2019

Redox-Sensitive Calcium/Calmodulin-Dependent Protein Kinase II α in Angiotensin II Intra-Neuronal Signaling and Hypertension

Urmi Basu
University of Nebraska Medical Center

Adam J. Case
University of Nebraska Medical Center, adam.case@unmc.edu

Jinxu Liu
University of Nebraska Medical Center, jinxu.liu@unmc.edu

Jun Tian
University of Nebraska Medical Center, jutian@nebraskamed.com

Yulong Li
University of Nebraska Medical Center, yulongli@unmc.edu

See next page for additional authors

Follow this and additional works at: https://digitalcommons.unmc.edu/com_cell_articles



Part of the [Cellular and Molecular Physiology Commons](#), [Medical Physiology Commons](#), and the [Systems and Integrative Physiology Commons](#)

Recommended Citation

Basu, Urmi; Case, Adam J.; Liu, Jinxu; Tian, Jun; Li, Yulong; and Zimmerman, Matthew C., "Redox-Sensitive Calcium/Calmodulin-Dependent Protein Kinase II α in Angiotensin II Intra-Neuronal Signaling and Hypertension" (2019). *Journal Articles: Cellular & Integrative Physiology*. 32.
https://digitalcommons.unmc.edu/com_cell_articles/32

This Article is brought to you for free and open access by the Cellular & Integrative Physiology at DigitalCommons@UNMC. It has been accepted for inclusion in Journal Articles: Cellular & Integrative Physiology by an authorized administrator of DigitalCommons@UNMC. For more information, please contact digitalcommons@unmc.edu.

Authors

Urmi Basu, Adam J. Case, Jinxu Liu, Jun Tian, Yulong Li, and Matthew C. Zimmerman



ELSEVIER

Contents lists available at ScienceDirect

Redox Biology

journal homepage: www.elsevier.com/locate/redox

Redox-sensitive calcium/calmodulin-dependent protein kinase II α in angiotensin II intra-neuronal signaling and hypertension

Urmi Basu^a, Adam J. Case^a, Jinxu Liu^a, Jun Tian^a, Yu-Long Li^{a,b}, Matthew C. Zimmerman^{a,*}

^a Department of Cellular and Integrative Physiology, University of Nebraska Medical Center, Omaha, NE, USA

^b Emergency Medicine, University of Nebraska Medical Center, Omaha, NE, USA

ARTICLE INFO

Keywords:

Angiotensin II
Reactive oxygen species
Calcium/calmodulin-dependent protein kinase II α
Neurons
Hypertension
Redox signaling

ABSTRACT

Dysregulation of brain angiotensin II (AngII) signaling results in modulation of neuronal ion channel activity, an increase in neuronal firing, enhanced sympathoexcitation, and subsequently elevated blood pressure. Studies over the past two decades have shown that these AngII responses are mediated, in part, by reactive oxygen species (ROS). However, the redox-sensitive target(s) that are directly acted upon by these ROS to execute the AngII pathophysiological responses in neurons remain unclear. Calcium/calmodulin-dependent protein kinase II (CaMKII) is an AngII-activated intra-neuronal signaling protein, which has been suggested to be redox sensitive as overexpressing the antioxidant enzyme superoxide dismutase attenuates AngII-induced activation of CaMKII. Herein, we hypothesized that the neuronal isoform of CaMKII, CaMKII- α (CaMKII α), is a redox-sensitive target of AngII, and that mutation of potentially redox-sensitive amino acids in CaMKII α influences AngII-mediated intra-neuronal signaling and hypertension. Adenoviral vectors expressing wild-type mouse CaMKII α (Ad.wtCaMKII α) or mutant CaMKII α (Ad.mutCaMKII α) with C280A and M281V mutations were generated to overexpress either CaMKII α isoform in mouse catecholaminergic cultured neurons (CATH.a) or in the brain subfornical organ (SFO) of hypertensive mice. Overexpressing wtCaMKII α exacerbated AngII pathophysiological responses as observed by a potentiation of AngII-induced inhibition of voltage-gated K⁺ current, enhanced *in vivo* pressor response following intracerebroventricular injection of AngII, and sensitization to chronic peripheral infusion of AngII resulting in a more rapid increase in blood pressure. In contrast, expressing the mutant CaMKII α in CATH.a neurons or the SFO failed to intensify these AngII responses. Taken together, these data identify neuronal CaMKII α as a redox-sensitive signaling protein that contributes to AngII-induced neuronal activation and hypertension.

1. Introduction

Elevated angiotensin II (AngII)-dependent signaling in the central nervous system has been implicated in the pathogenesis of several neuro-cardiovascular disorders, including hypertension and heart failure [1,2]. Numerous previous studies have shown that AngII signaling in central neurons leads to excessive influx of calcium, activation of NADPH oxidases causing increased production of reactive oxygen species (ROS), particularly superoxide (O₂⁻), and regulation of voltage-gated potassium (K⁺) channels [3–10]. Yin et al. demonstrated that the well-characterized AngII-mediated inhibition of outward K⁺ current (I_{Kv}) is mediated by O₂⁻ as overexpression of superoxide dismutases (SOD), which specifically scavenges O₂⁻, completely abolishes this AngII-response [8]. Several studies by Summers, Raizada, and

colleagues have demonstrated that the AngII-mediated reduction in I_{Kv} and increase in neuronal firing rate also involves specific kinases, such as calcium/calmodulin-dependent protein kinase II (CaMKII) [6,9]. However, the precise signaling mechanism(s) by which ROS and these kinases communicate with each other, if at all, to alter neuronal ion channel activity, neuronal activation, and ultimately an increase in systemic blood pressure remain unknown.

CaMKII is a multifunctional enzyme which couples increases in intracellular Ca²⁺ to regulation of ion channels, transcription of genes and apoptosis [11–13]. Under resting conditions CaMKII remains inactive, but upon binding with Ca²⁺/CaM the autoinhibitory regulatory domain of CaMKII is relieved from the kinase domain causing CaMKII activation. Autophosphorylation of a threonine residue (T286 in CaMKII α) within the autoinhibitory region of CaMKII then leads to a Ca²⁺/

* Corresponding author. Department of Cellular and Integrative Physiology, Free Radicals in Medicine Program, University of Nebraska Medical Center, 985850 Nebraska Medical Center, Omaha, NE, 68198-5850, USA.

E-mail address: mczimmerman@unmc.edu (M.C. Zimmerman).

<https://doi.org/10.1016/j.redox.2019.101230>

Received 9 March 2019; Received in revised form 16 May 2019; Accepted 19 May 2019

2213-2317/© 2019 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

CaM independent CaMKII activity by preventing the reassociation of the autoinhibitory region with the kinase domain. Studies have shown that AngII-mediated pathological responses in the heart involve a second Ca^{2+} /CaM independent activity regulated by pro-oxidant conditions [12,14]. More specifically, Erickson et al. have demonstrated that AngII-generated ROS oxidize paired methionine residues (M281/282) in CaMKII-delta (CaMKII δ), the cardiomyocyte-localized CaMKII isoform, leading to sustained CaMKII activity and contributing to the pathogenesis of cardiac disease [12]. Interestingly, and of particular relevance to the current study, the paired methionines (M281/282) in CaMKII δ are replaced by a cysteine (C280) neighboring a methionine (M281) in the neuronal CaMKII α isoform. Considering cysteine, like methionine, contains a redox-sensitive thiol, we hypothesize that C280 and M281 in CaMKII α are susceptible to ROS-mediated post-translational modification.

Within the brain, AngII acts on specific nuclei that are important in autonomic control of cardiovascular function, including the paraventricular nucleus (PVN) [16], rostral ventrolateral medulla (RVLM) [17], and nucleus tractus solitarius (NTS). AngII can also activate neurons located in the blood-brain barrier deficient circumventricular organs, such as the subfornical organ (SFO) [18], resulting in enhanced sympathoexcitation. Increased sympathetic output is known to contribute to hypertensive symptoms, including increased vasoconstriction, enhanced salt and water reabsorption, increased heart rate, and activation of T-lymphocytes and inflammatory cytokines [19–23]. Past studies have also demonstrated that scavenging O_2^- with adenoviral-mediated overexpression of CuZnSOD (AdCuZnSOD) in the SFO attenuates blood pressure in an AngII-infused hypertensive animal model [4]. However, the role of downstream effector molecules that may be sensitive to changes in the redox environment, such as CaMKII α , in the brain in mediating the AngII-induced hypertensive response is unclear.

In the present study, we tested the hypothesis that C280 and M281 in CaMKII α are redox sensitive in AngII-stimulated neurons leading to sustained CaMKII α activation, which drives the inhibition of I_{Kv} and contributes to AngII-induced hypertension. We observed that overexpression of wild-type CaMKII α in cultured neurons or in the brain SFO potentiated AngII-induced reduction of I_{Kv} and exacerbated the increase in blood pressure mediated by AngII stimulation of the SFO, respectively. In contrast, expression of a mutant CaMKII α (mutCaMKII α) in which C280 was mutated to an alanine (C280A) and M281 was mutated to a valine (M281V) failed to exacerbate the AngII-induced inhibition of I_{Kv} and also prevented the excessive AngII-induced increase in blood pressure. Overall, this study implicates that redox-sensitive post-translational modification of CaMKII α is involved in mediating AngII intra-neuronal signaling and the chronic AngII-mediated neurogenic hypertensive response.

2. Methods

2.1. Neuronal cell culture

Mouse catecholaminergic CATH.a neurons (American Type Culture Collection (ATCC), stock no. CRL-11179) were cultured in RPMI 1640 medium supplemented with 8% normal horse serum (NHS), 4% fetal bovine serum, and 1% penicillin-streptomycin at 37 °C with 5% CO_2 as recommended by ATCC. CATH.a neurons were differentiated for 6–8 days before experimentation by adding *N* [6],2'-*O*-dibutyryl adenosine 3',5' - cyclic monophosphate sodium salt (1 mM, Sigma-Aldrich, St. Louis, MO) to the culture medium as previously described [8]. It should be noted that CATH.a neurons express both AngII type 1 (AT1) and type 2 (AT2) receptors and have been widely used by various groups including ours to study AngII intra-neuronal signaling [6,8,24].

2.2. Construction and generation of wild-type CaMKII α adenovirus

Replication-deficient recombinant adenovirus (Ad5-CMV) encoding

mouse wild-type calcium/calmodulin-dependent protein kinase II α (Ad.wtCaMKII α) was generated. Briefly, CaMKII α plasmid (Origene) was amplified by conventional Polymerase Chain Reaction (PCR) using the HotStart PCR Master Mix (Qiagen, Venlo, Limburg) and the following primers: forward 5'- GAA TTC ATG GCT ACC ATC ACC TGC ACC C - 3'; reverse 5' - GGA TCC TCA ATG CGG CAG GAC GGA - 3'. The PCR product was then run on a 1% agarose gel, followed by gel extraction utilizing Qiagen gel-extraction kit (Qiagen). The CaMKII α insert was cloned using a pJet1.2 clone jet kit (Thermo Scientific), transformed into DH5 alpha bacteria and grown overnight onto ampicillin treated plates at 37°C in a bacterial incubator. Bacterial colonies were picked and grown in LB media overnight by shaking. PureLink Quick Plasmid DNA Miniprep Kits (Invitrogen, Grand Island, NY) was used to isolate CaMKII plasmid DNA as per manufacturer's instructions. Following quantification of the DNA using Nanodrop 200 spectrophotometer (Thermo Scientific, Waltham, MA) 1 μg DNA was digested with EcoRI and BamHI restriction enzymes and then run on a 1% agarose gel to confirm the presence of the correct insert. The plasmid samples were then sent off to the UNMC Gene Sequencing Core. Once the CaMKII α sequence was confirmed it was then cloned into an expression plasmid, transformed into bacteria, DNA isolated, digested using restriction enzymes and again sent to the UNMC Gene sequencing core to confirm the presence of correct CaMKII α sequence in the expression plasmid. Adenoviral vectors were then constructed, purified and provided by the University of Iowa Gene Vector Core, as previously described [25]. To determine the optimal overexpression of wtCaMKII α , CATH.a neurons were transduced with Ad.wtCaMKII α on day 3 of differentiation at 10, 25 and 50 MOI and Western blot analysis was performed to confirm CaMKII α overexpression. Empty adenoviral vectors (Ad.Empty) or GFP-expressing adenovirus (Ad.GFP) were used as controls.

2.3. Site-directed mutagenesis and mutant CaMKII α adenovirus generation

Replication-deficient recombinant adenovirus (Ad5-CMV) encoding mouse mutant calcium/calmodulin-dependent protein kinase II α (Ad.mutCaMKII α) was generated. Primer sequences were made specifically to mutate cysteine 280 to alanine (C280A) and methionine 281 to valine (M281V) with overlapped extension polymerase chain reaction (PCR). Briefly two sections of the wt-CaMKII α plasmid (Origene) were amplified by conventional PCR using HotStart PCR Master Mix (Qiagen, Venlo, Limburg) with the following two set of primers: Section 1 forward 5'- GAA TTC ATG GCT ACC ATC ACC TGC ACC C - 3'; reverse 5' - CCA CGG TCT CCT GTC TGT GCA - 3'. Section 2 forward 5'- GTG CAC AGA CAG GAG ACC ACC GTG - 3'; reverse 5' - GGA TCC TCA ATG CGG CAG GAC GGA - 3'. The next set of steps involving cloning using pJet1.2 PCR clone jet kit (Thermo Scientific) and bacterial colony culture was similar to the Ad.wtCaMKII α generation procedure described above. Adenoviral vectors encoding mutCaMKII α were constructed, purified and provided by the University of Iowa Gene Vector Core, as previously described [25].

2.4. Western blot analysis

Protein expression of CaMKII α and actin was determined using standard Western blot analysis in lysates prepared from differentiated CATH.a neurons either non-transduced or transduced (10–50 MOI) with Ad.wtCaMKII α or Ad.mutCaMKII α . Briefly, cells were harvested in lysis buffer containing protease inhibitor cocktail (Sigma-Aldrich) and phosphatase inhibitor (Thermo Scientific). Cell lysates (25–30 μg of protein per lane) were loaded onto gels, separated by SDS-PAGE and transferred to a nitrocellulose membrane. Membranes were incubated with primary antibodies directed against CaMKII α (1:1000 dilution, Abcam, Cambridge, UK) and actin (1:1000 dilution, Sigma-Aldrich) at 4 °C overnight. After incubation with anti-rabbit horseradish peroxidase-conjugated secondary antibody (1:5000 dilution, Thermo Scientific, Waltham, MA), bands were visualized using the Pierce

enhanced chemiluminescence detection system (Thermo Scientific, Rockford, IL). Densitometric analysis of band density was determined using NIH ImageJ analysis software. Values were normalized to actin to correct for any variations in protein loading.

2.5. Electrophysiological record of voltage-gated K^+ currents

Differentiated CATH.a neurons either non-transduced or transduced with control adenovirus (Ad.Empty; 25MOI), Ad.wtCaMKII α (25MOI) or Ad.mutCaMKII α (25MOI) were used to measure K^+ currents (I_{KV}). I_{KV} was recorded by the whole-cell configuration of the patch-clamp technique using an Axopatch 200-B patch-clamp amplifier (Axon Instruments, Inc, Union City, CA). Briefly, in the voltage-clamp experiments, resistance of the patch pipette was 4–6 M Ω when filled with the following solution (in mM): 130 KCl, 2 MgCl $_2$, 0.25 CaCl $_2$, 5 EGTA, 1 Mg-ATP, 0.1 Tris-GTP, 10 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES), and 8 glucose, pH 7.2. The extracellular solution consisted of (in mM): 137 NaCl, 5.4 KCl, 1.35 CaCl $_2$, 2 MgCl $_2$, 0.3 NaH $_2$ PO $_4$, 10 HEPES, 10 sucrose, pH 7.4. To selectively record I_{KV} , Na $^+$ and Ca $^{2+}$ channels were blocked by 0.5 μ M tetrodotoxin (TTX, Sigma-Aldrich) and 0.3 mM CdCl $_2$ (Sigma Aldrich) respectively. Current traces were sampled at 10 kHz and filtered at 5 kHz. Holding potential was -80 mV and current-voltage (I–V) relationship was elicited by 10 mV step increments to potentials between -80 and $+80$ mV for 400 ms. The change of I_{KV} with AngII was tested by superfusing CATH.a neurons with AngII (100 nM) for 5 min and repeating the voltage pulse regimen. To confirm the observed changes with AngII, I_{KV} was recorded again after 15 min of washout. Cells responded to AngII stimulation and then recovered with the washout were considered for data analysis. Current density of I_{KV} was calculated by dividing the respective current by cell membrane capacitance (C_m). The C_m of CATH.a neurons utilized in the study ranged from 10 to 60 pF and was not significantly different between the groups. The pClamp 10.2 program (Axon Instruments) was used for data acquisition and analysis. All experiments were carried out at room temperature.

2.6. Mouse studies

All animal procedures were reviewed and approved by the University of Nebraska Medical Center Institutional Animal Care and Use Committee. Adult C57BL/6 male mice (20–25 g body weight; Harlan, Indianapolis, IN) were used for physiological and immunohistochemical studies. Mice were provided standard chow and water ad libitum. Mice were surgically instrumented with intracerebroventricular (ICV) cannulas for administration of either AngII or adenoviruses to the brain SFO. Mean arterial pressure (MAP), systolic and diastolic blood pressure, and heart rate (HR) were recorded daily with radiotelemetry (Data Sciences International, DSI), as previously described [4,24]. Following 1 week of recovery from surgery, blood pressure and heart rate responses were recorded in conscious, unrestrained mice immediately prior to, during and after acute ICV injection of AngII (350 ng). Then, mice were ICV injected with Ad.GFP (control adenovirus) or Ad.wtCaMKII α (5×10^7 plaque forming units), and three days later mice were again ICV injected with AngII (350 ng) while blood pressure and heart rate were monitored.

To generate AngII-dependent hypertensive mice, osmotic minipumps (Alzet, Durect) were implanted subcutaneously 3 days after ICV injection of adenovirus (Ad.GFP, Ad.wtCaMKII α , and Ad.mutCaMKII α) and delivered AngII (400 ng/kg/min) for approximately 3 weeks [4,24]. Control mice received subcutaneous osmotic minipumps filled with saline.

2.7. Immunohistochemistry

Mice were euthanized with intraperitoneal injection of fatal plus (40 mg/kg, Henry-Schrein, Inc., Melville, NY) after 7 days of ICV

adenovirus injection. Brains were removed and fixed in 4% paraformaldehyde in 0.1 M phosphate buffer (PB) and then stored in 30% sucrose in PB overnight prior to sectioning. Cryostat brains sections (20 μ m) containing the SFO were prepared and immunohistochemistry was carried out, as previously described [24]. Briefly, sections were incubated for 1 h with blocking buffer (10% NHS in 0.1 M PB). Tissue sections were then incubated at 4 $^{\circ}$ C overnight with primary antibody against CaMKII α (1:200, rabbit monoclonal antibody, Abcam), prepared in 0.1 M PB containing 2% NHS and 0.3% Triton X-100. The sections were then washed in 0.1 M PB and incubated with Alexa Fluor 598 secondary antibody (1:200 dilution, Invitrogen, Carlsbad, CA) at room temperature for 2 h. Fluorescent images were detected by confocal laser scanning microscopy (Zeiss LSM 510 Meta).

2.8. Statistical analysis

All data are presented as means \pm standard error of the mean (SEM). Data were analyzed by Student's t-test for two group comparisons or by ANOVA for multiple group comparisons with Newman-Kuels post-hoc test. P -value less than 0.05 was considered statistically significant. Statistical analyses were performed using GraphPad Prism 5.0 statistical and graphical software.

3. Results

3.1. Wild-type CaMKII α overexpression, but not mutant CaMKII α , potentiates AngII-mediated inhibition of neuronal K^+ current (I_{KV}) in CATH.a neurons

To investigate the effect of wtCaMKII α overexpression on the AngII-dependent changes in neuronal electrophysiological properties, I_{KV} was measured in non-transduced, Ad.Empty (25 MOI) or Ad.wtCaMKII α (25 MOI)-transduced CATH.a neurons. Consistent with previous studies [8], acute AngII (100 nM) superfusion for 5 min inhibited I_{KV} in non-transduced and Ad.Empty-transduced CATH.a neurons, as seen in the representative traces of evoked K^+ currents (Fig. 1A). Interestingly, AngII superfusion caused a further significant reduction in I_{KV} (Fig. 1A) in Ad.wtCaMKII α -transduced CATH.a neurons ($n = 7$) compared to non-transduced ($n = 7$) or Ad.Empty-treated neurons ($n = 7$), as shown in the summary data (Fig. 1B). In contrast, expression of mutant CaMKII α in CATH.a neurons via adenovirus-mediated gene transfer (Ad.mutCaMKII α , 25 MOI) failed to potentiate the AngII-induced inhibition of I_{KV} (Fig. 1B). Western blot analysis confirmed overexpression of wtCaMKII α or mutant CaMKII α in neurons transduced with Ad.wtCaMKII α or Ad.mutCaMKII α , respectively, compared to non-transduced and Ad.Empty-treated cells (Fig. S1). These data indicate that overexpression of wtCaMKII α exacerbates the AngII-induced inhibition of neuronal K^+ channel current, and, more importantly, suggest that cysteine 280 and methionine 281 in CaMKII α are required to mediate this potentiated AngII response.

3.2. Overexpression of wtCaMKII α in the brain SFO potentiates the central acute AngII-induced increase in mean arterial pressure (MAP)

To determine the role of wtCaMKII α overexpression in the brain SFO on the acute, central actions of AngII *in vivo*, C57BL/6 male mice underwent radiotelemetry implantation and ICV cannula surgeries. Following a week's recovery, blood pressure and heart rate responses to ICV administered AngII (350 ng) were recorded in conscious mice. As seen in the representative recordings, mice receiving acute ICV AngII prior to ICV administration of Ad.GFP or Ad.wtCaMKII α exhibited the characteristic increase in blood pressure as compared to their baseline (Fig. 2A). Interestingly, 3 days post-adenovirus injection, the AngII-mediated rise in blood pressure was potentiated in mice ICV-injected with Ad.wtCaMKII α compared to Ad.GFP-injected mice. The peak change in MAP following acute ICV AngII injections in Ad.wtCaMKII α




Fig. 1. Wild-type CaMKII α over-expression, but not mutant CaMKII α , potentiates the AngII-mediated inhibition of outward K⁺ current in CATH.a neurons. A) Representative electrophysiological traces of evoked K⁺ currents in non-transduced, Ad.Empty (25 MOI) or Ad.wtCaMKII α (25 MOI)-transduced CATH.a neurons superfused with vehicle or 100 nM AngII. B) Percent inhibition of outward K⁺ current following 5 min of AngII (100 nM) superfusion in non-transduced, Ad.Empty (25 MOI), Ad.wtCaMKII α (25 MOI) or Ad.mutCaMKII α (25 MOI)-transduced CATH.a neurons. * $p < 0.05$ vs. non-transduced and Ad.Empty. # $p < 0.05$ vs. Ad.wtCaMKII α . $n = 4-8$ cells/group.

($n = 9$) and AdGFP ($n = 9$) mice are summarized in Fig. 2B. Heart rate did not change significantly between the groups (data not shown). In addition, the duration of the AngII-induced pressor response was longer in wtCaMKII α overexpressing mice as compared to mice with Ad.GFP (Fig. 2C). The duration was calculated from the start of the peak AngII-mediated rise in blood pressure until the time when blood pressures came back to half the maximal response. Together, these data demonstrate that overexpression of wild-type CaMKII α in the brain SFO exacerbates the central AngII-induced pressor response and suggest that CaMKII α in the brain is an important player in central AngII-mediated cardiovascular effects.

3.3. Wild-type CaMKII α overexpression, but not mutant CaMKII α , in the brain SFO sensitizes mice to an immediate and sustained increase in blood pressure during subcutaneous AngII infusion

We next evaluated the effect of overexpressing wtCaMKII α or mutCaMKII α in the brain SFO on the development and progression of AngII-induced hypertension. Similar to the acute AngII studies (Fig. 2),

mice were implanted with radiotelemeters for blood pressure monitoring and ICV cannulas for adenovirus injections. Following 1 week of recovery from surgery, Ad.GFP, Ad.wtCaMKII α or Ad.mutCaMKII α was administered ICV. Subcutaneous AngII (400 ng/kg/min) or saline osmotic minipumps were implanted 3 days post-adenovirus injection and blood pressure and heart rate were recorded daily in conscious mice for the next 3 weeks. As shown in Fig. 3, MAP, as well as systolic and diastolic blood pressure in Ad.GFP-injected mice ($n = 8$) increased gradually with AngII infusion with a peak at around day 16. Elevated blood pressures then returned to baseline levels at day 22–23 when the osmotic minipumps emptied. It should be noted that the gradual increase in blood pressure in Ad.GFP-treated mice during subcutaneous infusion of 400 ng/kg/min of AngII is characteristic of the slow-pressor AngII-induced hypertensive model [4,26,27]. Interestingly, in Ad-wtCaMKII α -injected mice ($n = 9$) AngII infusion resulted in a significant increase in blood pressure within 2 days that continued to increase until day 23 when the minipumps emptied and blood pressure returned back to basal levels. In mice expressing mutCaMKII α in the SFO ($n = 6$), there was an increase in blood pressure within 2 days of




Fig. 2. Overexpression of wtCaMKII α in the brain SFO potentiates the pressor response of central acute AngII. A) Representative blood pressure tracings and (B) summary data showing AngII-induced acute pressor response in mice ICV injected AngII (350 ng) both pre- and 3 days post-ICV injection of Ad.GFP or Ad.wtCaMKII α . C) Duration (in minutes) of the AngII-mediated rise in blood pressure in Ad.GFP and Ad.wtCaMKII α injected mice. * $p < 0.05$ vs. Ad.GFP. $n = 8-9$ mice/group.

starting AngII, which was similar to the response in Ad.wtCaMKII α -treated mice. However, blood pressures declined thereafter and continued to rise gradually similar to the Ad.GFP group. The peak increase in MAP (day 15–17 of AngII infusion) was significantly attenuated in

the mutCaMKII α mice (115 mmHg) as compared to the wtCaMKII α group (133 mmHg). Heart rate was not significantly different among the three groups infused with AngII (data not shown). Blood pressure and heart rate responses did not differ in saline-infused mice ICV




Fig. 3. Wild-type CaMKII α overexpression, but not mutant CaMKII α , in the brain SFO sensitizes mice to an immediate and sustained AngII-induced increase in blood pressure. Summary data of (A) mean arterial pressure, (B) systolic blood pressure, and (C) diastolic blood pressure, as measured by radiotelemetry in C57BL/6 male mice at baseline, following ICV injection of Ad.GFP, Ad.wtCaMKII α or Ad.mutCaMKII α , and after subcutaneous implantation of osmotic minipumps set to release AngII (400 ng/kg/min). * $p < 0.05$ vs. Ad.wtCaMKII α . # $p < 0.05$ vs. Ad.GFP mice. $n = 6-9$ mice/group.




Fig. 4. Adenoviral-mediated over-expression of wtCaMKII α and mutant CaMKII α in the brain SFO. Representative confocal microscopy images showing expression of GFP and CaMKII α in the SFO of mice 1 week after ICV injection of Ad.GFP, Ad.wtCaMKII α or Ad.mutCaMKII α . Coronal brain sections (20 μ m) were stained for GFP (green fluorescence), and CaMKII α (red fluorescence). Differential interference contrast (DIC) images show entire SFO tissue. Scale bar = 100 μ m. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

injected with Ad.GFP, Ad.wtCaMKII α or Ad.mutCaMKII α (data not shown).

Importantly, to confirm efficient adenovirus-mediated over-expression of wtCaMKII α and mutCaMKII α in the SFO following ICV administration of Ad.wtCaMKII α or Ad.mutCaMKII α , respectively, immunohistochemistry was performed on brain tissue sections 7 days post ICV adenovirus injections. Control mice received ICV Ad.GFP. As shown in the representative confocal microscopy images (Fig. 4), the SFO from Ad.wtCaMKII α - and Ad.mutCaMKII α -injected mouse brains exhibited higher levels of CaMKII α protein levels in comparison to the SFO from Ad.GFP-treated mice. Overall, these studies indicate that wtCaMKII α overexpression in the brain SFO sensitizes mice to an immediate increase in blood pressure at an initial subpressor dose of subcutaneous AngII. Furthermore, mutation of redox-sensitive cysteine and methionine residues in CaMKII α prevents the sensitization mechanism and attenuates the peak AngII-induced rise in blood pressure.

4. Discussion

Dysregulation in AngII intra-neuronal signaling mechanisms contributes to the pathogenesis of neuro-cardiovascular diseases, such as hypertension and heart failure [1,2]. In central neurons, AngII activates redox mechanisms including NADPH oxidases and increases generation of reactive oxygen species, particularly superoxide [5]. Many studies have clearly demonstrated that elevated levels of reactive oxygen species in neurons located in cardiovascular control brain regions mediate enhanced neuronal activation and drive sympathoexcitation, which are known to increase systemic blood pressure [3–5,23]. However, the underlying mechanism(s) by which an increase in neuronal reactive oxygen species following AngII stimulation leads to neuronal firing, sympathoexcitation, and elevated in blood pressure remains unclear. Herein, we report that adenoviral-mediated overexpression of wild-type CaMKII α (Ad.wtCaMKII α), the isoform of CaMKII highly expressed in neurons, in central neurons potentiates the AngII-induced reduction in neuronal K⁺ current and AngII-induced hypertension. In addition, mutation of specific cysteine (C280) and methionine (M281) residues in CaMKII α attenuates the potentiated AngII response observed with wtCaMKII α overexpression. The main findings of this study can be summarized as follows: (1) the potentiated AngII-mediated inhibition of

neuronal K⁺ current (I_{KV}) following wtCaMKII α overexpression in CATH.a neurons is absent in mutant CaMKII α -transduced neurons; (2) overexpression of wtCaMKII α in the brain SFO potentiates the central acute AngII-induced pressor response; (3) wtCaMKII α overexpression in the SFO sensitizes mice to an immediate increase in blood pressure at an initial subpressor dose of subcutaneous chronic AngII infusion; and (4) mutant CaMKII α in the SFO prevents the sensitization mechanism and attenuates the peak AngII-induced rise in blood pressure.

Previous studies have identified signaling proteins, such as CaMKII α and PKC, as downstream mediators of AngII signaling in central neurons [6,7,9]. These studies have shown a clear contribution of CaMKII α in mediating the AngII-induced inhibition of neuronal ion current, specifically K⁺ current, and the subsequent increase in neuronal firing both in primary neuronal cultures as well as in neuronal cell lines. Previous results from our lab suggest redox-dependent activation of CaMKII α , as measured by increased CaMKII α phosphorylation in AngII-stimulated CATH.a neurons that could be blocked by overexpressing superoxide dismutase [8]. Additionally, Erickson et al. have demonstrated that AngII can induce oxidation of cardiac CaMKII (CaMKII δ) at paired methionine residue (M281/282) and sustain CaMKII δ activity independent of Ca²⁺/CaM binding [12]. This sustained activity of CaMKII δ plays an important role in AngII-mediated apoptosis in ischemic heart and under myocardial infarction conditions [12–14]. Further studies have also elucidated that CaMKII α can undergo S-nitrosylation at cysteine 280 and 289, and mutation of either cysteine residue attenuates nitric oxide-induced excitotoxic neuronal cell death [15]. Although these studies indicate the importance of reactive oxygen species in CaMKII activation, the role of redox sensitive CaMKII α in AngII-dependent intra-neuronal signaling and AngII-mediated hypertension is unknown.

To determine the role of redox-dependent activation CaMKII α in mediating AngII-induced reduction in neuronal I_{KV} , we used adenoviral-mediated overexpression of wtCaMKII α in CATH.a neurons and measured changes in outward K⁺ current. AngII superfusion inhibited I_{KV} by almost 16% both in non-transduced as well as in Ad.Empty-transduced neurons. Interestingly, this effect was significantly exaggerated in Ad.wtCaMKII α -transduced neurons (37% inhibition). In contrast, CATH.a neurons that were transduced with mutCaMKII α adenovirus had a 13% inhibition in outward K⁺ current, which was similar to

control neurons. These results strongly suggest that cysteine 280 and methionine 281 are required for wild-type CaMKII α to potentiate the AngII-induced inhibition of outward I_{K_v} . Further studies are needed to determine if overexpression of wtCaMKII α in neurons results in changes in protein levels and/or membrane localization of K^+ channels, or if the potentiated AngII response is a result of wild-type CaMKII α directly acting on K^+ channels to regulate their activation.

Brain AngII is known as a potent regulator of arterial blood pressure and when administered acutely via ICV it elicits a well-characterized increase in blood pressure [16–19]. As seen in the representative blood pressure tracings in Fig. 2, acute ICV injection of AngII caused a rapid and relatively short-lived increase in mean arterial pressure in Ad.GFP-treated mice. This AngII response was significantly potentiated in mice overexpressing wtCaMKII α in the SFO. We did not observe changes in baseline blood pressure with wtCaMKII α overexpression. We next wanted to evaluate the role of CaMKII α in the SFO on the development and progression of AngII-dependent hypertension. In order to do these studies, we utilized the slow-pressor chronic AngII-induced hypertensive model that has been extensively used by our lab and others to investigate intra-neuronal signaling mechanisms in AngII-mediated hypertension [4,24,26,27]. Intriguingly, we observed a significant rise in blood pressure in Ad.wtCaMKII α -injected mice immediately (i.e. within 2 days) after the start of AngII infusion as compared to a more gradual rise in Ad.GFP-injected mice. Further, this robust and immediate increase in blood pressure was sustained throughout AngII infusion and resulted in the peak rise in blood pressure being significantly higher in Ad.wtCaMKII α injected mice as compared to Ad.GFP group (133 mmHg in Ad.wtCaMKII α vs. 125 mmHg in AdGFP group at days 15–17). We conclude that mice overexpressing wtCaMKII α in the brain SFO are sensitized to the initial subpressor dose of AngII infusion used in our study (i.e. 400 ng/kg/min). Although further studies are needed to determine the precise mechanism of action by which wtCaMKII α overexpression in the SFO sensitize mice to peripherally infused AngII, we predict the elevated levels of wtCaMKII enhance AngII-dependent intra-neuronal signaling in neurons residing in the blood-brain barrier deficient SFO. Further, we speculate that this enhanced signaling results in activation of SFO neurons, perhaps via potentiation of outward K^+ current, projecting to cardiovascular control regions behind the blood-brain barrier, such as the paraventricular nucleus (PVN), which are known to drive sympathoexcitation and increase blood pressure.

Lastly, in our current study, we observed that mice expressing mutant CaMKII α in the SFO had a similar rise in blood pressure within 2 days of AngII infusion as observed in the wtCaMKII α group. However, blood pressure declined after 2 days and followed a more gradual increase, similar to the Ad.GFP-injected mice infused with AngII. The peak AngII-induced increase in blood pressure in mice with mutCaMKII α was also significantly lower (115 mmHg). We speculate that the initial rise in blood pressure in mutCaMKII α is due to a similar sensitization mechanism as seen in wtCaMKII α mice. However, considering cysteine 280 and methionine 281 are mutated and previous studies suggest the redox modulation of CaMKII results in sustained activity even in the absence of Ca^{2+} /calmodulin [12,14], we predict this sensitization is lost as the mutant CaMKII α is unable to sustain its activity which results in the subsequent gradual increase in blood pressure similar to Ad.GFP mice.

Although the current study provides new insight into AngII intra-neuronal redox signaling mechanisms and indicates that C280 and M281 of neuronal CaMKII α are redox-sensitive targets of AngII, there are various limitations of this study that must be addressed. First, considering our mutant CaMKII α included mutation of C280 and M281, it remains unknown if one or both of these redox-sensitive amino acids contribute to the actions of CaMKII α in AngII stimulated neurons. Second, numerous previous studies from various groups, including ours [3–5,8,24], have identified superoxide as the primary ROS elevated in AngII-stimulated neurons. However, it is unlikely that the thiol in

cysteine and methionine is sensitive to oxidation by superoxide. As such, it remains unclear precisely how C280 and/or M281 in CaMKII α are oxidized in AngII activated neurons. Additional studies are needed to determine if hydrogen peroxide or some other ROS are involved in oxidizing C280 and/or M281. Lastly, future studies involving mass spectrometry and proteomic approaches are needed to definitively describe the oxidative post-translational modifications of CaMKII α in AngII-stimulated neurons.

In summary, the experimental data presented herein indicate that neuronal CaMKII α is an important redox-sensitive signaling protein contributing to AngII intra-neuronal signaling and hypertension. More specifically, we showed that wtCaMKII α overexpression exacerbates the AngII-mediated inhibition of neuronal K^+ current, which was blocked with mutation of redox-sensitive residues (i.e. C280 and M281) in CaMKII α . In addition, wtCaMKII α overexpression in the SFO potentiated the central acute AngII-mediated pressor response, and sensitized mice to an immediate increase in blood pressure at an initial subpressor dose of subcutaneous chronic AngII. Importantly, mice expressing mutCaMKII α in the SFO were protected from the sensitization mechanism and also had a significant lower peak AngII response. In conclusion, these studies may help to identify potential redox-sensitive therapeutic targets, such as CaMKII α , for the improved treatment of AngII-dependent hypertension.

Acknowledgments

This study was supported by a National Institutes of Health Grant R01-HL103942 (M.C. Zimmerman). We thank the University of Nebraska Medical Center Advanced Microscopy Core Facility for providing assistance with confocal microscopy. This Core facility receives partial support from the National Institute of General Medical Sciences INBRE - P20 GM103427 and COBRE - P30 GM106397 grants, as well as support from the National Cancer Institute for The Fred & Pamela Buffett Cancer Center Support Grant- P30 CA036727, and the Nebraska Research Initiative. This publication's contents and interpretations are the sole responsibility of the authors.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.redox.2019.101230>.

References

- [1] S.J. Veerasingham, M.K. Raizada, Brain renin-angiotensin system dysfunction in hypertension: recent advances and perspectives, *Br. J. Pharmacol.* 139 (2) (2003) 191–202.
- [2] I.H. Zucker, Novel mechanisms of sympathetic regulation in chronic heart failure, *Hypertension* 48 (6) (2006) 1005–1011.
- [3] M.C. Zimmerman, R.V. Sharma, R.L. Davison, Superoxide mediates angiotensin II-induced influx of extracellular calcium in neural cells, *Hypertension* 45 (4) (2005) 717–723.
- [4] M.C. Zimmerman, E. Lazartigues, R.V. Sharma, R.L. Davison, Hypertension caused by angiotensin II infusion involves increased superoxide production in the central nervous system, *Circ. Res.* 95 (2) (2004) 210–216.
- [5] M.C. Zimmerman, R.P. Dunlay, E. Lazartigues, et al., Requirement for Rac1-dependent NADPH oxidase in the cardiovascular and dipsogenic actions of angiotensin II in the brain, *Circ. Res.* 95 (5) (2004) 532–539.
- [6] C. Sun, J. Du, M.K. Raizada, C. Summers, Modulation of delayed rectifier potassium current by angiotensin II in CATH.a cells, *Biochem. Biophys. Res. Commun.* 310 (3) (2003) 710–714.
- [7] C. Summers, M. Zhu, C.H. Gelband, P. Posner, Angiotensin II type 1 receptor modulation of neuronal K^+ and Ca^{2+} currents: intracellular mechanisms, *Am. J. Physiol.* 271 (1 Pt 1) (1996) C154–C163.
- [8] J.X. Yin, R.F. Yang, S. Li, et al., Mitochondria-produced superoxide mediates angiotensin II-induced inhibition of neuronal potassium current, *Am. J. Physiol. Cell Physiol.* 298 (4) (2010) C857–C865.
- [9] C. Sun, C. Summers, M.K. Raizada, Chronotropic action of angiotensin II in neurons via protein kinase C and CaMKII, *Hypertension* 39 (2 Pt 2) (2002) 562–566.
- [10] L. Gao, Y. Li, H.D. Schultz, et al., Downregulated Kv4.3 expression in the RVLM as a potential mechanism for sympathoexcitation in rats with chronic heart failure, *Am. J. Physiol. Heart Circ. Physiol.* 298 (3) (2010) H945–H955.

- [11] J. Backs, K. Song, S. Bezprozvannaya, S. Chang, E.N. Olson, CaM kinase II selectively signals to histone deacetylase 4 during cardiomyocyte hypertrophy, *J. Clin. Investig.* 116 (7) (2006) 1853–1864.
- [12] J.R. Erickson, M.L. Joiner, X. Guan, et al., A dynamic pathway for calcium-independent activation of CaMKII by methionine oxidation, *Cell* 133 (3) (2008) 462–474.
- [13] Y. Yang, W.Z. Zhu, M.L. Joiner, et al., Calmodulin kinase II inhibition protects against myocardial cell apoptosis in vivo, *Am. J. Physiol. Heart Circ. Physiol.* 291 (6) (2006) H3065–H3075.
- [14] J.R. Erickson, B.J. He, I.M. Grumbach, M.E. Anderson, CaMKII in the cardiovascular system: sensing redox states, *Physiol. Rev.* 91 (3) (2011) 889–915.
- [15] S.J. Coultrap, K.U. Bayer, Nitric oxide induces Ca²⁺-independent activity of the Ca²⁺/calmodulin-dependent protein kinase II (CaMKII), *J. Biol. Chem.* 289 (28) (2014) 19458–19465.
- [16] J.S. Bains, A. Potyok, A.V. Ferguson, Angiotensin II actions in paraventricular nucleus: functional evidence for neurotransmitter role in efferents originating in subfornical organ, *Brain Res.* 599 (2) (1992) 223–229.
- [17] L. Gao, W. Wang, Y.L. Li, et al., Sympathoexcitation by central ANG II: roles for AT1 receptor upregulation and NAD(P)H oxidase in RVLm, *Am. J. Physiol. Heart Circ. Physiol.* 288 (5) (2005) H2271–H2279.
- [18] A.V. Ferguson, J.S. Bains, Actions of angiotensin in the subfornical organ and area postrema: implications for long term control of autonomic output, *Clin. Exp. Pharmacol. Physiol.* 24 (1) (1997) 96–101.
- [19] A.K. Johnson, R.L. Thunhorst, The neuroendocrinology of thirst and salt appetite: visceral sensory signals and mechanisms of central integration, *Front. Neuroendocrinol.* 18 (3) (1997) 292–353.
- [20] S. Matsukawa, L.C. Keil, I.A. Reid, Role of endogenous angiotensin II in the control of vasopressin secretion during hypovolemia and hypotension in conscious rabbits, *Endocrinology* 128 (1) (1991) 204–210.
- [21] P.J. Marvar, S.R. Thabet, T.J. Guzik, et al., Central and peripheral mechanisms of T-lymphocyte activation and vascular inflammation produced by angiotensin II-induced hypertension, *Circ. Res.* 107 (2) (2010) 263–270.
- [22] I.A. Reid, Interactions between ANG II, sympathetic nervous system, and baroreceptor reflexes in regulation of blood pressure, *Am. J. Physiol.* 262 (6 Pt 1) (1992) E763–E778.
- [23] M.C. Zimmerman, Angiotensin II and angiotensin-1-7 redox signaling in the central nervous system, *Curr. Opin. Pharmacol.* 11 (2) (2011) 138–143.
- [24] A.J. Case, S. Li, U. Basu, J. Tian, M.C. Zimmerman, Mitochondrial-localized NADPH oxidase 4 is a source of superoxide in angiotensin II-stimulated neurons, *Am. J. Physiol. Heart Circ. Physiol.* 305 (1) (2013) H19–H28.
- [25] M.C. Zimmerman, L.W. Oberley, S.W. Flanagan, Mutant SOD1-induced neuronal toxicity is mediated by increased mitochondrial superoxide levels, *J. Neurochem.* 102 (3) (2007) 609–618.
- [26] A. Edgley, M. Kett, W. Anderson, Slow pressor' hypertension from low-dose chronic angiotensin II infusion, *Clin. Exp. Pharmacol. Physiol.* 28 (12) (2001) 1035–1039.
- [27] G. Simon, G. Abraham, G. Cserep, Pressor and subpressor angiotensin II administration. two experimental models of hypertension, *Am. J. Hypertens.* 8 (6) (1995) 645–650.

