

NOTE

BRAZILIAN JOURNAL OF OCEANOGRAPHY, 58(special issue IICBBM):73-77, 2010

NEW HOST RECORD FOR *Livoneca redmanni* (LEACH, 1818)
(ISOPODA: CYMOTHOIDAE) IN THE BRAZILIAN COASTAL WATERS
WITH ASPECTS OF HOST-PARASITE INTERACTION*

Eudriano Florêncio dos Santos Costa^{1,**} and Sathyabama Chellappa^{1,**}

Universidade Federal do Rio Grande do Norte
Centro de Biociência - Programa de Pós-Graduação em Bioecologia Aquática
(Praia de Mãe Luiza, s/n, 59014-100 Natal, RN, Brasil)

**Corresponding authors: eudriano@yahoo.com.br; chellappa.sathyabama63@gmail.com

Cymothoids are among the largest parasites of fishes. These are the isopods commonly seen in numerous families and species of fishes of commercial importance, in tropical and subtropical waters, attached on the body surface, in the mouth or on the gills of their hosts (BRUSCA, 1981; BUNKLEY-WILLIAMS; WILLIAMS, 1998; LESTER; ROUBAL, 2005). Some cymothoids form pouches in the lateral musculature of few freshwater and marine fishes; are highly host and site specific (BRUSCA, 1981; BUNKLEY-WILLIAMS; WILLIAMS, 1998). They are protandrous hermaphrodites which are unable to leave their hosts after becoming females. All gravid females possess a marsupium or brood pouch on the ventral surface of the body, within which the young are held until they become manca, and there is no larval stage (BULLAR, 1876; BUNKLEY-WILLIAMS; WILLIAMS, 1985, 1998; WILLIAMS; WILLIAMS, 1998). The mancae have only six pairs of legs (compared to seven in juveniles and adults), large compound eyes and pleopods with setae with which they swim very rapidly. After a short free-swimming period they need to find a host fish to take the first meal within one to two days or they die (LESTER, 2005). Like most isopods, cymothoids are considered to feed principally on host blood, but they may consume the mucus, epithelium and subcutaneous tissues of their hosts (LANZING; O'CONNOR, 1975; GRABDA, 1991; LESTER; ROUBAL, 1995; BUNKLEY-WILLIAMS; WILLIAMS, 1998; RAMDANE et al., 2007).

The species *Livoneca redmanni* (Leach, 1818) has been found only on cero, *Scomberomorus regalis* and serra Spanish mackerel, *S. brasiliensis* (Osteichthyes: Scombridae), in pairs in the gill-chamber. *L. redmanni* causes extreme damages to these fishes and can kill them, causing significant loss of these valuable fishes. *L. redmanni* occurs in the Caribbean and the South American coasts to Rio de Janeiro, Brazil (WILLIAMS; BUNKLEY-WILLIAMS, 1996).

The fish *Chloroscombrus chrysurus* (Linnaeus, 1766) (Osteichthyes: Carangidae), commonly known as Atlantic bumper, has a wide distribution range along the shallow Brazilian tropical waters, mainly in bays and estuarine areas. In the western Atlantic there are records of this species from Massachusetts, USA, to Northern Argentina. This species is very common in Brazilian Southeast region where they reach a total body length of 300 mm (MENEZES; FIGUEIREDO, 1980). The length at first spawning for *C. chrysurus* has been registered at 95 mm of total length in Northeast region (CUNHA et al., 2000) and 115 mm of total length in Southeast region in Brazilian coastal waters (MAGRO et al., 2000).

C. chrysurus is here reported as a new host to isopod parasite *L. redmanni* captured in the coastal waters of Ponta Negra, Rio Grande do Norte, Brazil.

Samples of the Atlantic bumper, *C. chrysurus* were netted from various locations in the coastal waters of Ponta Negra, Rio Grande do Norte, located in the Brazilian Northeast region (Fig. 1). The fishes were captured on a monthly basis, with a help of local fishermen using a beach seine, from January to December 2006. To avoid artifacts of beach seine sampling (in which particularly the isopods abandon their original hosts, possibly live crawling onto another live fish); fishes were examined live, within 30 minutes after capture. All fishes captured were weighed to the nearest gram, measured to the nearest millimeter (the body size as the length from tip of snout to the fork of the caudal peduncle) and analyzed the parasitological aspects.

The body surface, buccal cavity and branchial chamber of the each fish were examined for isopod parasites. The parasites were dislodged from their host and preserved directly in a labeled tube with 70% ethyl alcohol. Sex of all parasites were identified, quantified and their sizes measured to the nearest millimeter (length from the tip of head to the end of telson). The sex of the host fishes was determined following Vazzoler (1996).

(*) Paper presented at the 2nd Brazilian Congress of Marine Biology, on 24-28 May. Búzios, RJ, Brazil. 2009.




Fig. 1. Location map showing the study area Ponta Negra, Rio Grande do Norte, Brazil.

The parasitological indices were calculated following Bush et al. (1997). The Chi-square (χ^2) test was applied to verify significant differences in the sexual proportion between all male and female fishes collected and the parasitized fishes. The Spearman's rank correlation test (r_s) was applied to verify the correlation between the fork length of the hosts and the size of the parasites. A significance level of 0.05 was applied in all tests. In the results mean \pm standard deviations (SD) are given.

Material analyzed. A total of 204 samples of *C. chrysurus* were captured and which ranged from 32 to 221mm (118 ± 29) in fork length. The length of the 12 parasitized fishes ranged from 93 to 135 mm, and from this total, 12 specimens of the isopod parasite *L. redmanni* were collected (9 females and 3 males).

Parasitological indices and site of fixation.

L. redmanni showed a prevalence of 5.9%, mean intensity of one parasite per host and a relative abundance of 0.05 parasites per fish captured. The isopods were attached to anterior-ventral portion of the branchial chamber and the heads of the parasites were always directed to the ventral side of the host. Four females of *L. redmanni* were registered carrying eggs and two incidents were registered showing the release

of mancae. Damages to the host *C. chrysurus* were caused by *L. redmanni*, and the most obvious effect caused by parasitism was the atrophy and the complete loss of the gill filaments (Fig. 2).

Host-parasite interaction. The sex ratio of the fishes collected was 2 males: 1 female ($\chi^2 = 12.45$, $df = 1$, $p < 0.05$). The male hosts were more parasitized than the females ($\chi^2 = 22.98$, $df = 1$, $p < 0.05$). The sex ratio of *L. redmanni* was 1 male to 3 females. Body size of the male isopods was 12 ± 1.6 in total length and ranged from 10 to 14 mm. Body size of the female isopods was 13 ± 1.7 and ranged from 11 to 15 mm. The relationship between the total length of the parasite (TL_{par}) and fork length of the hosts (FL_{hos}) showed a significant linear regression: $TL_{par} = 5.0693 + 0.0624 * FL_{hos}$ ($r_s = 0.67$, $p < 0.05$) (Fig. 3).

The results show that in the coastal waters of Ponta Negra, Rio Grande do Norte, Brazil, around 5.9% of *C. chrysurus* were infected by the isopod *L. redmanni* in the branchial chamber.

The length at first maturation for *C. chrysurus* is at 95 mm of total length in the Northeast region (CUNHA et al., 2000). In this study, the fork length of the parasitized fishes ranged from 93 to 135 mm which were all above the length at first maturation registered for this species.




Fig. 2. The marine fish *C. chrysurus* parasitized by *L. redmanni*. A) A female specimen of the parasite in the branchial chamber showing the release of mancae of *L. redmanni* (black spots below of the pectoral fin); B) Damages on the gill filament caused by parasitism; C) A female specimen of *L. redmanni* showing the ventral view marsupium or brood pouch and the dorsal view. Scale bars = 10 mm.




Fig. 3. Linear regression between the body size of the host *C. chrysurus* and the parasite *L. redmanni*.

The incidence and intensity of isopod parasite exhibit considerable variation (RAVICHANDRAN et al., 1999; RAVICHANDRAN et al., 2001; GRUTTER, 2003; CUYAS et al., 2004). BOWMAN (1960) registered the incidence of infection of *Livoneca puhi* on Hawaiian Moray eel, *Gymnothorax eurostus* which ranged from 15 to 17%. Sadzikowski and Wallace (1974) found a prevalence of 1.7% for the isopod *Livoneca ovalis* on white perch, *Morone americana* in the Delaware River. Alas et al. (2008) studying the cymothoid *Nerocila bivittata* on the fish *Parablennius sanguinolentus* (Perciformes, Blenniidae) in the Samsun coast of the Black Sea, found a prevalence of 7.4% and registered *P.*

sanguinolentus as a new host to the *N. bivittata*. However, low prevalence of isopod parasite is more frequently registered (SARTOR, 1986).

The female-male pair of *L. redmanni* occurs in each gill cavity of the hosts *S. regalis* and *S. brasiliensis* (Scombridae) thus showing host specificity (WILLIAMS; BUNKLEY-WILLIAMS, 1996). Lima et al. (2005) registered a prevalence of 86% and found up to four specimens of *L. redmanni* in the branchial chamber of the host *S. brasiliensis* in the coastal waters of the Rio Grande do Norte, Brazil. The present study registers one parasite *L. redmanni* per host *C. chrysurus*. Probably, this low prevalence of *L. redmanni* on the host *C. chrysurus* in relation to

Spanish mackerels can be explained based on the body size. Spanish mackerels (*S. regalis* and *S. brasiliensis*) have bigger body sizes which could afford to carry a heavy load of parasites as compared to the smaller size of *C. chrysurus*.

Males predominated the sexual proportion of the fishes collected in this study. Bello et al. (1997) found that equal numbers of male and female hosts *Atherina boyeri* carry the cymothoid *Mothocya epimerica*. Lima et al. (2005) found a sexual proportion of 1 male:1 female in the population of *S. brasiliensis* in the coastal waters of the Rio Grande do Norte and no relationship between *L. redmanni* and the sex of *S. brasiliensis* was established. Probably the preference of *L. redmanni* for male hosts of *C. chrysurus* is due to the larger proportion of the male hosts and a higher proportion of female parasites.

Isopods inhabiting the branchial chamber inflict damage to gills through attachment and feeding and the extent of damage is directly proportional to the size of the parasite and duration of settlement (ROMESTAND; TRILLES, 1977; RAVICHANDRAN et al., 2007). A positive and significant correlation was found between the size of both sexes of *L. redmanni* and the body size of the host *C. chrysurus*. Colorni et al. (1997) observed that the size of the females and males of *Livoneca* sp. correlated positively with the body size of the host Red Sea silverside, *Atherinomorus lacunosus* in the Gulf of Eilat. The isopods while in the manca stage usually locate and attaches to the hosts, particularly to young hosts (MENZIES et al., 1955; KROGER; GUTHRIE, 1972; GARREY; MAXWELL, 1982; ADLARD; LESTER, 1995; MARKS et al., 1996; COLORNI et al., 1997; LEONARDOS; TRILLES, 2003). In the definitive host, isopod growth is correlated to the host growth (FOGELMAN; GRUTTER, 2008). Leonardos and Trilles (2003) suggested two strategies in attempt to explain a linear correlation between the host and isopod parasite size. In the first strategy, infections occur early in the life of the fish and then the parasite grows with the host. In the second strategy, the isopod grows rapidly after the infection to a maximum size and then inhibits further growth due to the restricted space of the hosts branchial cavity. The first strategy is more coherent with the results of the present work. Furthermore, the body size affects survival probabilities, reproductive output and individual output of many organisms (TWOMBLY; TISH, 2000). The synchrony of parasite and host growth seems to be a natural strategy that increase its own body size and reproductive output (ÁLVAREZ; FLORES, 1997; Leonardos & Trilles, 2003; Chavez-Lopez et al., 2005).

The present study reports for the first time *C. chrysurus* as a host of the isopod parasite *L* in Brazilian coastal waters. Infections by *L. redmanni* on

the host *C. chrysurus* occur and the parasite grows with the host and enters in the reproductive phase at a latter stage.

ACKNOWLEDGEMENTS

The present study was carried out with financial support from the National Council for Scientific and Technological Development of Brazil (CNPq).

REFERENCES

- ADLARD, R. D.; LESTER, R. J. G. The Life-cycle and biology of *Anilocra pomacentri* (Isopoda, Cymothoidae), an ectoparasitic isopod of the coral-reef fish, *Chromis nitida* (Perciformes, Pomacentridae). **Aust. J. Zool.**, v. 43, p. 271-281, 1995.
- ALAS, A.; ÖKTENER, A.; ISCIMEN, A.; TRILLES, J. P. New host record, *Parablennius sanguinolentus* (Teleostei, Perciformes, Blenniidae) for *Nerocila bivittata* (Crustacea, Isopoda, Cymothoidae). **Parasitol. Res.**, v.102, p.645-646, 2008.
- Alvarez, F.; Flores, M. *Cymothoa exigua* (Isopoda: Cymothoidae) parasitando al pargo *Lutjanus peru* (Pisces: Lutjanidae) em Manzanillo, Colima, México. **Rev. Biol. Trop.**, v. 44, p.391-394, 1997.
- BELLO, G.; VAGLIO, A.; PISCITELLI, G. The reproductive cycle of *Mothocya epimerica* (Isopoda: Cymothoidae) a parasite of sand smelt *Atherina boyeri* (Osteichthyes: Atherinidae), in the Lesina Lagoon, Italy. **J. nat. Hist.**, v. 92, n.3, p. 1055-1066, 1997.
- BOWMAN, T. E. Description and notes on the biology of *Lironeca puhi*, n. sp. (Isopoda, Cymothoidae) parasite of the Hawaiian moray eel *Gymnothorax eurostus* (Abbott). **Crustaceana**, v.1, p.82-91, 1960
- BRUSCA, R. C. A monograph on the Isopoda: Cymothoidae (Crustacea) of the eastern Pacific. **Zool. J. Linn. Soc.**, v. 73, p.117-199, 1981.
- BULLAR, J. F. The generative organs of parasitic isopoda. **J. Anat. Physiol.**, v. 11, p.118-128, 1876.
- BUNKLEY-WILLIAMS, L; WILLIAMS JR., E. H. Brood pouch release of *Anilocra chromis* Williams & Williams (Isopoda, Cymothoidae) a parasite of brown chromis, *Chromis multilineatus* (Guichenot) in the Caribbean. **Crustaceana**, v. 49, n.1, 1985.
- BUNKLEY-WILLIAMS, L.; WILLIAMS JR., E. H. Isopods associated with fishes: a synopsis and corrections. **J. Parasitol.**, v.84, p.893-896, 1998.
- BUSH, A. O.; LAFFERTY, K. D.; LOTZ, J. M.; SHOSTAK, A.W. Parasitology meets ecology on its own terms: Margolis et al. revisited. **J. Parasitol.**, v. 83, p.575-583, 1997.
- CHAVEZ-LOPEZ, R.; ROCHA-RAMÍREZ, A.; ÁLVAREZ, F.; WETZER, R. *Elthusa alvaradoensis* Rocha-Ramirez, Chavez-Lopez & Bruce, 2005 (Isopoda, Cymothoidae) parasitizing the inshore lizardfish, *Synodus foetens* (Linnaeus, 1766) on the continental shelf off central Veracruz, Mexico. **Crustaceana**, v. 78, p. 865-872, 2005.

- COLORNI, A.; TRILLES, J. P.; GOLANI, D. *Livoneca* sp. (Flabellifera: Cymothoidae), an isopod parasite in the oral and branchial cavities of the Red Sea silverside *Atherinomorus lacunosus* (Perciformes, Atherinidae). **Dis. aquat. Org.**, v. 31, p.65-71, 1997.
- CUNHA, F. E. A.; FREITAS, J. E. P.; FITOSA, C. V.; MONTEIRO-NETO, C. *Biologia e biometria da palombeta, Chloroscombrus chrysurus* (Linnaeus, 1766) (Teleostei:Carangidae), em Fortaleza, Ceará, Brasil. **Arq.Ciê.Mar.**, v. 33, p.143-148, 2000.
- CUYAS, C.; CASTRO, J. J.; ORTEGA, S. A. T.; CARBONELL, E. Insular stock identification of *Serranus atricauda* (Pisces: Serranidae) through the presence of *Ceratothoa steindachneri* (Isopoda: Cymothoidae) and *Pentacapsula cutanea* (Myoxoa: Pentacapsulidae) in the Canary Islands. **Sci. Mar.**, v. 68, p. 159-163, 2004.
- FOGELMAN, R. M.; GRUTTER, A. S. Manceae of the parasitic cymothoid isopod, *Anilocra apogonae*: early life history, host-specificity, and effect on growth and survival of preferred young cardinal fishes. **Coral Reefs**, v.27, p.685-693, 2008.
- GARREY, J.; MAXWELL, H. Infestation of the jack mackerel, *Trachurus declivis* (Jenyns), with the cymothoid isopod, *Ceratothoa imbricatus* (Fabricus), in south eastern Australian waters. **J. Fish. Biol.**, v.20, p.341-349, 1982.
- GRABDA, J. Marine fish parasitology. In: GRABDA, E. (Ed.) **Marine fish pathology**. Warsaw: PWN- Polish Scientific Publ., 1991, p. 222-227.
- GRUTTER, A. S. Feeding ecology of the fish ectoparasite *Gnathia* sp. (Crustacea:Isopoda) from the Great Barrier Reef and its implications for fish cleaning behavior. **Mar. Ecol. Prog. Ser.**, v. 259, p. 295-302, 2003.
- KROGER, R. L.; GUTHRIE, J. F. Incidence of parasitic isopod, *Olencira praegustator*, in juvenile Atlantic menhaden. **Copeia**, v. 2, p.370-374, 1972.
- LANZING, W. J. R.; O'CONNOR, P. F. Infestation of luderick (*Girella tricuspidata*) populations with isopods. **Aust. J. mar. Freshw. Res.**, v. 26, p.355-361, 1975.
- LEONARDOS, I.; TRILLES, J. P. Host-parasite relationships: occurrence and effect of the parasitic isopod *Mothocya epimerica* on sand smelt *Atherina boyeri* in the Mesolongi and Etolikon Lagoons (W. Greece). **Dis. aquat. Org.**, v.54, p.243-251, 2003.
- LESTER, R. J. G.; ROUBAL, F. R. Phylum Arthropoda. Isopoda: Cymothoidae. WOO, P. (Ed.). **Fish diseases and disorders**. Wallingford: CAB International, 1995, p. 550-561.
- LESTER, R. J. G.; 2005ROUBAL, F. R. Isopod. In: ROHDE, K. (Ed.). **Marine Parasitology**. Collingwood: CSIRO Publishing, 2005. p. 138-144.
- LIMA, J. T. A. X.; CHELLAPPA, S.; THATCHER, V. E. *Livoneca redmanni* Leach (Isopoda, Cymothoidae) e *Rocinela signata* Schioedtei & Meinert (Isópoda, Aegidae), ectoparasitos de *Scomberomorus brasiliensis* Collette, Russo & Zavala-Camin (Osteichthyes, Scombridae) no Rio Grande do Norte. **Rev. Bras. Zool.**, v. 22, p.1104-1108, 2005.
- MAGRO, M.; CERGOLÉ, M. C.; ROSSI-WONGTSCHOWSKI, C. L. D. B. **Síntese do conhecimento dos principais recursos pesqueiros costeiros potencialmente explotáveis na costa Sudeste Sul do Brasil: peixes**. Rio de Janeiro: Graffline Editora, 2000. 143p.
- MARKS, R. E.; JUANES, F.; HARE, J. A.; CONOVER, D.O. Occurrence and effect of the parasitic isopod, *Livoneca ovalis* (Isopoda: Cymothoidae), on young-of-the-year bluefish, *Pomatomus saltatrix* (Pisces: Pomacentridae). **Can. J. Fish aquat. Sci.**, v.53, p. 2052-2057, 1996.
- MENEZES, N. A.; FIGUEIREDO, J. L. **Manual dos Peixes Marinhos do Sudeste do Brasil IV. Teleostei (3)**. São Paulo: USP, Museu de Zoologia, 1980. 96 p.
- MENZIES, R. J.; BOWMAN, T. E.; FRANKLIN, A. G. Studies of the biology of the fish parasite *Livoneca convexa* Richardson (Crustacea, Isopoda, Cymothoidae). **Wassman J. Biol.**, v. 13, p.277-295, 1955.
- RAMDANE, Z.; BENSOUILAH, M. A.; TRILLES, J. P. The Cymothoidae on marine fishes, from Algerian fauna. **Belg. J. Zool.**, v. 137, n.1, p.67-74, 2007.
- RAVICHANDRAN, S.; SINGH, A. J. A. R.; VEENAPPAN, N.; KANNUPANDI, T. Effect of isopod parasite *Joryma brachysoma* on *Illisha melastoma* from Parangipettai coastal waters (southeast coast of India). **Ecol. Environ. Conser.**, v.5, p. 95-101, 1999.
- RAVICHANDRAN, S.; SINGH, A. J. A. R.; VEENAPPAN, N. Parasite induced vibriosis in *Chirocentrus dorab* off Parangipettai coastal waters. **Curr. Sci.**, v. 80, p. 101-102, 2001.
- RAVICHANDRAN, S.; BALASUBRAMANIN, T.; KANNUPANDI, T. Incidence of parasitic isopods on the fish *Sphyrna obtusata*. **Res. J. Parasitol.**, v. 2, p.45-50, 2007.
- ROMESTAND, B.; TRILLES, J. P. Influence of Cymothoidae (Crustacea, Isopoda, Flabellifera) on some hematological constants of host-fishes. **Z. Parasitenkd.**, v. 52, n.1, p.91-95, 1977.
- SADZIKOSWSKI, M. R.; WALLACE, D. C. The incidence of *Livoneca ovalis* (Say) (Crustacea, Isopoda) and its effect on the Growth of white perch, *Morone Americana* (Gmelin), in the Delaware River near Artificial Island. **Chesapeake Sci.**, v.15, p.163-165, 1974.
- SARTOR, S. M. Incidência de isópodos parasitas (Cymothoidae) em peixes da plataforma continental brasileira. **Bolm Inst. oceanogr.**, v.34, p.1-12, 1986.
- TWOMBLY, S.; TISH, N. Body size regulation in copepod crustaceans. **Oecologia**, v. 122, p. 318-326, 2000.
- VAZZOLER, A. E. A. M. **Biologia reprodutiva de peixes Teleósteos: teoria e prática**. Maringá: EDUEM, 1996. 169 p.
- WILLIAMS JR., E. H. ; BUNKLEY-WILLIAMS, L. **Parasites off shore, big game sport fishes of Puerto Rico and the Western North Atlantic**. San Juan PR: Puerto Rico Department of Natural and Environmental Resources; Mayaguez: University of Puerto Rico, 1996. 382 p.

(Manuscript received 04 June 2009; revised 16 November 2009; accepted 27 April 2010)