

Brazilian Journal of Veterinary Research and Animal Science (2003) 40:341-348
ISSN printed: 1413-9596
ISSN on-line: 1678-4456

Development of a polymerase chain reaction and its comparison with agar gel immunodiffusion test in the detection of bovine leukemia virus infection

Desenvolvimento de uma reação em cadeia pela polimerase e comparação com a imunodifusão em gel de agar na detecção de infecções pelo vírus da leucemia bovina

Marcelo Fernandes CAMARGOS¹;
Daniel STANCEK¹;
Leandro Moreira LESSA¹;
Jenner Karlisson Pimenta REIS²;
Maurílio Andrade ROCHA¹;
Rômulo Cerqueira LEITE¹

1- Escola de Veterinária da Universidade Federal de Minas Gerais, Belo Horizonte – MG
2- Colégio Técnico da Universidade Federal de Minas Gerais, Belo Horizonte – MG

Correspondence to:

MARCELO FERNANDES CAMARGOS
Departamento de Medicina Veterinária Preventiva
Escola de Veterinária da UFMG
Avenida Antônio Carlos, 6627
30270-010 - Belo Horizonte – MG
e-mail: mcamargo@cyberpl.com.br

Recebido para publicação: 01/07/2002
Aprovado para publicação: 03/06/2003

Summary

Polymerase chain reaction (PCR) was used for bovine leukemia virus (BLV) detection in the peripheral leukocytes of the infected bovines. The primers used were designed to amplify a part of *env* gene of BLV. PCR products were analyzed by agarose gel electrophoresis stained by ethidium bromide. The analytical specificity of PCR was confirmed by enzymatic restriction analysis of the PCR product with Bam HI and also by nucleotide sequence analysis of three PCR samples. Sixty five animals were tested for anti-BLV antibody, by agar gel-immunodiffusion test (AGID) and for direct BLV detection by PCR. There was a 73.80% concordance rate between the two tests. Four animals positive in AGID were PCR negative, while 13 AGID negative animals were found PCR positive. PCR got a 0.87 diagnosis sensitivity and 0.62 specificity. The developed PCR may be complementary tool in the diagnosis of BLV infection, but should have its diagnosis sensitivity improved.

Key-words

Immunodiffusion.
Bovine leukemia virus.
Gene.
Polymerase chain reaction.

Introduction

The bovine leukemia virus (BLV), a Deltaretrovirus has a worldwide distribution in cattle herds and causes a disease known as Enzootic Bovine Leukosis (EBL). BLV infected animals may suffer lymphosarcoma, a fatal condition, bovine impairment in milk production and impairment in reproductive efficiency besides commercial restrictions^{1,2,3,4,5}.

The virus is able to infect, *in vitro*, cells of different species including human cells⁶ and it can also infect, *in vivo*, sheep, rabbits and non human primates⁶. It primarily infects bovine B lymphocytes but it is able to infect T lymphocytes and neutrophils. EBL is considered an experimental model for studies of human leukemias, caused by the Human T-Cell Leukemia virus types 1 and 2 (HTLV-1/HTLV-2, due to HTLV/BLV similarities in genome and

in pathogenesis⁷.

Various indirect and direct methods have been used for BLV detection, e.g. AGID, ELISA and PCR^{8,9,10,11,12,13}. In some cases the indirect methods fail in the detection of BLV infections^{9,14,15,16}. The goal of this work was to develop a PCR for BLV detection in blood samples obtained from cattle, to determine the specificity and sensitivity, and to compare the results of PCR and AGID, this one as the gold standard, applied in a BLV naturally infected dairy herd.

Material and Method

Blood samples

Blood was collected in order to have at least 30 samples from both EBL positive and negative animals. Sixty five samples were collected in tubes with and without EDTA. The farm, where the bovines (Holstein black and white) came from, was localized in the metropolitan region of Belo Horizonte, Minas Gerais State, in Southeast region of Brazil. All tested animals were in good health and their age ranged from 1 to 10 years.

Antibody detection

AGID test was used for anti-BLV antibody detection utilizing p24 and gp51 antigens commercially produced (EMBRAPA/CNPSA, Brazil), as described by Miller and Van der Maaten¹⁷.

DNA extraction

DNA extracted from fetal lamb kidney cells (FLK) persistently infected with BLV were used as PCR positive control¹². DNA extracted from MDBK non-infected cells (Marvin-Derby Bovine Kidney) was used to

verify the occurrence of annealing of the primers with bovine DNA. The controls DNA and the DNA from the infected animals were extracted from whole blood samples by the GFX Genomic Blood DNA Purification Kit (Amersham Pharmacia Biotech).

PCR primers

Primers were constructed using Oligo Primer Analysis program¹⁹. They were selected to amplify 521 base pairs (bp) fragments within the gp51 *env* gene of BLV. They are represented below, the number refers to the sequence accession number K02120:

Forward 5' -
GGCCATGGTCACATATGATTG-
3' (genome position 5128-5149)

Reverse 5' -
CGTTGCCTTGAGAAACATTGAAC-
3' (genome position 5627-5649)

PCR amplification

PCR was prepared in a 20 ml total volume, which contained 10 mM TrisHCL solution, pH 8.3; 50 mM KCl; 1.5 mM MgCl₂; 1,00% glycerol (v/v); 1,00% (v/v) dimetilsulfoxide (DMSO); 10 pmoles of each primer; 0.2 mM of each deoxinucleotide (Promega, USA); 0.5 Taq DNA polymerase (Gibco-BRL, USA) and 1 il of DNA preparation containing approximately 150 ng of DNA. One drop of mineral oil was then added to each tube. Amplification reaction was carried out in a PCR programmed thermocycler (MJ Research, Inc., USA), in the following cycles: initial incubation at 95° C 3 min., followed by repeated 35 cycles: denaturation 95° C 60 sec., annealing 59° C 60 sec. and extension by polymerase at 72° C 60 sec. The last cycle was run at 72° C for 10 min. Amplification products was analyzed in 1,00% agarose gels stained with ethidium bromide.

Estimation of PCR detection limit

One sample of DNA obtained from BLV seropositive bovine was diluted and DNA concentration was estimated spectrophotometrically (UV-160 A – Shimadzu spectrophotometer) at the waves length 260 nm. Tenfold DNA dilutions were prepared and then a PCR was ran to verify at which input DNA could be amplified.

Estimation of PCR analytical specificity

PCR specificity was evaluated by restriction-enzyme analysis and by nucleotide sequence analysis of the amplified fragments of three samples. The PCR products were purified by Wizard PCR Preps DNA Purification System Kit (Promega, USA) according to manufacturer recommendations. The samples were digested with Bam HI restriction enzyme (Promega, USA), as recommended by the manufacturer, by 60 min. incubation at 37° C. The obtained products were analyzed by electrophoreses in 1.00% agarose gels (w/v) stained by ethidium bromide. Sequential analysis was carried out in an automatic sequence analyzer (ALF, Pharmacia, Sweden) at Bioagro-UFV, Viçosa, Minas Gerais, Brazil.

Statistical evaluation

Association between PCR and AGID was statistically estimated by the kappa test (k)²⁰. The PCR diagnosis sensitivity and specificity were calculated using AGID as the gold standard²⁰.

Results

Detection of proviral DNA in whole blood of infected animals

The PCR described here, utilizing primers for BLV-*env*, gene

which coding for gp51 *env* protein, amplified a fragment of approximately 521 bp. The chosen sequence was localized in a rather conserved region of *env* gene and, additionally, primers annealed at quite high temperature which increased the specificity of BLV *env* gene amplification. Fragments lacking specificity were not detected. Figure 1 shows PCR results obtained with DNA extracted from bovine blood samples.

PCR detection limit

One DNA sample had its concentration estimated in 120 ng/il. This sample was diluted in deionized water by ten fold dilutions and analyzed by PCR at the same conditions previously described. Visible amplification was seen up to the dilution 10^{-3} (Figure 2) which corresponded to 120 pg of bovine DNA/il meaning that this is the minimal concentration of DNA able to give a positive result in our PCR test. However, we have analyzed just one sample and therefore it is not possible to generalize this result.

Determination of PCR analytical specificity

Specificity of the amplified product was confirmed by enzymatic restriction analysis with Bam HI and also by sequence analysis of three samples. The enzymatic restriction produced, as expected, two fragments with 422 bp and 99 bp respectively (Figure 3). Besides, no amplification could be seen when DNA isolated from MDBK was used which proves that the primers did not anneal with bovine genome (Figure 1). Sequence analysis showed that the amplified fragments correspond to the BLV *env* gene. The obtained sequences were deposited in GenBank, USA <http://www.ncbi.nlm.nih.gov:80/Genbank/>




Figure 1
 PCR results. The position of 521 bp band is pointed by the arrow. Lane 1: positive control, lane 2: negative control; lane 3: reagents control; lanes 4 and 9: BLV DNA negative samples; lanes 6, 7, 8, 10 and 11: BLV DNA positive samples; lane 5: molecular length markers (fx174, Life/USA). The corresponding molecular length is given on the right side of the figure




Figure 2
 Determination of detection limits of PCR. The position of 521 bp band is showed by the arrow. Lane 1: reagents control; lane 2: 120 ng/ul tube (estimated concentration); lane 3: 120.10⁻² ng/ul tube; lane 4: 120.10⁻³ ng/ul tube; lane 5: 120.10⁻⁴ ng/ul tube; lane 6: molecular length marker (fx174); lane 7: 120.10⁻⁵ ng/ul tube; lane 8: 120.10⁻⁶ ng/ul tube; lane 9: 120.10⁻⁷ ng/ul tube; lane 10: 120.10⁻⁸ ng/ul tube and lane 11: 120.10⁻⁹ ng/ul tube




Figure 3
 Determination of PCR analytical specificity by restriction enzyme analysis. The position of 521 bp band is pointed out by the arrow. Lane 1: purified BLV *env* gene fragment; lane 2: fragments obtained after digestion with the restriction enzyme BamHI; lane 3: molecular length marker (100 bp DNA Ladder, Gibco-BRL/USA)

[index.html](#)) and can be reached by the following access numbers: AF399702, AF399703 and AF399704. These were the first submission of BLV *env* sequences from Brazilian cattle.

Statistical evaluation

The results are shown in Table

Table 1

Comparison between PCR and AGID tests in BLV detection

	AGID positive	AGID negative	Total
PCR positive	27	13	40
PCR negative	4	21	25
Total	31	34	65

1. From 65 tested samples, 31 were positive in AGID test while 4 of these samples were PCR negative. On the other hand, 13 out of 34 negative samples in AGID test were positive by PCR. Showing a 73.80 % agreement between the two tests. The statistical kappa value (k) was 0.48 which reflects a moderate alignment between PCR and AGID. The PCR diagnosis sensitivity was 0.87 and specificity 0.62.

Discussion

The PCR developed for partial amplification of BLV *env* gene able to detect BLV in blood samples of infected cattle. Specific fragments were obtained as confirmed by BamHI enzyme digestion and nucleotide sequence analysis. It was designed as a supplementary test to be used mainly in herds at final phase of a EBL eradication program or when unclear results are present in serologic tests. This is the first study evaluating the usefulness of a PCR for field diagnosis of EBL on a naturally infected Brazilian herd.

So far, the PCRs described in literature, that could detect small quantities of proviral DNA, were obtained by PCR associated with hybridization, reamplification or semi-nested procedures^{10,11,21,22,23,24}.

We found a moderate 73.80% agreement between PCR and AGID tests, although up to 100.00% association between these two tests has

been described by other authors^{7,11,25,26,27}. We believe this high agreement may be due, however, to a small number of the samples utilized or due to the higher sensitivity of those PCRs. The PCR diagnosis sensitivity should be improved, maybe by the use of nested primers and/or improving in proviral DNA purification. The low diagnosis specificity reached it is not because of false positive results in PCR, but it is probably due to detection of recently infected animals by PCR, that AGID could not detect.

Higher BLV detection rate by PCR in blood screening tests is generally due to its higher sensitivity when compared with serologic tests. Early stages of BLV infection with no anti-BLV antibodies production, chronic BLV infections without or with low anti-BLV antibody levels as well as lower AGID sensitivity are further factors regarded as influencing the outcome of the PCR/AGID screening results^{9,14,15}. Roberts et al¹⁶ observed that animals persistently infected with the bovine diarrhea virus (BVD), presented an impairment in their immune response to BLV. Genetic variants of the BLV *env* gene can also influence the outcome of PCR/AGID testing by exhibiting atypical forms of BLV infection with decreased immune response²⁸. Our lab have been working in cloning and sequencing Brazilian isolates of BLV to figure out the main variation in *env* gene. This will give us the possibility of constructing primers more specific in conserved regions of the genome besides mapping *env* from different regions.

The findings of PCR negative, AGID positive samples could be explained by the presence of Taq DNA polymerase inhibitors in blood samples²⁹, by absence of BLV in circulating lymphocytes^{9,24,30}, small number of

provirus copies³¹ or restricted infection to lymphoid organs³².

Some authors showed large variations among PCR results utilizing samples collected from different geographical regions, which was probably due to non annealing of primers caused by genetic variations of the BLV provirus^{33,34}. Mamoun et al.³⁵, compared nucleotide sequences of BLV *env* gene isolated from different geographical regions and found a 3.50% rate of nucleotide variation among those isolates, with the main part being located between nucleotides 4945 and 5111 of BLV *env* gene. For this reason the primers used in this work were chosen from a more conserved *env* gene region as previously described^{35,36,37}.

On large scale BLV infection screenings, when PCR negative/AGID positive samples might be found. It is recommended to use the PCR as a complementary test. PCR will be also very important to screen AGID or ELISA negative samples as well as to determine BLV infections calves that received colostrum from BLV seropositive cows. Results of PCR are available within one day while for AGID tests three days are usually needed. Recent BLV infections can be detected by PCR into 2–4 weeks earlier than by AGID test^{38,39}. PCR should be used in BLV eradication programs, mainly in those herds with a low number of infected animals. Being also a reliable method when checking animals for export or selecting BLV free animals for vaccine preparations (e.g., anaplasmosis, babesiosis).

This work was supported by the FAPEMIG grant No. CAG-1084/98; CNPq fellowship No. 301404/88-7, CAPES fellowship, FEP/MVZ additional support.

Resumo

A reação em cadeia pela polimerase (PCR) foi utilizada para a detecção do vírus da leucemia bovina (VLB) em leucócitos periféricos de bovinos infectados. Os iniciadores utilizados foram construídos para amplificar uma parte do gene *env* do VLB. Os produtos da PCR foram analisados por eletroforese em gel de agarose corados por brometo de etídeo. A especificidade analítica da PCR foi confirmada por restrição enzimática dos produtos da reação com Bam HI e também pela análise da seqüência de três amostras. Sessenta e cinco animais foram testados para a presença de anticorpos anti-VLB, pela imunodifusão em gel de agar (IDGA) e pela PCR, para detecção direta do VLB. Houve 73,80% de concordância entre os dois testes. Quatro animais positivos na IDGA foram PCR negativos, enquanto 13 animais negativos na IDGA foram positivos na PCR. A sensibilidade diagnóstica obtida foi de 0,87 e a especificidade diagnóstica 0,62. A PCR desenvolvida pode ser uma ferramenta complementar no diagnóstico de infecções causadas pelo VLB, mas deve ter sua sensibilidade diagnóstica melhorada.

Palavras-chave

Imunodifusão.
Vírus da leucemia bovina.
Genes.
Reação em cadeia pela polimerase.

References

- 1- BRENNER, J.; VAN-HAAM, M.; SAVIR, D.; TRAININ, Z. The implication of BLV infection in the productivity, reproductive capacity and survival rate of a dairy cow. **Veterinary Immunology Immunopathology**, v. 22, n. 3, p. 299-305, 1989.
- 2- DA, Y.; SHANKS, R. D.; STEWART, J. A.; LEWIN, H. A. Milk and fat yields decline in bovine leukemia virus-infected Holstein cattle with persistent lymphocytosis. **Proceedings National Academy Sciences USA**, v. 90, n. 14, p. 6538-6541, 1993.
- 3- D'ANGELINO, J. L.; GARCIA, M.; BIRGEL, E. H. Productive and reproductive performance in cattle infected with bovine leukosis virus. **Journal Dairy Research**, v. 65, n. 4, p. 693-695, 1998.
- 4- HEALD, M. T. S.; WALTNER-TOEWS, D.; JACOBS, R. M.; MCNAB, W. B. The prevalence of anti-bovine leukemia virus antibodies in dairy cows and associations with farm management practices, production and culling in Ontario. **Preventive Veterinary Medicine**, v. 14, n. 1-2, p. 45-55, 1992.
- 5- REINHARDT, G.; HOCHSTEIN-MINTZEL, V.; RIEDMANN, S.; LEAL, H.; NIEDDA, M. Estudio serológico de Leucosis Enzoótica Bovina en un predio de la provincia de Valdivia y su relación a parámetros productivos y reproductivos. **Journal Veterinary Medicine B**, v. 35, n. 3, p. 178-185, 1988.
- 6- BURNY, A. Bovine leukaemia: facts and hypotheses derived from the study of an infectious cancer. **Veterinary Microbiology**, v. 17, n. 3, p. 197-218, 1988.
- 7- SHERMAN, M. P.; EHRlich, G. D.; FERRER, J. F.; SNINSKY, J. J.; ZANDOMENI, R.; DOCK, N. L.; POIESZ, B. J. Amplification and analysis of specific DNA and RNA sequences of bovine leukemia virus from infected cows by polymerase chain reaction. **Journal Clinical Microbiology**, v. 30, n. 1, p. 185-191, 1992.
- 8- BALLAGI-PORDANY, A.; BELAK, S. The use of mimics as internal standards to avoid false negatives in diagnostic PCR. **Molecular Cell Probes**, v. 10, n. 3, p. 159-164, 1996.
- 9- EAVES, F. W.; MOLLOY, J. B.; DIMMOCK, C. K.; EAVES, L. E. A field evaluation of the polymerase chain reaction procedure for the detection of bovine leukemia virus proviral DNA in cattle. **Veterinary Microbiology**, v. 39, n. 3-4, p. 313-321, 1994.
- 10- JACOBS, R. M.; SONG, Z.; POON, H.; HEENEY, J. L.; TAYLOR, J. A.; JEFFERSON, B.; VERNAU, W.; VALLI, V. E. O. Proviral detection and serology in bovine leukemia virus-exposed normal cattle and cattle with lymphoma. **Canadian Journal Veterinary Research**, v. 56, n. 4, p. 339-348, 1992.
- 11- NAIF, H. M.; BRANDON, R. B.; DANIEL, R. C.; LAVIN, M. F. Bovine leukaemia proviral DNA detection in cattle using the polymerase chain reaction. **Veterinary Microbiology**, v. 25, n. 2-3, p. 117-129, 1990.
- 12- ROLA, M.; KUZMAK, J. The detection of bovine leukemia virus proviral DNA by PCR – ELISA. **Journal Virological Methods**, v. 99, n. 1-2, p. 33-40, 2002.
- 13- SIMARD, C.; RICHARDSON, S.; DIXON, P.; BELANGER, C.; MAXWELL, P. Enzyme-linked immunosorbent assay for the diagnosis of bovine leukosis: comparison with agar gel immunodiffusion test approved by the

- Canadian Food Inspection Agency. **Canadian Journal Veterinary Research**, v. 64, n. 2, p. 101-106, 2000.
- 14- COCKERELL, G. L.; ROVNAK, J. The correlation between the direct and indirect detection of bovine leukemia virus infection in cattle. **Leukemia Research**, v. 12, n. 6, p. 465-469, 1988.
- 15- DE BOER, G. F.; BOERRIGTER, H. M.; GROEN, J.; OSTERHAUS, A. D. M. E. Identification of bovine leukemia virus (BLV) infected cattle by complex-trapping-blocking (CTB) ELISA employing monoclonal antibodies directed against BLV-p24. **Journal Veterinary Medicine B**, v. 34, p. 717-728, 1987.
- 16- ROBERTS, D. H.; LUCAS, M. H.; WIBBERLEY, G.; WESTCOTT, D. Response of cattle persistently infected with bovine virus diarrhoea virus to bovine leukosis virus. **Veterinary Record**, v. 122, n. 13, p. 293-296, 1988.
- 17- MILLER, J. M.; VAN DER MAATEN, M. J. Use of glycoprotein antigen in the immunodiffusion test for bovine leukemia virus antibodies. **European Journal Cancer**, v. 13, n. 12, p. 1369-1375, 1977.
- 18- VAN DER MAATEN, M. J.; MILLER, J. M. Replication of bovine leukemia virus in monolayer cell cultures. **Bibliotheca Haematologica**, v. 43, p. 360-362, 1975.
- 19- RYCHLIK, W.; RHOADS, R. E. A computer program for choosing optimal oligonucleotides for filter hybridization, sequencing and in vitro amplification of DNA. **Nucleic Acids Research**, v. 17, n. 21, p. 8543-8551, 1989.
- 20- SMITH, R. D. **Veterinary clinical epidemiology**. London: CRC Press, 1995, 279 p.
- 21- AGRESTI, A.; PONTI, W.; ROCCHI, M.; MENEVERI, R.; MAROZZI, A.; CAVALLERI, D.; PERI, E.; POLI, G.; GINELLI, E. Use of polymerase chain reaction to diagnose bovine leukemia virus infection in calves at birth. **American Journal Veterinary Research**, v. 54, n. 3, p. 373-378, 1993.
- 22- BRANDON, R. B.; NAIF, H.; DANIEL, R. C. W.; LAVIN, M. F. Early detection of bovine leukosis virus DNA in infected sheep using the polymerase chain reaction. **Research Veterinary Science**, v. 50, n. 1, p. 89-94, 1991.
- 23- MIRSKY, M. L.; DA, Y.; LEWIN, H. A. Detection of bovine leukemia virus proviral DNA in individual cells. **PCR Methods Applications**, v. 2, n. 4, p. 333-340, 1993.
- 24- MURTAUGH, M. P.; LIN, G. F.; HAGGARD, D. L.; WEBER, A. F.; MEISKE, J. C. Detection of bovine leukemia virus in cattle by the polymerase chain reaction. **Journal Virological Methods**, v. 33, n. 1-2, p. 73-85, 1991.
- 25- BALLAGI-PORDANY, A.; KLINTEVALL, K.; MERZA, M.; KLINGEBORN, B.; BELAK, S. Direct detection of bovine leukemia virus infection: practical applicability of a double polymerase chain reaction. **Zentralblatt Veterinarmed [B]**, v. 39, n. 1, p. 69-77, 1992.
- 26- DUBE, S.; BACHMAN, S.; SPICER, T.; LOVE, J.; CHOI, D.; ESTEBAN, E.; FERRER, J. F.; POIESZ, B. J. Degenerate and specific PCR assays for the detection of bovine leukaemia virus and primate T cell leukaemia/lymphoma virus *pol* DNA and RNA: phylogenetic comparisons of amplified sequences from cattle and primates from around the world. **Journal General Virology**, v. 78, n. 6, p. 1389-1398, 1997.
- 27- KUZMAK, J.; SKORUPSKA, A.; MOUSSA, A.; GRUNDBOECK, J. Application of non-radioactive method of DNA detection in the diagnosis of bovine leukemia virus infection. **Bulletin Veterinary Institute Pulawy**, v. 37, n. 1, p. 3-8, 1993.
- 28- FECHNER, H.; BLANKENSTEIN, P.; LOOMAN, A. C.; ELWERT, J.; GEUE, L.; ALBRECHT, C.; KURG, A.; BEIER, D.; MARQUARDT, O.; EBNER, D. Provirus variants of the bovine leukemia virus and their relation to the serological status of naturally infected cattle. **Virology**, v. 237, n. 2, p. 261-269, 1997.
- 29- PANNACIO, M.; LEW, A. PCR based diagnosis in the presence of 8%(v/v) blood. **Nucleic Acids Research**, v. 19, n. 5, p. 1151, 1991.
- 30- REICHEL, M. P.; THAM, K. M.; BARNES, S.; KITTELBERGER, R. Evaluation of alternative methods for the detection of bovine leukaemia virus in cattle. **New Zealand Veterinary Journal**, v. 46, n. 4, p. 140-6, 1998.
- 31- MOLLOY, J. B.; WALKER, P. J.; BALDOCK, F. C.; RODWELL, B. J.; COWLEY, J. A. Control of bovine leukaemia virus transmission by selective culling of infected cattle on the basis of viral antigen expression in lymphocyte cultures. **Veterinary Microbiology**, v. 39, n. 3-4, p. 323-333, 1994.
- 32- KLINTEVALL, K.; BALLAGI-PORDÁNY, A.; NÄSLUND, K.; BELÁK, S. Bovine leukaemia virus: rapid detection of proviral DNA by nested PCR in blood and organs of experimentally infected calves. **Veterinary Microbiology**, v. 42, n. 2-3, p. 191-204, 1994.
- 33- FECHNER, H.; KURG, A.; BLANKENSTEIN, P.; MEWES, G.; GEUE, L.; ALBRECHT, C.; EBNER, D. Direct use of cell lysates in PCR-based diagnosis of bovine leukemia virus infection. **Berliner Munchener Tierärztliche Wochenschrift**, v. 109, n. 11-12, p. 446-450, 1996.
- 34- MARSOLAIS, G.; DUBUC, R.; BERGERON, J.; MORREY, J. D.; KELLY, E. J.; JACKSON, M. K. Importance of primer selection in the

- application of PCR technology to the diagnosis of bovine leukemia virus. **Journal Veterinary Diagnostic Investigation**, v. 6, n. 3, p. 297-301, 1994.
- 35- MAMOUN, R. Z.; MORISSON, M.; REBEYROTTE, N.; BUSETTA, B.; COUEZ, D.; KETTMANN, R.; HOSPITAL, M.; GUILLEMAIN, B. Sequence variability of bovine leukemia virus env gene and its relevance to the structure and antigenicity of the glycoproteins. **Journal Virology**, v. 64, n. 9, p. 4180-4188, 1990.
- 36- RICE, N. R.; STEPHENS, R. M.; COUEZ, D.; DESCHAMPS, J.; KETTMANN, R.; BURNY, A.; GILDEN, R. V. The nucleotide sequence of the env gene and post-env region of bovine leukemia virus. **Virology**, v. 138, n. 1, p. 82-93, 1984.
- 37- SAGATA, N.; YASUNAGA, T.; TSUZUKU-KAWAMURA, J.; OHISHI, K.; OGAWA, Y.; IKAWA, Y. Complete nucleotide sequence of the genome of bovine leukemia virus: its evolutionary relationship to other retroviruses. **Proceedings National Academy Sciences USA**, v. 82, n. 3, p. 677-681, 1985.
- 38- KELLY, E. J.; JACKSON, M. K.; MARSOLAIS, G.; MORREY, J. D.; CALLAN, R. J. Early detection of bovine leukemia virus in cattle by use of the polymerase chain reaction. **American Journal Veterinary Research**, v. 54, n. 2, p. 205-209, 1993.
- 39- NAIF, H. M.; DANIEL, R. C. W.; COUGLE, W. G.; LAVIN, M. F. Early detection of bovine leukemia virus by using an enzyme-linked assay for polymerase chain reaction-amplified proviral DNA in experimentally infected cattle. **Journal Clinical Microbiology**, v. 30, n. 3, p. 675-679, 1992.