

Poisoning by Diacetoxyscirpenol in cattle fed citrus pulp in the state of São Paulo, Brazil

Intoxicação por Diacetoxyscirpenol em bovinos alimentados com polpa cítrica no Estado de São Paulo, Brasil

Marcelo GALHARDO¹; Eduardo Harry BIRGEL Júnior¹; Lúcia Maria Valente SOARES²; Régina Prado Zanes FURLANI²; Eduardo Harry BIRGEL.

Short communication

CORRESPONDENCE TO:
Eduardo Harry Birgel Júnior
Departamento de Clínica Veterinária
Faculdade de Medicina Veterinária e
Zootecnia da USP
Av. Prof. Dr. Orlando Marques de
Paiva, 87 - Cidade Universitária
Armando Salles de Oliveira
05508-900 - São Paulo - SP - Brasil
e-mail: ehbirgel@usp.br

1 - Departamento de Clínica
Veterinária
Faculdade de Medicina
Veterinária e Zootecnia da
USP - SP
2 - Departamento de Ciências
de Alimentos
Faculdade de Engenharia de
Alimentos - UNICAMP - SP

SUMMARY

The death of cows poisoned by the trichothecene is reported from a cattle herd of animals fed with citrus pulp containing 7 mg/kg of the micotoxin diacetoxyscirpenol.

UNITERMS: Poisoning; Bovine; Diacetoxyscirpenol; Citrus; Pulps

INTRODUCTION

Diacetoxyscirpenol, T-2 toxin, zearalenone, nivalenol, deoxynivalenol (vomitoxin) are mycotoxins (trichothecenes) isolated from strains of the following *Fusarium* species: *F. episphae*, *F. lateridium*, *F. nivale*, *F. oxysporum*, *F. tricinctum*; species of *Cephalosporium*, *Myrothecium*, *Trichoderma* and *Stachybotrys* have also been reported to produce trichothecenes (World Health Organization⁹, 1979). They grow on cereal grains such as wheat, barley, oats, rye, millet and corn (Oehme⁶, 1990).

Dried citrus pulp can replace 67% of the maize in concentrate mixtures for lactating cows at pasture, without affecting neither the volume nor fat content of milk production (Veloso⁸, 1985).

Poisoning by diacetoxyscirpenol has several characteristic, clinical signs which may include: anorexia, digestive disorders, diarrhea, progressive weight loss and even death with hemorrhagic lesions in the stomach, heart, intestines, lungs, urinary bladder and kidneys (Smalley⁷, 1973). Pigs (Coppock *et al.*², 1989; Macri *et al.*³, 1991), dogs (Coppock *et al.*², 1989), cattle (Coppock *et al.*², 1989) and mice (Mayura *et al.*⁵, 1987) are susceptible to the toxin. The DL-50 for mammals range from 0.5 to 1.0 mg/kg (Busby; Wogang¹, 1988)

During November 1994 the Hospital Veterinário da Faculdade de Medicina Veterinária e Zootecnia da Universidade de São Paulo attended 2 adult, Holstein cows with alopecia of the head and neck, moderate bloody diarrhea, signs of light, yellowish and/or reddish discoloration of the ocular mucosa, increased hepatic area, dehydration, lack of milk production, prostration, weakness, decumbency and death after 10-15 days. Serum levels of urea, creatinine, aspartate aminotransferase (AST) and gamma glutamyltransferase (GGT) were moderately increased. Discrete hypochromic anemia was diagnosed in one case. At necropsy, liver and kidneys were congested and hemorrhages were observed in the abomasum and bowel. The owner reported the deaths of 6 other cows, of a total of 200 bovines on the farm, which had shown similar symptoms. These signs appeared 30 days after the introduction of citrus pulp into the diet.

Clinical signs, gross lesions and the history of the disease suggested poisoning by trichothecenes contained in the citrus pulp. Examination of the citrus pulp for fungi revealed only *Aspergillus* sp., *Penicillium* sp and *Mucor* sp. Chemical analysis by thin-layer chromatography (Marochi ; Soares⁴, 1993) detected 7 mg/kg of Diacetoxyscirpenol. The fact that *Fusarium* was not found in the citrus pulp, but only the toxin, suggest that the contamination occurred in the orange groove.

RESUMO

Relata-se a morte de bovinos causada pela ingestão de polpa cítrica contaminada pela micotoxina Diacetoxyscirpenol (7mg/kg), pertencente ao grupo dos tricotecenos.

UNITERMOS: Envenenamento; Bovinos; Diacetoxyscirpenol; Polpa cítrica

REFERENCES

- 1-BUSBY JR., W.F.; WOGANG, G.N. Trichotechenes. In: SHANK, R. C. **Mycotoxin and n-nitroso compounds; environmental risk**. Boca Raton, CRC Press, p.29-45, 1988.
- 2-COPPOCK, R.W.; HOFFMANN, W.E.; GELBERG, H.B.; BASS, D.; BUCK, W.B. *American Journal of Veterinary Research*, v.50, n.3, p. 411-5, 1989.
- 3-MACRI, B.; PANEBIANCO, A.; GALOFARO, V.; DUGO, G.; CRISEO, G. Micotossicosi da diacetossiscirpenolo (DAS) in suinetti. Aspetti anatomo-istopatologici e considerazioni patogenetiche. *Selezione Veterinaria*, v.32, p.321-34, 1991. Suppl. 1.
- 4-MAROCHI, M.A.; SOARES, M.L.V. Metodologia para determinação de tricotecenos e zearalenona em grãos. *Boletim da Sociedade Brasileira de Ciência e Tecnologia de Alimentos*, v.27, n.1, p.1-8, 1993.
- 5-MAYURA, K.; SMITH, E.E.; CLEMENT, B.A.; HARVEY, R.B.; KUBENA, L.F.; PHILLIPS, T.D. *Toxicology*, v.45, p.245-55, 1987.
- 6-OEHME, F.W. Disorders caused by toxins. In: SMITH, B.P. **Large animal internal medicine**, St. Louis, C.V. Mosby: diseases of horses, cattle, sheep and goats, 1990, p.1661.
- 7-SMALLEY, B.E. T-2 Toxin. *Journal of the American Veterinary Medical Association*, v.163, n.11, p.1278-81, 1973.
- 8-VELOSO, L. Uso da polpa cítrica na alimentação animal. *Comunicações Científicas da Faculdade de Medicina Veterinária e Zootecnia da Universidade de São Paulo*, v.9, n.2, p.163-80, 1985.
- 9-WORLD HEALTH ORGANIZATION. **Mycotoxins**. Geneva, 1979, 127 p. (Environmental Health Criteria, 11).

Recebido para publicação: 27/12/94
Aprovado para publicação: 24/06/96