

Pediatric medicines and their relationship to dental caries

Valdenice Aparecida de Menezes^{1,*}, Gabriela Cavalcanti², Cristiana Mora², Ana Flávia Granville Garcia³, Rossana Barbosa Leal⁴

¹Dentistry College, University of Pernambuco, ²Dentistry College of the Caruaruense Higher Education Association, ³Department of Dentistry, State University of Paraíba, ⁴Dentistry College of the Caruaruense Higher Education Association

The aim of this study was to evaluate mothers' knowledge about the cariogenic potential of pediatric medicines. A total of 111 mothers were interviewed using a standardized form containing 15 questions relating to the association of pediatric drug use with dental caries and oral hygiene care. Descriptive and inferential statistics (Chi-square test and Fisher's exact) were used at a significance of 5%. Most of the mothers were aged 40 years or under (77.4%), high school educated (30.6%) and not working (50.5%). The association between medication use and dental caries or defects in teeth structure was mentioned by 35 (43.2%) mothers, 33 of whom (40.7%) cited this was due to the presence of sugar in the formulations. Only 32 mothers (28.8%) performed oral hygiene for the child after drugs ingestion, although 81.1% (n = 90) had never received guidance on the importance of this practice. The type of occupation and maternal education level were not significant in these issues (p > 0.05). Pediatric medicines can create problems for the teeth and a high percentage of mothers are unable to establish a clear cause and effect relationship with this association. Therefore, the pharmaceutical industry needs to be more aware of this and should prepare pediatric medicines without the presence of sucrose.

Uniterms: Pediatric medicine/cariogenic potential. Dental caries/prevention.

O objetivo do estudo foi verificar o conhecimento materno sobre o potencial cariogênico de medicamentos infantis. Foram entrevistadas 111 mães por meio de formulário padronizado contendo 15 perguntas relativas à associação do uso de fármacos pediátricos com a cárie dentária e aos cuidados com a higiene bucal. Foi usada a estatística descritiva e inferencial (Qui-quadrado de Pearson e exato de Fischer), com significância de 5%. A maioria das mães tinha até 40 anos (77,4%), ensino médio (30,6%) e não trabalhava (50,5%). A associação entre uso de medicamentos e cárie dental ou defeitos na estrutura dos dentes foi apontada por 35 (43,2%) mães, das quais 33 (40,7%) devido à presença de açúcar nas formulações. Apenas 32 (28,8%) realizavam a higienização bucal da criança após a ingestão de medicamentos; 81,1% (n=90) nunca receberam orientações quanto à importância desta prática. O tipo de ocupação e o grau de escolaridade materno não foram significativos nestas questões (p>0,05). Os medicamentos pediátricos podem trazer problemas para os dentes e alto percentual de mães não consegue estabelecer claramente a relação de causa e efeito existente em tal associação, portanto, é necessária maior conscientização das indústrias farmacêuticas para elaboração de medicamentos pediátricos sem a presença de sacarose.

Unitermos Medicamento infantil/potencial cariogênico. Cárie dental/prevenção.

INTRODUCTION

Dental caries is one of the most common oral diseases in childhood (Brazil, 2004). Its etiology is multifactorial and related to an imbalance between the tooth

structure and oral environment, of which microbiota, diet and host are responsible for the disease initiation and progression (Shaw, Glenwright, 1989). Updated concepts also consider social and behavioral factors of the individual (Gill, 2003).

Concerning diet components, sugar and especially sucrose serves as a substrate for fermentation of the oral microbiota, in addition to influencing the production of acids and the type and amount of biofilm (Pinto, 2000).

*Correspondence: V. A. Menezes. Faculdade de Odontologia, Universidade de Pernambuco. Av. Gal Newton Cavalcanti, n.1650 - 54753-020 - Camaragibe - PE, Brasil. E-mail: valdenicemenezes@terra.com.br

Therefore, sugar addition to various medications for children is a supplementary source of carbohydrates for pediatric patients. Its purpose is to mask the unpleasant taste of some active constituents, allowing for better acceptance by patients (Shaw, Glenwright, 1989; Hebling, 2002; Neves, 2006). Studies have emphasized that a significant portion of the population ignores the type and concentration of sugars added to foods and/or beverages, including pediatric formulations (Mentes, 2001; Pierro, *et al.*, 2005).

Studies show a positive association between intake of these drugs and dental caries. This represents a cause for concern for children's oral health, since many studies have shown increased prevalence of the disease and its relationship with the habitual intake of these liquid formulations (Shaw, Glenwright, 1989; Mackie *et al.*, 1993; Bigeard, 2000; Neves 2006; Neves, Pierro, Maia, 2007). However, there are studies in which the true relationship between drug intake and caries rates in the population assessed was not proven. (Campos *et al.*, 2006).

Thus, children with chronic or recurrent health problems who make frequent use of medicines are particularly at risk (Souza, *et al.*, 2002, Peres *et al.*, 2005), whereas caries is a sucrose-dependent disease. Moreover, as many liquid medicines for children have low endogenous pH, they can also promote dental erosion, particularly if they stay for prolonged periods in contact with the tooth surface (Neiva, 2001; Pierro, *et al.* 2005; Neves, Pierro, Maia, 2007).

The use of sugar products (syrups, expectorants, antibiotics in solution, tonic and homoeopathic products) and those that cause reduced salivary flow (benzodiazepines, antihistamines, appetite suppressants, antiparkinsonian, hypotension, muscle relaxants, diuretics, etc.) become potentially cariogenic with regular use as does any other product containing sugar, especially when used during the night (Silva, Guimarães, 2001; Neves, Pierro, Maia, 2007).

Against this background, this study aimed to evaluate the mothers' knowledge about the use of pediatric medicines and their cariogenic potential.

METHODS

This was a cross-sectional study, with exploratory clinical-epidemiological characteristics. The project was approved by the Committee of Ethics in Research of the Caruaruense Higher Education Association under the No. 071/07. The study was conducted at the Center for Dental Specialties (CEO) of the Early Childhood Care and Clinical Dentistry, Faculty of Dentistry, Caruaru (FOC), located in Caruaruense Association of Higher Education (ASCES). The study involved interviewing 111 mothers of children aged between 0 and 12 years of age whose

children were screened or receiving dental care from February to July 2008. The interviews were conducted by two researchers in a private room, at the facilities of the institution using a questionnaire containing 15 questions, consisting of two parts: the first recorded mothers' identification and demographic data and the second part evaluated data relevant to the proposed objectives.

First, the purpose of the research and the participation that would be expected of the mothers was explained to them, after which they signed a consent form.

The responses were noted during the interview, allowing for greater fidelity and accuracy of the information, and avoiding inaccuracies due to memory failure. The reliability of responses was tested by the method of validation of "face" in 10% of the respondents (Frankfort-Nachimir, Nachimir, 1992). In this method, the researcher asks decision makers to make clear in their own words, what they understood by each question.

The data obtained was calculated using descriptive statistics by means of absolute distribution, percentage of measures and techniques of inferential statistics. The software used was EPI-INFO version 3.3 and Microsoft® Office Excel 2003.

RESULTS

Table I highlights that the majority of mothers were aged younger than 40 years (77.4%), high school educated (30.6%) and had children under the age of 6 years (62.1%).

Of the diseases that required the use of medicine, the most cited were "Asthma/ bronchitis" (33.1%), "Allergy" (23.8%) and infection (21%), for which antibiotics (45.6%) and antihistamines (33.3%) were the most used. The presentation of medicines in "suspension" and "syrup" corresponded to (52.6%), as shown by results presented in Table II.

In Table III, no significant association is evident between issues relating to the mothers' knowledge about the use of pediatric medicines and educational level of mothers surveyed.

Table IV shows that the percentage of mothers who reported problems when the child took medicine and those who brush with toothpaste was higher in the group of mothers who worked outside the home, at 41.5% and 39.6%, respectively. A significant association of the cited items with occupation was identified ($p < 0.05$).

DISCUSSION

Oral health care should be initiated at an early age, with the preventive approach to maintaining health as

TABLE I - Sample Distribution by sociodemographic variables

Variable	N	%
• Maternal age (years)		
21 to 30	35	31.5
31 to 40	51	45.9
41 or older	25	22.5
• Number of children		
One	39	35.1
Two	43	38.7
Three or more	29	26.1
• Mother's educational level		
Functional illiterate	4	3.6
Incomplete elementary	16	14.4
Elementary	33	29.7
High school	34	30.6
Undergraduate	24	21.6
• Mother's occupation		
Housewife	56	50.5
Working	53	47.7
Not reported	2	1.8
• Childrens' Age (years)		
Under 3	34	30.6
Between 3 and 6	35	31.5
Older than 6	42	37.8
TOTAL	111	100.0

the overall goal of this education. As mothers are often responsible for the administration and care related to the use of pediatric medicines, they should be counseled about the risk factors associated with caries, confirming the important role mothers play in establishing the oral health of their children.

In Brazil, parents frequently report the poor state of the oral health of their children as the main issue related to the use of antibiotics during childhood (Smith, Silva, Maia, 2002). In this study, it was found that the need to use drugs regularly for children was high (51.4%). Among these, antibiotics (45.6%) and antihistamines (33.3%), used to treat respiratory problems (33.1%), allergies (23.8%) and infections (21%), showed a higher percentage. In a recent study (Neves, Pierro, Maia, 2007), antibiotics were used with high frequency in children (88.4%), although at a lower percentage in relation to antipyretics (98.8%). These were not mentioned by the mothers, probably because of the drugs common use, this however makes their cariogenic role no less important.

TABLE II - Sample distribution by variables related to "Maternal Knowledge on Use of Pediatric Medicines"

Variable	N	%
• Does your child use medicines often?		
Yes	57	51.4
No	54	48.6
TOTAL	111	100.0
• Type of Disease		
Allergy	14	23.8
Asthma/ bronchitis	18	33.1
Cancer	1	1.7
Epilepsy	3	5.1
Inflammation	6	10.2
Infection	12	21.0
Others	3	5.1
TOTAL	57	100.0
• What was the drug used?		
Analgesic	3	5.3
Antifreeze	-	-
Antibiotic	26	45.6
Antitussive	2	3.5
Anticonvulsant	3	5.3
Antihistamine	19	33.3
Anti-inflammatory	1	1.8
Not reported	2	3.5
Others	3	5.3
BASE ⁽¹⁾	57	
• What form of presentation of these medicines have you used?		
Drops	-	-
Suspension	19	33.3
Syrup	11	19.3
Tablets	27	47.4
Capsules	-	-
Other	-	-
TOTAL	57	100.0

(1): Whereas a single patient could have taken more than one product, only the basis is registered for the calculation of percentages and not the total.

These results are corroborated by other studies in which children at greatest risk of caries development secondary to use of products containing sugar were those with chronic diseases such as asthma, epilepsy, ear in-

TABLE III - Evaluation of issues on “Maternal Knowledge about Use of Pediatric Medicines” by educational level of mothers interviewed

Study variables	Educational level						Group Total		p value
	Up to elementary		High school		Higher education				
	n	%	n	%	n	%	N	%	
• Does your child use medicines often?									
Yes	23	43.4	18	52.9	16	66.7	57	51.4	p ⁽¹⁾ =0.173
No	30	56.6	16	47.1	8	33.3	54	48.6	
• Does your child have trouble taking medicine?									
Yes	20	37.7	12	35.3	7	29.2	39	35.1	p ⁽¹⁾ =0.766
No	33	62.3	22	64.7	17	70.8	72	64.9	
TOTAL	53	100	34	100	24	100	111	100	
• Conduct when the child refuses to take the medicine									
Ignore treatment proposed	2	4.0	-	-	-	-	2	1.9	p ⁽²⁾ =0.122
Contact the doctor	1	2.0	-	-	1	4.5	2	1.9	
Child is forced to take the medicine	35	70.0	25	75.8	20	90.9	80	76.2	
Add sugar to taste	12	24.0	6	18.2	1	4.5	19	18.1	
Other	-	-	2	6.1	-	-	2	1.9	
TOTAL	50	100	33	100	22	100	105	100	
• Can medicines for children cause problems for their teeth?									
Yes	36	67.9	29	85.3	16	66.7	81	73.0	p ⁽¹⁾ =0.151
No	17	32.1	5	14.7	8	33.3	30	27.0	
TOTAL	53	100	34	100	24	100	111	100	
• If yes, what?									
Dental caries	16	44.4	9	31.0	8	50.0	33	40.7	p ⁽¹⁾ =0.386
Weaker Teeth	18	50.0	17	58.6	6	37.5	41	50.6	p ⁽¹⁾ =0.397
Structural defects	1	2.8	1	3.4	-	-	2	2.5	p ⁽²⁾ =1.000
Others	2	5.6	5	17.2	3	18.8	10	12.3	p ⁽²⁾ =0.206
BASIS⁽³⁾	36		29		16		81		
• Action after child took drugs containing sugar:									
Brushing	4	7.5	6	17.6	5	20.8	15	13.5	p ⁽¹⁾ =0.168
Brushing + tooth paste	18	34.0	7	20.6	7	29.2	32	28.8	p ⁽²⁾ =0.405
Washing with running water	12	22.6	5	14.7	2	8.3	19	17.1	p ⁽²⁾ =0.275
Use of chewing gum	-	-	1	2.9	-	-	1	0.9	p ⁽¹⁾ =0.271
None	20	37.7	15	44.1	10	41.7	45	40.5	p ⁽²⁾ =0.833
BASIS⁽³⁾	53		34		24		111		
• Guidance on care after use of sugary drugs:									
Yes	9	17.0	4	11.8	8	33.3	21	18.9	p ⁽²⁾ =0.105
No	44	83.0	30	88.2	16	66.7	90	81.1	
TOTAL	53	100	34	100	24	100	111	100	

(1): Using the Chi-square test. (2): Using Fisher's Exact test. (3): Whereas a single survey could cite more than one alternative, it is only registered in the basis for the percentages calculation and not the total.

TABLE IV – Evaluation of issues on “Maternal Knowledge about Use of Pediatric Medicines”, by occupation of mothers interviewed

Study variables	Occupation (work outside home)				Total group		p value
	YES		NO		n	%	
	N	%	n	%			
• Does your child use medicines often?							
Yes	29	54.7	28	50.0	57	52.3	p ⁽¹⁾ =0.622
No	24	45.3	28	50.0	52	47.7	
• Does your child have trouble taking medicine?							
Yes	22	41.5	16	28.6	38	34.9	p ⁽¹⁾ =0.157
No	31	58.5	40	71.4	71	65.1	
TOTAL	53	100	56	100	109	100	
• Conduct when the child refuses to take the medicine							
Ignore treatment proposed	1	1.9	1	2.0	2	1.9	p ⁽¹⁾ =0.974
Contact doctor	1	1.9	1	2.0	2	1.9	
Child is forced to take the medicine	39	75.0	40	78.4	79	76.7	
Add sugar to taste	10	19.2	8	15.7	18	17.5	
Other	1	1.9	1	2.0	2	1.9	
TOTAL	52	100	51	100	103	100	
• Can medicines for children cause problems for their teeth?							
Yes	38	71.7	41	73.2	79	72.5	p ⁽¹⁾ = 0.859
No	15	28.3	15	26.8	30	27.5	
TOTAL	53	100	56	100	109	100	
• If yes, what?							
Dental carie	16	42.1	16	39.0	32	40.5	p ⁽¹⁾ = 0.780
Weaker Teeth	17	44.7	23	56.1	40	50.6	
Structural defects	1	2.6	1	2.4	2	2.5	p ⁽²⁾ = 1.000
Others	5	13.2	4	9.8	9	11.4	
BASIS⁽³⁾	38		41		79		
• Action after her child took drugs containing sugar:							
Brushing	2	3.8	12	21.4	14	12.8	p ⁽¹⁾ =0.006*
Brushing + tooth paste	21	39.6	11	19.6	32	29.4	
Washing with running water	7	13.2	11	19.6	18	16.5	p ⁽¹⁾ = 0.366
Use of chewing gum	1	-	-	-	1	0.9	
None	23	43.4	22	39.3	45	41.3	p ⁽¹⁾ = 0.700
BASIS⁽³⁾	53		56		109		
• Guidance on care after use of sugary drugs:							
Yes	12	22.6	9	16.1	21	19.3	p ⁽¹⁾ = 0.385
No	41	77.4	47	83.9	88	80.7	
TOTAL	53	100	56	100	109	100	

(*): Significant association at the level of 5.0%. (1): Using the Chi-square test. (2): Using Fisher’s Exact test. (3): Whereas a single survey could cite more than one alternative, it is only registered in the basis for the percentages calculation and not the total.

flammation or upper respiratory tract infection (Durward, Thou, 1997), heart, respiratory or renal (Silva Santos, 1994) disease who make frequent use of these drugs (Smith, Silva, Maia, 2002).

The presence of sucrose and/or other fermentable carbohydrates in the formulation of pediatric medicines and the low pH values interfere with the cariogenic and erosive potential displayed by these drugs. Other factors are

also cited such as: acidity, frequency of administration and formulation (Costa *et al.*, 2004; Marquezan *et al.*, 2007).

While many parents recognize that sugar causes tooth decay, they do not normally associate this change with the sugar added to pediatric liquid medicines (Mentes, 2001). This fact was clearly seen in the present study: of 81 mothers (73%) who reported that pediatric medicines can cause problems for their child's teeth, only 33 (40.7%) associated them with the occurrence of dental caries, correctly linking them with the presence of sugars in the formulations. Note that 50.6% of mothers believed medicines can make teeth weaker, with no necessary connection with the sugar addition (Table III). The reported association with defects in teeth structure was low (2.5%) in contrast to another study (Neves, Pierro, Maia, 2007) which detected a percentage of 70.9%. The type of occupation and level of maternal education did not differ on these issues ($p > 0.05$).

Regarding the medicine intake, it was found that the percentage of mothers who reported problems was higher in the group who worked outside the home (41.5%). Moreover, the practice of forcing children to swallow the medicine when they refused was higher among mothers with a higher education (90.9%). The percentage of mothers who add sugar to improve the taste of the medicine was lower among those with a higher education (4.5%) but very similar between groups of mothers with up to elementary school and high school education (70.0% and 75.8%, respectively) ($p > 0.05$). This data clearly demonstrates the importance of education in maternal behavior.

The risk of developing caries associated with the use of sugar products becomes aggravated when no measure of oral hygiene is performed (Bigéard, 2000). However, despite all the risk involved, 40.5% of mothers did not adopt any action after the administration of the dose prescribed, a higher rate compared to the findings of another study (28.6%) (Neves, Pierro, Maia, 2007). In addition, only 28.8% of mothers had an adequate oral hygiene routine for their children, using a brush and toothpaste. The percentage of those who performed oral hygiene was lower among those who had elementary education (7.5%).

It is noteworthy that 81.1% (90) of mothers had never received guidance on the importance of oral cleaning after intake of medicines. Other authors found similar results (Paiva, 1991; Souza, *et al.*, 2002; Neves, Pierro, Maia, 2007). The type of occupation and level of maternal education were not significant factors influencing these issues ($p > 0.05$).

Further, drugs administered in the evening or at bedtime, the period when there is a decrease in salivary flow as well as reduced reflexes of swallowing and muscle

movement, remain longer in the oral cavity (Durward, Thou, 1997; Bigéard, 2000; Costa *et al.*, 2004).

It is noteworthy that data showed that most (52.6%) drugs had been prescribed in liquid form (syrup or suspension) for which elimination occurs more slowly. However, most mothers had children under the age of six years (62.1%), a phase in which the indication of use of tablets is not common, and therefore the adoption of proper practice of oral hygiene after ingesting certain drugs by children is essential. Another aspect to consider is the use of a variety of sugar products without prescription such as throat lozenges, cough syrup, lickers, chewable vitamins and liquid tonics, which require dietary advice, and intensification of preventive methods against caries (Paiva, 1991).

Reducing the cariogenic potential of pediatric medicines should be a concern among all health professionals. Although the market has already made available sugar-free drugs or sweetened with other substances, medicinal products containing sucrose are still among the most prescribed and do not warn about the risk of developing carious lesions (Durward, Thou, 1997). Moreover, the manufacture of pediatric medicines not containing fermentable carbohydrates, could be the best health policy. In addition, research should be conducted to find acceptable levels of fermentable carbohydrates to preserve the taste of medicines (Peres *et al.*, 2005).

Therefore, it is essential that clear instructions regarding oral hygiene be provided for each prescription of medicines containing sugar, in order to avoid the development of caries in many children, particularly those who consume medicines on a routine basis and especially those who have chronic diseases.

Although the relationship between the continued use of sugar products and the development of dental caries is not clearly established, the administration of these drugs should be done preferably during meals and before sleeping (Bigéard, 2000) and should be followed by instructions for cleaning the oral cavity.

CONCLUSIONS

Although most mothers recognize that pediatric medicines can cause teeth problems, a high percentage are unable to establish a clear cause and effect relationship with this association. Pediatric formulations without added sugar should be available to those responsible, and professionals could provide better and more intensive teaching on the proper oral hygiene care after use of these medicines. Therefore, the pharmaceutical industry needs to be more aware of this problem and should prepare pediatric medicines without the presence of sucrose.

REFERENCES

- BIGEARD, L. The role of medication and sugars in pediatric dental patients. *Dent. Clin. North Am.*, v.44, n.3, p.443-456, 2000.
- BRASIL. Ministério da Saúde. Secretaria de Atenção a Saúde. Departamento de Atenção Básica. Projeto SB Brasil 2003: condições de saúde bucal da população brasileira 2002-2003: resultados principais. Brasília: Ministério da Saúde, 2004. Available at: <http://bvsmms.saude.gov.br/bvs/publicacoes/projeto_sb2004.pdf>. Accessed on: 06 out. 2008.
- CAMPOS, J. A. D. B.; GIRO, E. M. A.; ORRICO, S. R. P.; OLIVEIRA, A. P. C.; LORENA, S. M. Correlação entre a prevalência de cárie e a utilização de medicamentos em pacientes com necessidades especiais institucionalizados e não institucionalizados. *Salusvita*, v.25, n.1, p.35-42, 2006.
- COSTA, C. C.; ALMEIDA, I. C.; RAYMUNDO M. S.; FETT, R. Análise do pH endógeno, da acidez e da concentração de sacarose de medicamentos pediátricos. *Rev. Odonto Ciênc.*, v.19, n.44, p.164-169, 2004.
- DURWARD, C.; THOU, T. Dental caries and sugar containing liquid medicines for children in New Zealand. *N. Z. Dent. J.*, v.93, n.414, p.124-129, 1997.
- FRANKFORT-NACHIMIAS, C.; NACHIMIAS, D. *Research methods in the social*. 4.ed. London: Edward Arnold, 1992. 144 p.
- GONÇALVES, M. R. Cárie dental: uma doença psicossomático-ambiental. *J. Bras. Clin. Odontol. Integr.*, v.7, n.39, p.251-255, 2003.
- HEBLING, J. ; TEIXEIRA, H. M.; PIZZOLITTO, A. C. Potencial cariogênico de medicamentos infantis. *Robrac*, v.11, n.31, p.35-39, 2002.
- MACKIE, I. C.; WORTHINGTON, H. V.; HOBSON, P. An investigation into sugar containing and sugar-free over-the counter medicines stocked and recommended by pharmacists in the north western region of England. *Br. Dent. J.*, v.175, n.3, p.93-98, 1993.
- MARQUEZAN, M.; MARQUEZAN, M., POZZOBON, R. T., OLIVEIRA, M. D. M. Medicamentos utilizados por pacientes odontopediátricos e seu potencial cariogênico. *RPG Rev. Pós Grad., Santa Maria*, v.13, n.4, p.334-339, 2007.
- MENTES, A. pH changes in dental plaque after using sugar-free pediatric medicine. *J. Clin. Pediatr. Dent.*, v.25, n.4, p.307-312, 2001.
- NEIVA, A.; SILVA, V. S.; MAIA, L. C.; SOARES, E. L.; TRUGO, L. C. Análise *in vitro* da concentração de sacarose e pH de antibacterianos de uso pediátrico. *Pesqui. Bras. Odontopediatria Clín. Integr.*, v.1, n.1, p.9-16, 2001.
- NEVES, B. G. *Avaliação dos fatores relacionados ao potencial cariogênico e erosivo de medicamentos líquidos infantis*. Rio de Janeiro, 2006. 99 p. [Dissertação de Mestrado. Faculdade de Odontologia. Universidade Federal do Rio de Janeiro].
- NEVES, B. G.; PIERRO, V. S.; SILVA, M. L. C. Perceptions and attitudes among parents and guardians on the use of pediatric medicines and their cariogenic and erosive potential. *Ciênc. Saúde Coletiva*, v.12, n.5, p.1295-1300, 2007.
- PAIVA, M. S. Cariogenicidade dos medicamentos: um risco a saúde bucal. *Rev. Gaúcha Odontol.*, v.39, n.5, p.336-338, 1991.
- PERES, K. G.; OLIVEIRA, C. T.; PERES, M. A.; RAYMUNDO, M. S.; FETT, R. Sugar content in liquid oral medicines for children. *Rev. Saúde Pública*, v.39, n.3, p.486-489, 2005.
- PIERRO, V. S. S.; ABDELNUR, J. P.; MAIA, L. C.; TRUGO, L. C. Free sugar concentration and pH of pediatric medicines in Brazil. *Community Dent. Health*, v.22, n.3, p.180-183, 2005.
- PINTO, V. G. *Saúde bucal coletiva*. 4.ed. São Paulo: Editora Santos, 2000. 541 p.
- SHAW, L.; GLENWRIGHT, H. D. The role of medications in dental caries formation: need for sugar-free medication for children. *Pediatrician*, v.16, n.3-4, p.153-155, 1989.
- SILVA, M. F. A.; GUIMARÃES, J. A. T. L. Potencial cariogênico de medicamentos utilizados no tratamento de doenças do trato respiratório e alergia. *JBP, J. Bras. Odontopediatr. Odontol. Bebê*, v.4, n.21, p.383-386, 2001.
- SILVA, S. M. B.; SANTOS, C. F. Pediatric Medicines and Caries Risk – A Review. *Rev. Fac. Odont. Bauru*, v.2, n.4, p.15-21, 1994.

- SILVA, V. S.; SILVA, A. N.; MAIA, L. C. Antibacterianos pediátricos e cárie dental em crianças: mitos e realidades. *Rev. Paul. Pediatr.*, v.20, n.4, p.191-195, 2002.
- SOUZA, M. I. C.; SEGAL, M. T.; MEDEIROS, U. V.; BARBOSA, A. R. S. Potencial cariogênico dos xaropes infantis para doenças respiratórias prolongadas. *JBP, J. Bras. Odontopediatr: Odontol. Bebê*, v.5, n.25, p.209-214, 2002.

Received for publication on 18th December 2008

Accepted for publication on 11th October 2009