

CLINICAL SCIENCE

Macroscopic placental changes associated with fetal and maternal events in diabetes mellitus

Ana Karina Marques Salge,¹ Karlla Morgana Nunes Rocha,¹ Raphaela Maioni Xavier,¹ Wilzianne Silva Ramalho,¹ Érika Lopes Rocha,¹ Janaína Valadares Guimarães,¹ Renata Calciolari Rossi e Silva,¹ Karina Machado Siqueira,¹ Douglas Reis Abdalla,² Márcia Antoniazzi Michelin,² Eddie Fernando Candido Murta²

¹Goias Federal University, Nursing School, Goiania/GO, Brazil. ²Oncology Institute Research (IPON) of Triângulo Mineiro Federal University, Uberaba/MG, Brazil.

OBJECTIVES: The current study sought to identify macroscopic placental changes associated with clinical conditions in women with or without diabetes and their newborns.

METHODS: The study population consisted of 62 pregnant women clinically diagnosed with diabetes and 62 healthy women (control group).

RESULTS: Among the subjects with diabetes, 43 women (69.3%) were diagnosed with gestational diabetes mellitus, 15 had diabetes mellitus I (24.2%), and four had diabetes mellitus II (6.5%). The mean age of the women studied was 28.5 ± 5.71 years, and the mean gestational age of the diabetic women was 38.51 weeks. Of the 62 placentas from diabetic pregnancies, 49 (79%) maternal surfaces and 59 (95.2%) fetal surfaces showed abnormalities, including calcium and fibrin deposits, placental infarction, hematoma, and fibrosis. A statistical association was found between newborn gender and fetal and maternal placental changes ($p=0.002$). The mean weight of the newborns studied was $3,287 \pm 563$ g for women with diabetes mellitus, $3,205 \pm 544$ g for those with gestational diabetes mellitus, $3,563 \pm 696$ g for those with diabetes mellitus II, and $3,095 \pm 451$ g for those with diabetes mellitus I.

CONCLUSIONS: Infarction, hematoma, calcification, and fibrin were found on the maternal and fetal placental surfaces in women with diabetes. Women with gestational diabetes and post-term infants had more calcium deposits on the maternal placental surface as compared to those with type I and type II diabetes.

KEYWORDS: Diabetes Mellitus; Newborn; Placenta; Pregnancy.

Salge AK, Rocha KM, Xavier RM, Ramalho WS, Rocha ÉL, Guimarães JV, et al. Macroscopic placental changes associated with fetal and maternal events in diabetes mellitus. *Clinics*. 2012;67(10):1203-1208.

Received for publication on April 15, 2012; First review completed on May 2, 2012; Accepted for publication on June 22, 2012

E-mail: karllamorgan@hotmail.com

Tel.: 55 62 3521-1822

INTRODUCTION

The human placenta regulates maternal and fetal physiological changes during pregnancy. It also plays a major role in pregnancy-related complications, such as diabetes mellitus (DM) (1).

DM is a common clinical condition that affects approximately 1 to 15% of pregnant women. Maternal metabolic control has been directly associated with complications and deaths in pregnant women with DM (2). Diabetes during pregnancy can lead to or aggravate placental abnormalities, which will impact the systemic balance between the pregnant woman and her fetus. DM can increase the frequency of placental dysfunction and alter its structure

and function, and it is also associated with increased rates of maternal and perinatal morbidity and miscarriage. The nature, extent, and specificity of these abnormalities depend on the gestational period and the severity and type of diabetes.

Although the mechanisms of pathogenesis are not fully understood, it is believed that hormonal adjustments during pregnancy, including increased levels of estrogen, progesterone, cortisol, prolactin, and human placental lactogen, play an important role. These changes lead to metabolic imbalance and may result in increased insulin resistance and DM onset in susceptible women and uncontrolled blood glucose levels in pre-diabetic women, especially during the last two trimesters of pregnancy (3).

DM complications during pregnancy include polyhydramnios (clinically diagnosed in pregnant women with poorly controlled DM); premature rupture of membranes and pre-term births (4); hypertensive disorders of pregnancy, particularly preeclampsia (5); and higher rates of spontaneous abortion, macrosomia (6), birth defects, stillbirth, respiratory distress syndrome, and neonatal hypoglycemia (4).

Copyright © 2012 **CLINICS** – This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

No potential conflict of interest was reported.

Placental changes are believed to be partly responsible for the high incidence of fetal complications observed in pregnant women with DM (1). However, the relationship between DM and the macroscopic placental changes that occur during pregnancy is not well established. Thus, the current study aimed to identify macroscopic placental changes associated with clinical outcomes in women with and without DM and their newborns.

MATERIALS AND METHODS

This prospective study was conducted at a public maternity hospital and diabetes reference center in the city of Goiânia in Central-West Brazil between August 2009 and July 2010. The study population included 62 pregnant women clinically diagnosed with DM. These women were matched to a control group that comprised 62 healthy women considered to be at a low obstetric risk who had undergone vaginal delivery or cesarean section. These women had no maternal complications during pregnancy, their obstetric and laboratory tests were normal, and they had clinically normal babies born at term.

The following exclusion criteria were applied: women with infectious diseases, placentas in formaldehyde, cases of stillbirths and intrauterine fetal deaths, pregnant women with autoimmune diseases or other diseases that could lead to immunological changes, and those who received corticosteroid therapy during labor. Stillborn fetuses were defined as any fetus at a gestational age of at least 22 weeks and/or fetal weight ≥ 500 g whose death occurred before complete separation from the maternal body and who did not show any of the following signs compatible with life at birth: heartbeat, umbilical cord pulsation, or voluntary muscle movements (7). The control group comprised pregnant women who did not experience complications during pregnancy and who had normal laboratory tests and clinically normal newborns.

We evaluated all macroscopic changes on the fetal and maternal placental and umbilical cord surfaces. All lesions were examined, recorded, and photographed.

The evaluation of macroscopic placental parameters was performed according to protocols published by Driscoll and Langston (1991) (8) and Benirschke and Kaufmann (1995) (9).

Data were collected from medical records and transcribed into study forms. We collected information about obstetric history; potential neonatal/fetal and maternal obstetric complications; neonatal anthropometric measurements, 1- and 5-minute Apgar scores; laboratory tests for the evaluation of DM; parity; and gestational age, which was determined according to the date of the last menstrual period, the first trimester ultrasound examination, and the Capurro method. When the results of the gestational age assessment were inconsistent, the results of the Capurro method (10) were used. Underlying maternal conditions were grouped according to the 10th Revision of International Classification of Diseases (ICD-10) criteria, and neonatal/fetal underlying conditions were grouped according to the Carlidge and Stewart criteria (7).

Statistical analyses were performed with SigmaStat® version 2.0 (San Jose, California, USA), and an electronic spreadsheet was created. Categorical variables are presented as absolute (n) and relative values (percentages). Proportions were compared using the Yates' correction or chi-squared

(χ^2) and Fisher's exact tests. Differences of $p < 0.05$ were considered statistically significant.

The research project was approved by the Research Ethics Committee of the Federal University of Goiás (UFG) and was registered in the U.S. National Institutes of Health website (protocol number 101/2008).

RESULTS

Regarding the type of diabetes, 43 women (69.3%) were diagnosed with gestational diabetes mellitus (GDM), 15 were diagnosed with DM I (24.2%) and 4 were diagnosed with DM II (6.5%). The mean age of the women studied was 28.5 ± 5.71 years, and there was no statistical correlation between maternal age and maternal placental changes. There was a higher proportion of multigravida than secundigravida and primigravida women (23 cases [37.1%], 11 cases [17.7%], and 9 cases [14.5%], respectively) among women with GDM.

Of the 62 placentas that were macroscopically examined, 49 (79%) maternal surfaces and 59 (95.2%) fetal surfaces showed changes, including calcium and fibrin deposits, placental infarction, hematoma, and fibrosis (Table 1).

Calcification was predominantly observed on the maternal surface of 35 placentas (56.4%), although hematoma and fibrosis were not detected on the maternal surface. A greater number of placental changes were observed in pregnant women with GDM: 31 (50%) were detected on the maternal surface (Table 1) and 40 (64.5%) were detected on the fetal surface (Table 2).

The most common pathological change found on the fetal surface was the presence of fibrin, which was observed in 51 placentas (82.3%). Placental infarction and fibrosis were not found on the fetal surface (Table 2). The assessment of an association between GDM and placental changes showed that all 43 women with GDM (100%) demonstrated some sort of change in either the fetal or maternal placental surface. The most common placental changes included calcification (19; 44.1%), fibrin (11; 25.6%), and placental infarction (1; 2.3%) on the maternal surface; and fibrin (35; 81.4%), calcification (4; 9.3%), and hematoma (1; 2.3%) on the fetal surface. There were no placental changes on the maternal surface in 12 cases (28%) and on the fetal surface in three cases (7%).

Table 1 - Distribution of placental macroscopic changes on the maternal surface in 62 pregnant women with diabetes mellitus from Goiânia, Central-West Brazil, between August 2009 and July 2010.

Maternal surface alterations	DM		
	GDM	DM I	DM II
	n (%)	n (%)	n (%)
Calciphylaxis	19 (44.2)	13 (86.6)	3 (75)
Fibrin	11 (25.6)	1 (6.7)	1 (25)
Infarction	1 (2.3)	0 (0)	0 (0)
No alterations	12 (27.9)	1 (6.7)	0 (0)

DM = diabetes mellitus, GDM = gestational diabetes mellitus, DM I = type I diabetes mellitus, DM II = type II diabetes mellitus, n = number of cases.

Table 2 - Distribution of placental macroscopic changes on the fetal placental surface in 62 pregnant women with diabetes mellitus from Goiânia, Central-West Brazil, between August 2009 and July 2010.

Fetal surface alterations	DM		
	GDM	DM I	DM II
	n (%)	n (%)	n (%)
Calciphylaxis	4 (9.3)	3 (20)	0 (0)
Fibrin	35 (81.4)	12 (80)	4 (100)
Infarction	1 (2.3)	0 (0)	0 (0)
No alterations	3 (7.0)	0 (0)	0 (0)

DM = diabetes mellitus, GDM = gestational diabetes mellitus, DM I = type I diabetes mellitus, DM II = type II diabetes mellitus, n = number of cases.

Women with GDM had a greater amount of hydroxyapatite (calciphylaxis) deposited on the maternal placental surface as compared to those without DM ($p = 0.008$) (Table 3).

Among women with DM, there was a higher proportion of full-term births (45 cases, 72.6%), followed by pre-term (15 cases, 24.2%) and post-term births (two cases, 3.2 %) (Table 4).

No statistically significant relationship was found between fetal and maternal placental changes and gestational age. The mean gestational age among women with DM was 38.51 weeks. Table 5 shows the associations between gestational age and types of DM. Full-term births were more common among women with all types of DM: 29 full-term births (64.4%) occurred in women with GDM; 12 (26.7%) occurred in those with DM I; and four (8.9 %) occurred in those with DM II. There was no statistically significant relationship between gestational age and type of DM (Table 5).

A statistical association was found between newborn gender and fetal and maternal placental changes. Women who gave birth to females had more calcification on the maternal placental surface as compared to other groups

($p = 0.002$). In addition, larger fibrin deposits on the maternal placental surface were associated with mothers of male newborns ($p < 0.001$), and placental macroscopic changes were less common in mothers with female newborns ($p = 0.021$). There was also a predominance of male (33, 53.2%) as compared to female (29, 46.8%) newborns born to the women in our study.

The mean newborn weight was $3,287 \pm 563$ g in women with DM; $3,205 \pm 544$ g in those with GDM, $3,563 \pm 696$ g in those with DM II, and $3,095 \pm 451$ g in those with DM I. There was no statistically significant association between newborn weight and type of maternal DM (Table 6).

The mean 1- and 5-minute Apgar scores in newborns born to mothers with DM were 7.6 ± 1.6 and 9.5 ± 0.6 , respectively. The mean 1- and 5-minute Apgar scores of newborns of women with GDM, DM I and DM II were 7.8 ± 1.3 and 9.2 ± 0.6 , 7.4 ± 2.4 and 9.5 ± 0.7 , and 7.7 ± 1.2 and 9.7 ± 0.5 , respectively.

DISCUSSION

Several changes occur in the human placenta during pregnancies that are complicated by DM. These abnormalities are directly associated with intrauterine blood glucose levels that lead to reduced maternal-placental blood circulation and maternal-fetal exchange (11). Both macroscopic and microscopic pathological changes are associated with complications during pregnancy. For example, clinical and obstetric complications accelerate placental calcification due to stress mechanisms (12).

The results of the current study demonstrated a significant association between fibrin deposits on the fetal surface of the placenta and full-term newborns. However, a previous study with a similar sample size presented findings that contrast with those presented here (13), and another publication reported that fertility in women with DM may be affected through different mechanisms. Preimplantation human conceptuses are sensitive to insulin and may be directly affected by maternal insulin resistance, and rat models have shown that excess

Table 3 - Association between macroscopic changes on the fetal and maternal placental surfaces and gestational diabetes mellitus in 43 women from Goiânia, Central-West Brazil, between August 2009 and July 2010.

MACROSCOPIC ALTERATIONS	GDM					
	MATERNAL SURFACE		p-value	FETAL SURFACE		p-value
Calcification	Yes	No	0.008	Yes	No	0.665
	19	24		4	39	
Fibrin	Yes	No	0.310	Yes	No	1.000
	11	32		35	8	
Placental infarction	Yes	No	1.000	Yes	No	1.000
	1	42		0	43	
Hematoma	Yes	No	1.000	Yes	No	1.000
	0	43		1	42	
Fibrosis	Yes	No	1.000	Yes	No	1.000
	0	43		0	43	
No alterations	Yes	No	0.050	Yes	No	0.546
	12	31		3	40	
	1	18		0	19	

Fisher's exact test.

Table 4 - Distribution of gestational age for 62 women with or without (control of) diabetes mellitus, according to diabetes type, from Goiânia, Central-West Brazil, between August 2009 and July 2010.

GESTATIONAL AGE	GDM		DM I		DM II		CONTROL	
	n (%)		n (%)		n (%)		n (%)	
Pre-term	12 (28)		3 (20)		0 (0)		25 (40.3)	
Full-term	29 (67.4)		12 (80)		4 (100)		23 (37.1)	
Post-term	2 (4.6)		0 (0)		0 (0)		14 (22.6)	

DM = diabetes mellitus, GDM = gestational diabetes mellitus, DM I = type I diabetes mellitus, DM II = type II diabetes mellitus, n=number of cases CONTROL: women without DM. ANOVA: F=7.282 (p=0.02), followed by Dunnett’s method (p<0.05) (control vs. DM II, control vs. DM I).

insulin leads to reduced preimplantation. Although findings in rat models cannot be directly applied to humans, they provide information on how insulin resistance may affect fertility (14).

Studies of vascular systems have shown that high glucose and insulin levels profoundly affect vascular endothelial growth factor, nitric oxide and protein kinase C levels, as well as alterations in junctional adhesion molecules, such as occludin and vascular endothelial-cadherin, which cause vascular albumin leakage. Increased superoxide and nitric oxide production may also induce placental oxidative stress via the effects of prooxidant peroxynitrite, which itself leads to vascular dysfunction (15). Nitrotyrosine residues are found in placental vessels of preeclamptic and diabetic pregnancies, which further highlight the effects of oxidative stress in these conditions (16). Moreover, the abnormal molecular mechanisms underlying vascular changes in the diabetic human placenta may be consequences of high glucose and hyperinsulinemia (15).

With regard to gestational age, Sgrott (17) reported a mean gestational time of 38.51 weeks at birth in women with DM, which is similar to what we observed in the current study. However, some studies have claimed that DM may cause pre-term births (less than 38 weeks of gestation) (18).

Pregnant women diagnosed with DM are at an increased risk of clinical and obstetric complications, such as pre-term delivery, gestational hypertension, urinary tract and other infections, periodontal disease, cesarean delivery, and obstetric trauma. Pre-term delivery before 35 weeks of gestation is directly related to the severity of diabetes and proteinuria in early gestation (19). Furthermore, one-third of all premature births in women with DM are a consequence of hypertensive complications. The factors involved in increased spontaneous pre-term births in women with DM are not yet clearly understood, although they have been associated with poor blood glucose control, polyhydramnios, and infection (20). It is

believed that hospital management of women with DM may have contributed to a decrease in the number of premature births, although no reports on the relationship between DM and post-term births could be found in the literature.

In the current study, the control group demonstrated a high number of premature births, which we believe may have been because the study hospital serves a population of high-risk pregnant women. Although these women did not have clinical DM, they may have had other abnormalities that could have predisposed them to pre-term delivery and premature rupture of membranes, including urinary tract infections, polyhydramnios, and high blood pressure.

Regarding gestational age, one study found that newborns of mothers with DM I were born at lower gestational ages than those of mothers with GDM (21). In a similar study, the 1- and 5-minute Apgar scores of infants from women with DM I, DM II and GDM were 5±2 and 9±0.5, 8±1.5 and 10±0.5, and 9±1.5 and 10±0.5, respectively (22). In our study, there was no statistically significant relationship between Apgar scores and the type of maternal DM.

The literature provides conflicting results on Apgar scores. For example, one study found no significant difference in 1- and 5-minute Apgar scores of newborns from mothers with DM as compared to a control group (23), whereas another publication showed that placental changes consistent with reduced blood flow were associated with lower 1- and 5-minute Apgar scores (24).

The mean weight of newborns from women with DM II was higher than that of newborns from women with GDM and DM I. Neither our study nor that of Szylit et al. (21) found any significant differences in the mean weights of newborns born to women with different types of DM.

Fetal macrosomia, which is commonly defined as a birth weight of more than 4.000 g, is the most common fetal

Table 5 - Gestational ages for 62 pregnant women with different types of diabetes mellitus from Goiânia, Central-West Brazil, between August 2009 and July 2010.

GESTATIONAL AGE	GDM		p-value	DM I		p-value	DM II		p-value
	Yes	No		Yes	No		Yes	No	
<38 weeks of gestation	Yes	12	0.356	Yes	3	0.214	Yes	0	0.564
	No	31		No	12		No	15	
38-40 weeks of gestation	Yes	29	0.291	Yes	12	0.528	Yes	4	0.568
	No	14		No	3		No	41	
>40 weeks of gestation	Yes	2	1.000	Yes	0	1.000	Yes	0	1.000
	No	41		No	15		No	2	

Fisher’s exact test. DM = diabetes mellitus, GDM = gestational diabetes mellitus, DM I = type I diabetes mellitus, DM II = type II diabetes mellitus, n = number of cases.

Table 6 - Characteristics of infants born to 62 women with or without (control of) diabetes mellitus, according to type, from Goiânia, Central-West Brazil, between August 2009 and July 2010.

ANTHROPOMETRIC PARAMETERS	GDM	DM I	DM II	CONTROL GROUP
	n	n	n	n
Weight (g)	3.205	3.095	3.563	3.072
Head circumference (cm)	34.6	34.6	35.5	33.7
Chest circumference (cm)	33.0	34	33.2	32
Abdominal circumference (cm)	31.2	32.2	32.6	31.5
1-min Apgar score	7.8	7.4	7.7	7.4
5-min Apgar score	9.2	9.5	9.7	9.2

DM = diabetes mellitus; GDM = gestational diabetes mellitus; DM I = type I diabetes mellitus; DM II = type II diabetes mellitus; CONTROL: women without DM; min: minute; n = number of cases. ANOVA: $F = 5.351$ ($p = 0.864$) followed by Dunnett's method: control vs. DM II, control vs. DM I, control vs. GDM.

complication of maternal DM. This condition increases the risk of perineal lacerations and delivery complications (fetal dystocia) that often necessitate cesarean section. There is also a high risk of immediate complications for infants born to mothers with DM, including intracranial hemorrhage, shoulder dystocia, neonatal hypoglycemia, jaundice, and respiratory distress (25). These infants are also at an increased risk of long-term complications. Early nutritional adjustments during gestation can permanently alter carbohydrate metabolism and cause adult-onset disorders. In addition, high birth weight is a predisposing factor for insulin resistance, obesity and DM II during childhood, as it reflects inadequate glucose control that is often due to delayed prenatal care (18). Childhood obesity has become a major public health concern, and the population of overweight children in Brazil increases by 0.5% annually (26).

In conclusion, we observed infarction, hematoma, calcification, and fibrin on the maternal and fetal placental surfaces of women with DM. In addition, women with GDM and post-term infants had more calcium deposits on the maternal placental surface of the placenta, and mothers of female newborns had a larger number of calcium and fibrin deposits on the maternal placental surface.

ACKNOWLEDGMENTS

This study was conducted in the Oncology Institute of Research (IPON) of Triângulo Mineiro Federal University, Uberaba, Minas Gerais, Brazil and received grants from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq); Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (Capes); Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG); Fundação de Ensino e Pesquisa de Uberaba (FUNEPU) and Goiás Federal University, Goiânia, Goiás, Brazil.

AUTHOR CONTRIBUTIONS

Rocha KM, Xavier RM, Ramalho WS, Rocha EL, Guimarães JN, Rossi and Silva RC, Siqueira KM and Abdalla DR conceived the study, participated in its design, coordination and participants recruitment, drafted the manuscript, performed the statistical analysis and interpretation of data, and read and approved the final manuscript. Salge AK, Michelin MA and Murta EF conceived the study, participated in the study design, coordination, and participants recruitment, drafted the manuscript, performed the statistical analysis and interpretation of data, and read and approved the final manuscript.

REFERENCES

- Hayati AR, Cheah FC, Tan AE, Tan GC. Insulin-like growth factor-1 receptor expression in the placenta of diabetic and normal pregnancies. *Early Hum Dev.* 2007;83(1):41-6. <http://dx.doi.org/10.1016/j.earlhumdev.2006.04.002>.
- Maganha CA, Nomura RMY, Zugaib M. Associação entre perfil glicêmico materno e o índice de líquido amniótico em gestações complicadas pelo diabetes mellitus pré-gestacional. *Rev Assoc Med Bras.* 2009;55(2):169-74. <http://dx.doi.org/10.1590/S0104-42302009000200021>.
- Kitzmiller JL, Davidson MB. Diabetes e gravidez. In: Davidson MB. *Diabetes mellitus: diagnóstico e tratamento.* Rio de Janeiro: Revinter. 2001;p. 277-303.
- Menicatti M, Fregonesi CEPT. Diabetes Gestacional: Aspectos fisiopatológicos e tratamento. *Arq Ciênc Saúde Unipar.* 2006;10(2):105-11.
- Kitzmiller JL. Insulin lispro and the development of proliferative diabetic retinopathy during pregnancy. *Am J Obstet Gynecol.* 2001;185(3):774-5. <http://dx.doi.org/10.1067/mob.2001.117472>.
- Rodacki M, Zajdenverg L, Albernaz MS, Bencke-Gonçalves MR, Milech A, Oliveira JE. Relationship between the prevalence of anti-glutamic acid decarboxylase autoantibodies and duration of type 1 diabetes mellitus in Brazilian patients. *Braz J Med Biol Res.* 2004;37(11):1645-50.
- Cartledge PHT, Stewart JH. Effect of changing the stillbirth definition on evaluation of perinatal mortality rates. *Lancet.* 1995;346:4486-8. [http://dx.doi.org/10.1016/S0140-6736\(95\)91327-0](http://dx.doi.org/10.1016/S0140-6736(95)91327-0).
- Driscoll SG, Langston C. Placental examination in a clinical setting. *Arc Pathol Lab Med.* 1991;115(7):668-71.
- Benirschke J, Kaufmann P. *Pathology of the human placenta*, 3rd ed. New York: Springer-Verlag; 1995.
- Capurro H. Simplified method for diagnosis of gestational age in the newborn infant. *J Pediatr.* 1978;93:120-2. [http://dx.doi.org/10.1016/S0022-3476\(78\)80621-0](http://dx.doi.org/10.1016/S0022-3476(78)80621-0).
- Del Nero U, Rudge MVC, Novo NF, Calderon IMP, Brasil MAM. Metodologia para estudo do volume e densidade absoluta da placenta humana de termo. *RBGO.* 2002;24(10):669-73.
- Salafia CM, Yampolsky M. Metabolic scaling law for fetus and placenta. *Placenta.* 2009;30(5):468-71. <http://dx.doi.org/10.1016/j.placenta.2008.12.013>.
- Moretto VL, Lautert L. Features of pregnant women with diabetes mellitus. *Rev Gaucha Enferm.* 2004;25(3):334-45.
- Whitworth KW, Baird DD, Stene LC, Skjaerven R, Longnecker MP. Fecundability among women with type 1 and type 2 diabetes in the Norwegian Mother and Child Cohort Study. *Diabetologia.* 2011;54(3):516-22. <http://dx.doi.org/10.1007/s00125-010-2003-6>.
- Leach L, Taylor A, Sciota F. Vascular dysfunction in the diabetic placenta: causes and consequences. *J Anat.* 2009;215(1):69-71. <http://dx.doi.org/10.1111/j.1469-7580.2009.01098.x>.
- Myatt L, Kossenjans W, Sahay R, Eis A, Brockman D. Oxidative stress causes vascular dysfunction in the placenta. *J Matern Fetal Med.* 2000;9(1):79-82. [http://dx.doi.org/10.1002/\(SICI\)1520-6661\(200001/02\)9:1<79::AID-MFM16>3.0.CO;2-O](http://dx.doi.org/10.1002/(SICI)1520-6661(200001/02)9:1<79::AID-MFM16>3.0.CO;2-O).
- Sgrott I, Neto JAS, Schweitzer LC, Battisti R. Análise dos resultados maternos e perinatais do diabetes mellitus na gravidez. *Arq Catarinense Med.* 2009;38(3):17-23.
- Silva AM, de Almeida MF, Matsuo T, Soares DA. Risk factors for preterm birth in Londrina, Paraná State, Brazil. *Cad Saude Publica.* 2009;25(10):2125-38.
- Sibai BM, Hauth J, Caritis S, Lindheimer MD, MacPherson C, Klebanoff M, et al. Hypertensive disorders in twin versus singleton gestations. *Am J Obstet Gynecol.* 2000;182(4):938-42. [http://dx.doi.org/10.1016/S0002-9378\(00\)70350-4](http://dx.doi.org/10.1016/S0002-9378(00)70350-4).
- Golbert A, Campos MAA. Diabetes Mellitus Tipo 1 e Gestação. *Arq Bras Endocrinol Metab.* 2008;52(2):307-14.
- Szylił NA, Segre CA de M, Machado MV. Diabetes e gestação: análise das gestantes submetidas à ecocardiografia fetal em um período de dez anos. *Einstein.* 2008;6(1):42-50.
- Mota M, Melo A, Burak C, Daltro C, Rodrigues B, Lucena R. Antropometria crânica de recém-nascidos normais. *Arq Neuropsiquiatr.* 2004;62(3):626-29. <http://dx.doi.org/10.1590/S0004-282X2004000400011>.

23. Nomura RMY, Martinez NA, Teshima LK, Miyadahira S, Zugaib M. Movimentos respiratórios fetais em gestações com *diabetes mellitus* pré-gestacional. RBCO. 2007;29(7):354-59.
24. Chakraborty SK, Yousuf BMA, Islam S, Banu LA, Khan R, Shamim KM, et al. Impacts of Gestational Diabetes Mellitus on the Mother and the Neonate – A Descriptive Study. Bangladesh J Anatomy. 2010;8(2):64-8.
25. Corrêa RRM, Salge AKM, Ribeiro GA, Ferraz MLF, Reis MA, Castro ECC, et al. Alterações anatomopatológicas da placenta e variações do índice de Apgar. Rev Bras Saúde Matern Infant. 2006;6(2):239-43, <http://dx.doi.org/10.1590/S1519-38292006000200012>.
26. Brasil. Ministério da Saúde. Secretaria de Assistência à Saúde. Gestação de alto risco. 4ªed. Brasília: Divisão Nacional de Saúde Materno-Infantil. 2000.