

Papéis Avulsos de Zoologia

Museu de Zoologia da Universidade de São Paulo

Volume 52(35):423-429, 2012

www.mz.usp.br/publicacoes
<http://portal.revistasusp.sibi.usp.br>
www.scielo.br/paz

ISSN impresso: 0031-1049
ISSN on-line: 1807-0205

RICHNESS OF HYMENOPTEROUS GALLS FROM SOUTH AMERICA

VALÉRIA CID MAIA¹

ABSTRACT

An overview of hymenopterous galls from South America is presented here based on literature as well as on data from the insect gall collection of the Museu Nacional/UFRJ. Seventy-three galls have been recorded on 71 host plants. Myrtaceae and Fabaceae are the plant families with the greatest number of galled species and gall morphotypes. Gall richness per plant species varied from one to two. The stem and bud were the most galled plant organs. The gallers comprise eight hymenopteran families, the best-represented being Eulophidae, Eurytomidae, and Cynipidae. Geographic records are restricted to six countries, and the majority is from Brazil.

KEY-WORDS: Diversity; Geographical distribution; Insect plant interaction.

INTRODUCTION

Galls are predictable and consistent plant deformations that occur in response to feeding or other stimuli by foreign organisms (Gagné, 1994). The gall formation is characterized by abnormal growth of plant tissues by cell hypertrophy and hyperplasia (Price, 2005). Insect galls are considered the most sophisticated herbivore interactions of nature (Shorthouse *et al.*, 2005). Galling insects have the ability to manipulate the development of plant tissue and promote its growth (Stone & Schönrogge, 2003). The development of these structures has resulted from an adaptive strategy of many insects to obtain food and even protection against predators (Stone & Schönrogge, 2003). Among the insects, Diptera, Lepidoptera, Hemiptera, Coleoptera, Hymenoptera and Thysanoptera include galling species (Mani, 1964).

Gall midges (Diptera, Cecidomyiidae) are the most common galling insects in all biogeographic regions (Felt, 1940). In the Neotropical region,

Hemiptera are the second most diversified and frequent gallers, followed by Lepidoptera and Coleoptera, being responsible for about 135, 110 and 80 gall morphotypes (Maia, 2006 and 2012). Thysanoptera are the least frequent ones with only 14 records in the Neotropics (Maia, 2006). Hymenopteran gallers are very frequent in Europe and North America, where records of galls induced by Tenthredinidae (sawflies), Cynipidae (gall wasps), Agaonidae (fig wasps), Braconidae, Eurytomidae, Eulophidae, and others families are found (Felt, 1940; Wehrmaker, 1998; Zinovjev, 1998).

Nevertheless, little is known about the diversity of hymenopterous galls in South America, as data are scattered in several insect gall inventories. There is a single previous compilatory study, published by Houard in 1933, which includes Hymenopterous galls. This author organized a catalog of galls induced by arthropods and nematodes. But even in Houard, 1933, the information about Hymenopteran gallers is pulverized. The present work is the first attempt to gather and upgrade this information.

1. Museu Nacional, Departamento de Entomologia, Quinta da Boa Vista, São Cristóvão, CEP 20940-040, Rio de Janeiro, RJ, Brasil.
E-mail: maiaavid@acd.ufrj.br

This paper aims to answer the following questions: (1) How many hymenopterous galls have been recorded in South America? (2) How many plant families are galled by Hymenoptera? (3) Which plant families are the most galled? (4) Is there a preference for any plant organ? (5) What is known about the taxonomy of these gallers? (6) Which hymenopteran families induce galls in this region? (7) Among them, which are the most common? and (8) What is known about the geographic distribution of these gallers?

MATERIAL AND METHODS

Data on hymenopterous galls were compiled based on data from the insect galls collection of Museu Nacional/UFRJ as well as on literature. The catalog of Houard, 1933 was used as a starting point and reference to previous publications. From 1933 to 2010, the research was based on the Web of Science, focusing upon host plant, galled plant organ, galling wasp and locality. The following papers were examined: Rübssaamen (1899, 1907, 1908), Tavares (1909, 1914), Kieffer & Jörgensen (1910), Brèthes (1915, 1918, 1922), Ferrière (1924), Houard (1933), Gomes (1943), Costa-Lima (1962), Fernandes *et al.* (1988), Arduin *et al.* (1989), Monteiro *et al.* (1993), Kraus *et al.* (1998), Kraus & Tanque (1999), Gonçalves-Alvim & Fernandes (2001), Ronquist & Liljeblad (2001), DalMolin *et al.* (2004), Maia & Fernandes (2004), Urso-Guimarães & Scareli-Santos (2006), Gates & Delvare (2008), Ros-Farré & Pujade-Villar

(2009), Pentead-Dias & Carvalho (2008), Coelho *et al.* (2009), Pereira (2010) and Leite *et al.* (2011).

The Hymenopteran species names were checked in the Universal Chalcidoidea Database (Noyes, 2011). The botanical names and authors were checked in the websites Tropicos® (Tropicos.org) and Angiosperm Phylogeny Group III.

RESULTS AND DISCUSSION

Seventy-three hymenopterous galls have been recorded in South America. These galls occurred on 71 plant species distributed among 21 plant families of angiosperms. No galls were recorded on gymnosperms. The great majority of the host plants (about 90%) belong to dicotyledons. Only two families of monocotyledons were galled: Araceae and Orchidaceae. Myrtaceae and Fabaceae are the plant families with the greatest number of host plant (18 and 16, respectively) and gall morphotypes (18 and 19, respectively). These two families comprise about 50% of the recorded galls (Table 1).

The gall richness per plant species varied little (from one to two). The great majority (97%) presented only one gall morphotype; the only exceptions are *Prosopis alba* Griseb and *P. campestris* Griseb (Fabaceae) with two morphotypes respectively (Table 1). The plant genera with the greatest number of galled species are *Eugenia* L. (Myrtaceae) (09 spp.), *Erythroxylum* P. Browne (Erythroxylaceae) (05 spp.) and *Psidium* L. (Myrtaceae) (04 spp.) (Table 1).

TABLE 1: Distribution of hymenopterous galls from South America per host plant families and species.

Host family (n = 21)	Host plant (n = 71)	Number of Hymenopterous galls (n = 73)
Anacardiaceae	<i>Schinus dependens</i> Ortega	01
	<i>S. weinmanniaefolia</i> Engl.	01
Annonaceae	<i>Duguetia furfuracea</i> (St. Hil.) Benth. & Hook.	01
Araceae	<i>Philodendron selloum</i> C. Koch	01
	<i>P. dubium</i> Chodat & Vischer	01
	<i>P. petraeum</i> Chodat & Vischer	01
	<i>Philodendron</i> sp.	01
Boraginaceae	<i>Cordia curassavica</i> (Jacq.) R. & S.	01
Caryocaraceae	<i>Caryocar brasiliense</i> Camb.	01
Erythroxylaceae	<i>Erythroxylum campestre</i> St. Hill.	01
	<i>E. citrifolium</i> A.St.-Hil.	01
	<i>E. frangulifolium</i> St. Hill.	01
	<i>E. ovalifolium</i> Peyr	01
	<i>E. vacciniifolium</i> Mart.	01
Euphorbiaceae	<i>Colliguaya brasiliensis</i> Müll. Arg.	01
	<i>Colliguaya odorifera</i> Mol.	01
	<i>Sapium</i> sp.	01
	<i>Tragia volubilis</i> L.	01

Continuation of Table 1.

Host family (n = 21)	Host plant (n = 71)	Number of Hymenopterous galls (n = 73)
Fabaceae	<i>Acacia cavenia</i> Bert.	01
	<i>Calliandra bicolor</i> Benth.	01
	<i>Canavalia ensiformis</i> (L.) DC.	01
	<i>Copaifera</i> sp.1	01
	<i>Copaifera</i> sp.2	01
	<i>Copaifera langsdorfii</i> Desf.	01
	<i>Cratylia mollis</i> Mart. ex Benth	01
	<i>Gourliea decorticans</i> Gill.	01
	<i>Inga</i> sp.	01
	<i>Machaerium</i> sp.	01
	<i>Mimosa biuncifera</i> Benth.	01
	<i>Peltogyne</i> sp.	01
	<i>Prosopis alba</i> Griseb	02
	<i>P. alata</i> Phil.	01
	<i>P. campestris</i> Griseb	02
	<i>Swartzia</i> sp.	01
	Fagaceae	<i>Nothofagus antarctica</i> Oerst
Lauraceae	<i>Nectandra lanceolata</i> Ness	01
	<i>Ocotea opifera</i> Mart	01
Loranthaceae	<i>Sruthanthus vulgaris</i> Mart.	01
Malpighiaceae	<i>Byrsonima coccolobifolia</i> H.B. & K.	01
Moraceae	<i>Ficus doliaria</i> Mart.	01
	<i>Ficus noronhae</i> Oliver	01
	<i>Ficus</i> sp.	01
Myrsinaceae	<i>Cybianthus</i> sp.	01
Myrtaceae	<i>Blepharocalyx salicifolium</i> (Kunth) O. Berg.	01
	<i>Corymbia citriadora</i> Hill & Johnson (introduced species)	01
	<i>Eucalyptus camaldulensis</i> Dehn. (introduced species)	01
	<i>Eugenia acuminatissima</i> Berg.	01
	<i>E. axillaris</i> (Willd.)	01
	<i>E. copacabanensis</i> Kiaersk	01
	<i>E. jaboticaba</i> (Vell.) Kiaersk	01
	<i>E. ovalifolia</i> Camb.	01
	<i>E. puniceifolia</i> (H.B. e K.) DC	01
	<i>Eugenia</i> sp.1	01
	<i>Eugenia</i> sp.2	01
	<i>Eugenia</i> sp.3	01
	<i>Myrcia torta</i> DC.	01
	<i>Neomitranthes obscura</i> (DC) N.J.E. Silveira	01
	<i>Psidium cattleianum</i> Sab.	01
	<i>P. cinereum</i> Mart. ex DC	01
	<i>P. guayava</i> L.	01
<i>Psidium</i> sp.	01	
Nyctaginaceae	<i>Guapira opposita</i> (Vell.) Reitz.	01
Orchidaceae	<i>Cattleya gutata</i> Lindl.	01
	<i>Cattleya</i> sp.	01
	<i>Laelia</i> sp.	01
Rhamnaceae	<i>Scutia buxifolia</i> Reiss.	01
Sapindaceae	<i>Paullinia elegans</i> Camb.	01
	<i>Serjania</i> sp.	01
	<i>Urvillea uniloba</i> Radlk.	01
Vochysiaceae	<i>Qualea parvifolia</i> Mart.	01
Winteraceae	<i>Drymis brasiliensis</i> Miers.	01




FIGURE 1: Distribution of Hymenopteran galls from South America per plant organ.

The galls were recorded on several plant organs, such as stem, bud, leaf, fruit, flower and aerial root. Stem and bud were the most galled plant organs (around 48%), and flower and aerial root were the least galled (around 9.4%) (Figure 1).

The taxonomy of the galling wasps is still incipient. Only 28 galls (about 38%) are identified at species level (Tables 2 and 3). The majority of the records are in wider categories. The galls comprise eight hymenopteran families (Agaonidae, Cynipidae, Eulophidae, Eurytomidae, Figitidae, Pteromalidae, Scelionidae, and Tanaostigmatidae), the best-represented being Eulophidae, Eurytomidae, and Cynipidae with 13, 08 and 06 species, respectively. They also comprise 14 genera: *Aditrochus* Rübsaamen, 1902, *Chrysocharis* Förster, 1856, *Eurytoma* Illiger, 1807, *Neohyperteles* De Santis, 1957, *Proseurytoma* Kieffer, 1910, *Tanaostigma* Howard, 1890, and *Tetrastichus* Haliday, 1844 (all represented by only one species), *Minapis* Brèthes, 1916, and *Tanaostigmodes* Ashmead, 1896 (each represented by two species), *Eschatocerus* Mayr, 1881, *Myrtopsen* Rübsaamen, 1908, *Prodecatoma* Ashmead, 1904, and *Rileyia* Ashmead, 1888 (each represented by three species). *Aprostocetus* Westwood, 1833 the best represented genus, has five galling species in South America (Table 2).

These hymenopteran genera differ in species richness as well as in geographic distribution. *Eurytoma*, *Aprostocetus*, and *Tetrastichus* are very large and cosmopolitan genera, with about 700, 750 and 500 described species (Gates & Delvare, 2008; Noyes, 2011). *Chrysocharis*, *Tanaostigmodes*, *Rileyia*, and *Prodecatoma* are also worldwide, but less diversified with about 140, 65, 65, and 50 known species (Hanson,

1987, Noyes, 2011). The others (*Tanaostigma*, *Myrtopsen*, *Aditrochus*, *Minapis*, *Eschatocerus*, *Neohyperteles*, and *Proseurytoma*) comprise few species (about 11, 09, 03, 03, 02 and 01 species, respectively), and are mainly or exclusively Neotropical genera (Noyes, 2011; Ronquist & Liljeblad, 2001; Ros-Farré & Pujade-Villar, 2009).

Regarding the families of Hymenoptera, Eulophidae are associated with 10 plant families (Araceae, Euphorbiaceae, Myrtaceae, Sapindaceae, Lauraceae, Annonaceae, Boraginaceae, Caryocaraceae, Erythroxylaceae, and Winteraceae), Araceae being the plant family with the greatest number of galled species (04), followed by Euphorbiaceae (03), Lauraceae, Myrtaceae and Sapindaceae (02 each). Only one species of the other plant families was galled. The majority of the Eulophidae galling species was monophagous. Only three were oligophagous: *Chrysocharis gallicola* (Costa-Lima, 1930) which induced galls on two species of *Psidium* L. (Myrtaceae), *Aprostocetus gallicola* (Ferrière, 1924) on four species of *Philodendron* Schott (Araceae), and *A. verticillata* (Brèthes, 1913) on two genera of Euphorbiaceae (Table 2). Eurytomidae are associated with four plant species: Fabaceae, Orchidaceae, Myrtaceae, and Euphorbiaceae, galling two species of each family, except Euphorbiaceae, with a single host plant (Table 2). Cynipidae are associated mainly with Fabaceae, but there is a single species associated with Anacardiaceae. Four species of Cynipidae are monophagous, and two are oligophagous, namely: *Eschatocerus myriadeus* Kieffer & Jörgensen, 1910, and *E. niger* Kieffer & Jörgensen, 1910, both inducing galls on two species of *Prosopis* L., (Fabaceae) (Table 2). Tanaostigmatidae are associated with two plant species: Fabaceae (three galling species) and Rhamnaceae (a single galling species). All Tanaostigmatidae galls are monophagous, except for *Tanaostigma chapadae* (Ashmead, 1904), which induce galls on three genera of Fabaceae.

The South American fauna of hymenopteran galls differ from Holarctic fauna, mainly by the absence of gall forming sawflies (Tenthredinidae) and by the higher diversity of gall-forming Eulophidae, Eurytomidae and Cynipidae. The geographic records are restricted to six countries, and the majority is from Brazil (around 68%; Figure 2), where most of the surveys have been developed.

CONCLUSIONS

Hymenoptera galls are less diversified in South America than in Nearctic and Palearctic regions. The


TABLE 2: Distribution of Hymenopteran gallers from South America per plant, plant organ and country.

Galling family	Galling species	Host plant	Plant organ	Country
Agaonidae (n = 01)	sp.	<i>Ficus</i> sp.	Flower	Brazil
Chalcidoidea (n = 03)	sp.1	<i>Acacia cavenia</i>	Bud	Paraguay
	sp.2	<i>Gourliea decorticans</i>	Bud and stem	Argentina
	sp.3	Not determined	Leaf	French Guyana
Cynipidae (n = 06)	<i>Eschatocerus acaciae</i> Mayr, 1881	<i>Acacia cavenia</i>	Stem	Uruguay
	<i>E. myriadeus</i> Kieffer & Jörgensen, 1910	<i>Prosopis alpacato</i> and <i>P. campestris</i>	Stem Stem	Argentina Argentina
	<i>E. niger</i> Kieffer & Jörgensen, 1910	<i>P. alba</i> and <i>P. campestris</i>	Bud and stem Bud	Argentina Argentina
		sp.1	<i>Schinus weinmanniaefolia</i>	Stem
	sp.2	<i>Peltogyne</i> sp.	Leaf	Brazil
	sp.3	<i>Machaerium</i> sp.	Bud	Brazil
	Eulophidae (n = 13)	<i>Aprostocetus bahiensis</i> (Costa-Lima, 1959)	<i>Ocotea opifera</i>	Fruit
<i>A. colliguayae</i> (Philippi, 1873)		<i>Colliguaya odorifera</i>	No data	Chile
<i>A. bondari</i> (Costa-Lima, 1959)		<i>Paullinia elegans</i>	Fruit	Brazil
<i>A. gallicola</i> (Ferrière, 1924)		<i>Philodendron selloum</i> ,	Flower	Paraguay
		<i>P. petraeum</i> ,	Flower	Paraguay
		<i>P. undulatum</i> ,	Flower	Paraguay
<i>Philodendron</i> sp.		Flower	Paraguay	
<i>A. verticillata</i> (Brèthes, 1913)		<i>Sapium</i> sp. <i>Colliguaya brasiliensis</i>	No data No data	Argentina Brazil
<i>Neohyperteles euplectriformis</i> De Santis, 1957		<i>Nectandra lanceolata</i>	No data	Argentina
<i>Chrysocharis gallicola</i> (Costa-Lima, 1930)		<i>Psidium guayaba</i> and <i>Psidium</i> sp.	No data No data	Brazil Brazil
		<i>Tetratichus urvilleae</i> (De Santis, 1957)	<i>Urvillea uniloba</i>	Fruit
sp.1		<i>Cordia curassavica</i>	Stem	Brazil
sp.2		<i>Erythroxylum frangulifolium</i>	Stem	Brazil
sp.3		<i>Drimys brasiliensis</i>	Stem	Brazil
sp.4		<i>Duguetia furfuracea</i>	Leaf	Brazil
sp.5	<i>Caryocar brasiliense</i>	Leaf	Brazil	
Eurytomidae (n = 08)	<i>Rileya haumani</i> (Brèthes, 1918)	<i>Tragia volubilis</i>	Flower	Argentina
	<i>R. nigra</i> (Gomes, 1943)	<i>Cattleya gutata</i>	Root	Brazil
	<i>R. cearae</i> (Crawford, 1910)	Not determined	No data	Brazil
	<i>Eurytoma orchideadrum</i> (Westwood, 1869)	<i>Cattleya</i> sp.	Stem	Brazil
	<i>Prodecatoma carpophaga</i> DalMolin, Melo & Perioto, 2004	<i>Psidium cattleianum</i>	Fruit	Brazil
	<i>P. parodii</i> Brèthes, 1922	<i>Prosopis alba</i>	Stem	Argentina
	<i>P. petrodoma</i> DalMolin, Melo & Perioto, 2004	<i>Psidium cinereum</i>	Fruit	Brazil
	<i>Proseurytoma gallarum</i> Kieffer, 1910	<i>Gourliea decorticans</i>	Bud	Argentina
	Figitidae (n = 03)	<i>Myrtopsen mayri</i> Rübsaamen, 1907	<i>Eugenia acuminatissima</i> and <i>Eugenia</i> sp.	Leaf Leaf
<i>M. luderwalti</i> Dettmer, 1930		<i>Eugenia jaboticaba</i>	No data	Brazil
<i>M. rodvalhoi</i> Dettmer, 1930		<i>Mimosa biuncifera</i>	No data	Brazil
Pteromalidae (n = 01)	<i>Aditrochus fagicolus</i> Rübsaamen, 1902	<i>Nothofagus antarctica</i>	Leaf	Argentina
Scelionidae (n = 01)	sp.	<i>Struthanthus vulgaris</i>	Leaf	Brazil
Tanaostigmatidae (n = 05)	<i>Minapis maculiventris</i> Gomes, 1941	<i>Inga</i> sp.	Leaf	Brazil
	<i>M. nigra</i> Brèthes, 1916	<i>Scutia baccifolia</i>	Leaf	Brazil
	<i>Tanaostigma chapadae</i> (Ashmead, 1904)	<i>Canavalia ensiformis</i> ,	Seed	Brazil
		<i>Cratylia mollis</i> and <i>Inga</i> sp.	Seed Seed	Brazil Brazil
		<i>Tanaostigmodes ringueleti</i> (Brèthes, 1924)	<i>Calliandra bicolor</i>	Bud
	<i>T. kiefferi</i> (Mayr, 1905)	Not determined	No data	Paraguay

TABLE 3: Distribution of Hymenopteran galls (records at order category) from South America per plant, plant organ and country.

Host Plant		N° of galls (n = 22)	Plant organ	Country
Family	Species			
Anacardiaceae	<i>Schinus dependens</i>	01	Bud	Brazil
Erythroxylaceae	<i>Erythroxylum campestre</i>	01	Bud	Brazil
Fabaceae	<i>E. ovalifolium</i>	01	Leaf	Brazil
	<i>Copaifera langsdorfii</i>	01	Stem	Brazil
	<i>Copaifera</i> sp.1	01	Bud	Brazil
	<i>Copaifera</i> sp.2	01	Leaf	Brazil
Malpighiaceae	<i>Swartzia</i> sp.	01	Bud	Brazil
	<i>Byrsonima coccolobifolia</i>	01	Bud	Brazil
	<i>Ficus doliaria</i>	01	Bud	Brazil
Moraceae	<i>F. noronhae</i>	01	Fruit	Brazil
	<i>Ficus</i> sp.	01	Leaf	Brazil
	<i>Ficus</i> sp.	01	Fruit	Brazil
Myrsinaceae	<i>Cybianthus</i> sp.	01	Bud	Brazil
Myrtaceae	<i>Eugenia copacabanensis</i>	01	Bud	Brazil
	<i>E. ovalifolia</i>	01	Stem	Brazil
	<i>Eugenia</i> sp.1	01	Stem	Brazil
	<i>Eugenia</i> sp.2	01	Bud	Brazil
	<i>Myrcia torta</i>	01	Bud	Brazil
	<i>Neomitranthes obscura</i>	01	Bud	Brazil
Nyctaginaceae	<i>Guapira opposita</i>	01	Stem	Brazil
Sapindaceae	<i>Serjania</i> sp.	01	Stem	No data
Vochysiaceae	<i>Qualea parvifolia</i>	01	Stem	Brazil

galls are represented by eight families. The most common of these are Eulophidae, Eurytomidae, and Cynipidae. There is no record of Tenthredinidae galls. Hymenopteran galls were more frequent on stems and buds. The taxonomy of the galling species is still incipient.

**FIGURE 2:** Distribution of the records of Hymenopteran galls from South America per country.

RESUMO

Uma abordagem geral de galhas de himenópteros da América do Sul é apresentada aqui com base na literatura e na coleção de galhas de insetos do Museu Nacional/ UFRJ. Setenta e três galhas estão registradas em 71 espécies de plantas hospedeiras. Myrtaceae e Fabaceae são as famílias de planta com maior número de espécies hospedeiras e de morfotipos de galhas. A riqueza de galhas por espécie botânica variou de um a dois. Caule e gema foram os órgãos da planta mais atacados. Os galhadores compreendem oito famílias de Hymenoptera, tendo Eulophidae, Eurytomidae, e Cynipidae maior representatividade. Os registros geográficos estão restritos a seis países, sendo a maioria do Brasil.

PALAVRAS-CHAVE: Diversidade; Distribuição geográfica; Interação inseto-planta.

ACKNOWLEDGMENTS

I am grateful to CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico) for financial support (Proc. 372060/2011-0).

REFERENCES

- ARDUIN, M.; KRAUS, J.E.; OTTO, P.A. & VENTURELLI, M. 1989. Caracterização morfológica e biométrica de galhas foliares em *Struthanthus vulgaris* Mart. (Loranthaceae). *Revista Brasileira de Biologia*, 49:817-823.
- BRÈTHES, J. 1915. Hyménoptères parasites de l'Amérique Méridionale. *Anales del Museo Nacional de Historia Natural*, Buenos Aires, 27:401-430.
- BRÈTHES, J. 1918. Description d'un Chalcidien gallicole de la République Argentina. *Bulletin de La Société Entomologique de France*, (3):82-84.
- BRÈTHES, J. 1922. Himenopteros y Dípteros de varias procedências. *Anales de La Sociedad Científica Argentina*, 93:119-146.
- COELHO, M.S.; ALMADA, E.D.; FERNANDES, G.W.; CARNEIRO, M.A.A.; SANTOS, R.M. DOS; QUINTINO, A.V. & SANCHEZ-AZOFEIFA, A. 2009. Gall inducing arthropods from a seasonally dry tropical forest in Serra do Cipó, Brazil. *Revista Brasileira de Entomologia*, 53:404-414.
- COSTA-LIMA, A. 1962. *Insetos do Brasil. Himenópteros*. Escola Nacional de Agronomia, Rio de Janeiro. v. 12, 2. pt. 393 pp. (série didática, 14)
- DALMOLIN, A.; MELO, G.A.R. & PERIOTO, N.W. 2004. Novas espécies de *Prodecatoma* (Hymenoptera, Eurytomidae) associadas a galhas em frutos de duas espécies de *Psidium* L. (Myrtaceae), com comentários sobre *Prodecatoma spermophaga* Costa-Lima. *Revista Brasileira de Entomologia*, 48(4):519-528.
- FELT, E.P. 1940. *Plant galls and gall makers*. Ithaca, N.Y. 364 p.
- FERNANDES, G.W.; TAMEIRÃO-NETO, E. & MARTINS, R.P. 1988. Ocorrência e caracterização das galhas entomógenas na vegetação do Campus Pampulha da Universidade Federal de Minas Gerais. *Revista Brasileira de Zoologia*, 5:11-29.
- FERRIÈRE, C. 1924. Note sur deux nouveaux Chalcidiens phytophages du Paraguay, suivie d'une étude sur la structure de leur tarière. *Annales de La Société de Entomologie de France*, 93:1-21.
- GAGNÉ, R.J. 1994. *The gall midges of the Neotropical region*. Cornell University Press, Ithaca. 352 p.
- GATES, M. & DELVARE, G. 2008. A new species of *Eurytoma* (Hymenoptera: Eurytomidae) attacking *Quadrastichus* spp. (Hymenoptera: Eulophidae) galling *Erythrina* spp. (Fabaceae), with a summary of the African *Eurytoma* biology and species checklist. *Zootaxa*, 1751:1-24.
- GOMES, J.G. 1943. Um novo euritomídeo em orquídea (Hymenoptera, Chalcidoidea). *Boletim da Sociedade Brasileira de Agronomia*, 6:244-246.
- GONÇALVES-ALVIM, S.J. & FERNANDES, G.W. 2001. Comunidades de insetos galhadores (Insecta) em diferentes fisionomias do cerrado em Minas Gerais, Brasil. *Revista Brasileira de Zoologia*, 18:289-305.
- HANSON, C. 1987. Revision of the New World species of *Chrysocharis* Förster (Hymenoptera: Eulophidae). *Entomologica Scandinavia*, (Suppl. 31):1-86 p.
- HOUARD, C. 1933. *Les Zoocécidies des Plantes de L'Amérique du Sud et de L'Amérique Centrale*. Hermann et Cie, Paris. 549 p.
- KIEFFER, J.J. & JÖRGENSEN, P. 1910. Gallen und gallentiere aus Argentinien. *Zentralblatt für Bakteriologie und Parasitenkunde*, 2:362-444.
- KRAUS, J.E. & TANQUE, M. 1999. Morpho-ontogenetic aspects of entomogenous galls in roots of *Cattleya guttata* (Orchidaceae). *Lindleyana*, 14(4):204-213.
- KRAUS, J.E.; SOLORZANO-FILHO, J.A.; ARDUIN, M. & ISAIAS, R.M.S. 1998. Respostas morfogênicas de plantas brasileiras a insetos galhadores. *Monographs in Systematic Botany from the Missouri Botanical Garden*, 68:345-354.
- LEITE, G.L.D.; D'ÁVILA, V. DE A.; CERQUEIRA, V.M.; NASCIMENTO, A.F. DO & FERNANDES, G.W. 2011. Spatial distribution of a spherical gall (Hymenoptera, Eulophidae) on *Caryocar brasiliense* (Caryocaraceae). *Revista Brasileira de Entomologia*, 55:396-400.
- MAIA, V.C. 2006. Galls of Hemiptera, Lepidoptera and Thysanoptera from Central and South America. *Publicações Avulsas do Museu Nacional*, 110:3-22.
- MAIA, V.C. 2012. Coleopterous galls from the Neotropical region. *Papéis Avulsos de Zoologia*, 52(15):175-184.
- MAIA, V.C. & FERNANDES, G.W. 2004. Insect galls from Serra de São José (Tiradentes, MG, Brazil). *Brazilian Journal of Biology*, 64(3A):423-445.
- MANI, M.S. 1964. *Ecology of plant galls*. Junk, The Hague. 434 p.
- MONTIÉRO, R.F.; FERRAZ, F.F.; MAIA V.C. & AZEVEDO, M.A.P. 1993. Galhas entomógenas em restinga: uma abordagem preliminar. In: Simpósio de Ecossistemas da Costa Brasileira, 3º. Anais. São Paulo, ACIESP. p. 210-220.
- NOYES, J.S. 2011. Universal Chalcidoidea Database. World Wide Web electronic publication. Available at: <www.nhm.ac.uk/chalcidoidea>. Access in: 11/11/2011.
- PENTEADO-DIAS, A.M. & CARVALHO, F.M. 2008. New species of Hymenoptera associated with galls on *Calliandra brevipes* Benth. (Fabaceae, Mimosoidea) in Brazil. *Revista Brasileira de Entomologia*, 52:305-310.
- PEREIRA, R.A. 2010. *Aspectos morfo-bioecológicos de Epichrysocharis burwelli* (Eulophidae, Hymenoptera), *vespas-das-galhas das folhas de Corymbia citriodora*. (Tese de Doutorado). Escola Superior de Agricultura "Luiz de Queiroz", Piracicaba, SP. 71 p.
- PRICE, P.W. 2005. Adaptive radiation of gall-inducing insects. *Basic Applied Ecology*, 6(5):413-421.
- RONQUIST, F. & LILJEBLAD, J. 2001. Evolution of the gall wasp-host plant association. *Evolution*, 55:2503-2522.
- ROS-FARRÉ, P. & PUJADE-VILLAR, J. 2009. Revisión del género *Myrtopsen* Rübсаamen, 1908 (Hymenoptera: Figitidae: Thrasorinae). *Dugesiana*, 16:21-33.
- RÜBSAAMEN, E.H. 1899. Mittheilungen über neue und bekannte Gallen aus Europa, Asien, Afrika und Amerika. *Entomologische Nachrichtenblatt*, 25:225-282.
- RÜBSAAMEN, E.H. 1907. Beiträge zur Kenntniss ausereuropäischer Zoocécidien. III Beitrag: Gallen aus Brasilien und Peru. *Marcellia*, 6:110-173.
- RÜBSAAMEN, E.H. 1908. Beiträge zur Kenntniss ausereuropäischer Zoocécidien. III Beitrag: Gallen aus Brasilien und Peru. *Marcellia*, 7:15-79.
- SHORTHOUSE, J.D.; WOOL, D. & RAMAN, A. 2005. Gall-inducing insects – Nature's most sophisticated herbivores. *Basic Applied Ecology*, 6(5):407-411.
- STONE, G.N. & SCHÖNROGGE, K. 2003. The adaptive significance of insect gall morphology. *Trends in Ecology and Evolution*, 18(10):512-522.
- TAVARES, J.S. 1909. Contributio primo ad cognitionem Cecidologiae Braziliae. *Brotéria*, 8:5-29.
- TAVARES, J.S. 1914. Le Goyavier (*Psidium guyava* Radd.) au Brésil. *Brotéria*, 12:148-151.
- TROPICOS.ORG [BOTANICAL NAMES AND AUTHORS] MISSOURI BOTANICAL GARDEN. 2011. Available at: <www.tropicos.org>. Access in: 11/11/2011.
- URSO-GUIMARÃES, M.V. & SCARELI-SANTOS, C. 2006. Galls and Gall makers in plants from the Pé-de-Gigante Cerrado Reserve, Santa Rita do Passa Quatro, SP, Brazil. *Brazilian Journal of Biology*, 66:357-369.
- WEHRMAKER, A. 1998. On the beginnings of cecidology in 19th century North America: Bassett's discovery of heterogony in oak gall wasps (Hymenoptera: Cynipidae). In: Csóka, G.; Mattsom, W.J.; Stone, G.N. & Price, P.W. (Eds.) *The Biology of Gall-Inducing Arthropods*. Minnesota, Department of Agriculture, North Central Forest Experiment Station. p. 106-110.
- ZINOVJEV, A.G. 1998. Palearctic sawflies of the genus *Pontania* Costa (Hymenoptera: Tenthredinidae) and their host-plant specificity. In: Csóka, G.; Mattsom, W.J.; Stone, G.N. & Price, P.W. (Eds.) *The Biology of Gall-Inducing Arthropods*. Minnesota, Department of Agriculture, North Central Forest Experiment Station. p. 204-225.

Aceito em: 23.10.2012

Publicado em: 20.12.2012



Publicado com o apoio financeiro do Programa de Apoio às Publicações Científicas Periódicas da USP

EDITORIAL COMMITTEE

Publisher: Museu de Zoologia da Universidade de São Paulo. Avenida Nazaré, 481, Ipiranga, CEP 04263-000, São Paulo, SP, Brasil.

Editor-in-Chief: Carlos José Einicker Lamas, Serviço de Invertebrados, Museu de Zoologia, Universidade de São Paulo, Caixa Postal 42.494, CEP 04218-970, São Paulo, SP, Brasil. E-mail: editormz@usp.br.

Associate Editors: Mário César Cardoso de Pinna (*Museu de Zoologia, Universidade de São Paulo, Brasil*); Luis Fábio Silveira (*Museu de Zoologia, Universidade de São Paulo, Brasil*); Marcos Domingos Siqueira Tavares (*Museu de Zoologia, Universidade de São Paulo, Brasil*); Sérgio Antonio Vanin (*Instituto de Biociências, Universidade de São Paulo, Brasil*); Hussam El Dine Zaher (*Museu de Zoologia, Universidade de São Paulo, Brasil*).

Editorial Board: Rüdiger Bieler (*Field Museum of Natural History, U.S.A.*); Walter Antonio Pereira Boeger (*Universidade Federal do Paraná, Brasil*); Carlos Roberto Ferreira Brandão

(*Universidade de São Paulo, Brasil*); James M. Carpenter (*American Museum of Natural History, U.S.A.*); Ricardo Macedo Corrêa e Castro (*Universidade de São Paulo, Brasil*); Mario de Vivo (*Universidade de São Paulo, Brasil*); Marcos André Raposo Ferreira (*Museu Nacional, Rio de Janeiro, Brasil*); Darrel R. Frost (*American Museum of Natural History, U.S.A.*); William R. Heyer (*National Museum of Natural History, U.S.A.*); Ralph W. Holzenthal (*University of Minnesota, U.S.A.*); Adriano Brilhante Kury (*Museu Nacional, Rio de Janeiro, Brasil*); Gerardo Lamas (*Museo de Historia Natural "Javier Prado", Lima, Peru*); John G. Maisey (*American Museum of Natural History, U.S.A.*); Naércio Aquino Menezes (*Universidade de São Paulo, Brasil*); Christian de Muizon (*Muséum National d'Histoire Naturelle, Paris, France*); Nelson Papavero (*Universidade de São Paulo, Brasil*); James L. Patton (*University of California, Berkeley, U.S.A.*); Richard O. Prum (*University of Kansas, U.S.A.*); Olivier Rieppel (*Field Museum of Natural History, U.S.A.*); Miguel Trefaut Urbano Rodrigues (*Universidade de São Paulo, Brasil*); Randall T. Schuh (*American Museum of Natural History, U.S.A.*); Ubirajara Ribeiro Martins de Souza (*Universidade de São Paulo, Brasil*); Paulo Emilio Vanzolini (*Universidade de São Paulo, Brasil*); Richard P. Vari (*National Museum of Natural History, U.S.A.*).

INSTRUCTIONS TO AUTHORS - (April 2007)

General Information: *Papéis Avulsos de Zoologia (PAZ)* and *Arquivos de Zoologia (AZ)* cover primarily the fields of Zoology, publishing original contributions in systematics, paleontology, evolutionary biology, ontogeny, faunistic studies, and biogeography. *Papéis Avulsos de Zoologia* and *Arquivos de Zoologia* also encourage submission of theoretical and empirical studies that explore principles and methods of systematics.

All contributions must follow the International Code of Zoological Nomenclature. Relevant specimens should be properly curated and deposited in a recognized public or private, non-profit institution. Tissue samples should be referred to their voucher specimens and all nucleotide sequence data (aligned as well as unaligned) should be submitted to GenBank (www.ncbi.nih.gov/Genbank) or EMBL (www.ebi.ac.uk).

Peer Review: All submissions to *Papéis Avulsos de Zoologia* and *Arquivos de Zoologia* are subject to review by at least two referees and the Editor-in-Chief. All authors will be notified of submission date. Authors may suggest potential reviewers. Communications regarding acceptance or rejection of manuscripts are made through electronic correspondence with the first or corresponding author only. Once a manuscript is accepted providing changes suggested by the referees, the author is requested to return a revised version incorporating those changes (or a detailed explanation of why reviewer's suggestions were not followed) within fifteen days upon receiving the communication by the editor.

Proofs: Page-proofs with the revised version will be sent to e-mail the first or corresponding author. Page-proofs *must be returned to the editor, preferentially within 48 hours*. Failure to return the proof promptly may be interpreted as approval with no changes and/or may delay publication. Only necessary corrections in proof will be permitted. Once page proof is sent to the author, further alterations and/or significant additions of text are permitted only at the author's expense or in the form of a brief appendix (note added in proof).

Submission of Manuscripts: Manuscripts should be sent to the **SciELO Submission** (<http://submission.scielo.br/index.php/paz/login>), along with a submission letter explaining the importance and originality of the study. Address and e-mail of the corresponding author must be always updated since it will be used to send the 50 reprints in titled by the authors. Figures, tables and graphics **should not** be inserted in the text. Figures and graphics should be sent in separate files with the following formats: ".JPG" and ".TIF" for figures, and ".XLS" and ".CDR" for graphics, with 300 DPI of minimum resolution. Tables should be placed at the end of the manuscript.

Manuscripts are considered on the understanding that they have not been published or will not appear elsewhere in substantially the same or abbreviated form. The criteria for acceptance of articles are: quality and relevance of research, clarity of text, and compliance with the guidelines for manuscript preparation.

Manuscripts should be written preferentially in English, but texts in Portuguese or Spanish will also be considered. Studies with a broad coverage are encouraged to be submitted in English. All manuscripts should include an abstract and key-words in English and a second abstract and key-words in Portuguese or Spanish.

Authors are requested to pay attention to the instructions concerning the preparation of the manuscripts. Close adherence to the guidelines will expedite processing of the manuscript.

Manuscript Form: Manuscripts should not exceed 150 pages of double-spaced, justified text, with size 12 and source Times New Roman (except for symbols). Page format should be A4 (21 by 29.7 cm), with 3 cm of margins. The pages of the manuscript should be numbered consecutively.

The text should be arranged in the following order: **Title Page, Abstracts with Key-Words, Body of Text, Literature Cited, Tables, Appendices, and Figure Captions**. Each of these sections should begin on a new page.

(1) **Title Page:** This should include the **Title, Short Title, Author(s) Name(s) and Institutions**. The title should be concise and, where appropriate, should include mention of families and/or higher taxa. Names of new taxa should not be included in titles.

(2) **Abstract:** All papers should have an abstract in **English** and another in **Portuguese or Spanish**. The abstract is of great importance as it may be reproduced elsewhere. It should be in a form intelligible if published alone and should summarize the main facts, ideas, and conclusions of the article. Telegraphic abstracts are strongly discouraged. Include all new taxonomic names for referencing purposes. Abbreviations should be avoided. It should not include references. Abstracts and key-words should not exceed 350 and 5 words, respectively.

(3) **Body of Text:** The main body of the text should include the following sections: **Introduction, Material and Methods, Results, Discussion, Conclusion, Acknowledgments, and References at end**. Primary headings in the text should be in capital letters, in bold and centered. Secondary headings should be in capital and lower case letters, in bold and centered. Tertiary headings should be in capital and lower case letters, in bold and indented at left. In all the cases the text should begin in the following line.

(4) **Literature Cited:** Citations in the text should be given as: Silva (1998) *or* Silva (1998:14-20) *or* Silva (1998: figs. 1, 2) *or* Silva (1998a, b) *or* Silva & Oliveira (1998) *or* (Silva, 1998) *or* (Rangel, 1890; Silva & Oliveira, 1998a, b; Adams, 2000) *or* (Silva, *pers. com.*) *or* (Silva *et al.*, 1998), the latter when the paper has three or more authors. The reference need not be cited when authors and date are given only as authority for a taxonomic name.

(5) **References:** The literature cited should be arranged strictly alphabetically and given in the following format:

- **Journal Article** - Author(s). Year. Article title. *Journal name*, volume: initial page-final page. Names of journals must be spelled out in full.
- **Books** - Author(s). Year. *Book title*. Publisher, Place.
- **Chapters of Books** - Author(s). Year. Chapter title. *In: Author(s) ou Editor(s), Book title*. Publisher, Place, volume, initial page-final page.
- **Dissertations and Theses** - Author(s). Year. *Dissertation title*. (Ph.D. Dissertation). University, Place.
- **Electronic Publications** - Author(s). Year. *Title*. Available at: <electronic address>. Access in: date.

Tables: All tables must be numbered in the same sequence in which they appear in text. Authors are encouraged to indicate where the tables should be placed in the text. They should be comprehensible without reference to the text. Tables should be formatted with vertical (portrait), not horizontal (landscape), rules. In the text, tables should be referred as Table 1, Tables 2 and 4, Tables 2-6. Use "TABLE" in the table heading.

Illustrations: Figures should be numbered consecutively, in the same sequence that they appear in the text. Each illustration of a composite figure should be identified by capital letters and referred in the text as: Fig. 1A, Fig. 1B, for example. When possible, letters should be placed in the left lower corner of each illustration of a composite figure. Hand-written lettering on illustrations is unacceptable. Figures should be mounted in order to minimize blank areas between each illustration. Black and white or color photographs should be digitized in high resolution (300 DPI at least). Use "Fig(s)," for referring to figures in the text, but "FIGURE(S)" in the figure captions and "fig(s)," when referring to figures in another paper.

Responsibility: Scientific content and opinions expressed in this publication are sole responsibility of the respective authors.
Copyrights: The journals *Papéis Avulsos de Zoologia* and *Arquivos de Zoologia* are licensed under a Creative Commons Licence (<http://creativecommons.org>).

For other details of manuscript preparation of format, consult the CBE Style Manual, available from the Council of Science Editors (www.councilscienceeditors.org/publications/style).
Papéis Avulsos de Zoologia and *Arquivos de Zoologia* are publications of the Museu de Zoologia da Universidade de São Paulo (www.mz.usp.br). Always consult the Instructions to Authors printed in the last issue or in the electronic home pages: www.scielo.br/paz or www.mz.usp.br/publicacoes.