

Gall-inducing insects of restinga areas (Atlantic Forest) in Brazil: economic importance

Valéria Cid Maia¹

¹ Universidade Federal do Rio de Janeiro (UFRJ), Museu Nacional. Rio de Janeiro, RJ, Brasil.
ORCID: 0000-0001-9396-5618. E-mail: maiavcid@acd.ufrj.br

Abstract. Many gall-inducing insects have been reported as pests in the Old World and North America, although few such examples are known from South America. A list of gall-inducing insects of potential economic importance, with a focus on those of restinga environments, was compiled using Maia (2013a) as starting point and updated with the database “Thompson ISI”, using ‘Insect (title) and gall (topic)’ as keywords. Botanical names were updated using the site “Flora do Brasil, 2020”, while potential economic significance of host plant species was acquired from Santos *et al.* (2009) and the site “Useful Tropical Plants”. Fifty-eight galling species were associated with 29 economically important plant species of 18 families in Brazilian restingas. The gallers were found to belong to Diptera (Cecidomyiidae and Agromyzidae) and Hemiptera (Eriococcidae and Psyllidae), among which Cecidomyiidae were the most important, with 55 gall-inducing species distributed among 28 genera. Six of the found genera are endemic to the Atlantic Forest and, until now, have been exclusively reported in restingas. About 78% of the gallers have been recorded only in Southeast Brazil and about 64% only in the state of Rio de Janeiro. Most gallers were found to be mainly associated with edible and/or medicinal plant species. Data on natural enemies are scarce with most records having been published at the taxonomic level of family or genus. Natural enemies were found associated with 43 gall-inducing species and included parasitoids, predators and inquilines. The first were the most diverse, being represented by 13 hymenopteran families, but the impacts of all of these guilds on galler populations are poorly known. Although 58 gall-inducing species were identified in the present study, the number of insect galls associated with plants of economic interests in restinga environments is about three times greater, since a total of 186 gall morphotypes have been reported. Nonetheless, many gallers are still undetermined, thus revealing how deficient their taxonomical knowledge remains.

Key-Words. Cecidomyiidae; Diptera; Hemiptera; Natural enemies; Geographic distribution.

INTRODUCTION

Galls induced by insects are pathological neoplasms on a plant (Mamaev & Krivosheina, 1993), which provide offspring with nutritionally better food and protection against extreme environmental conditions and several natural enemies (Price *et al.*, 1987; Fernandes & Price, 1992). Although gallers benefit from this relationship, host plants are negatively impacted (Price *et al.*, 1986; Guimarães *et al.*, 2014). Gall induction reduces leaf-water potential and photosynthetic and transpiration rates. It also affects mineral concentrations, such as that of boron, chloride, magnesium, and zinc (Florentine *et al.*, 2005). Physiologically, galls may act as nutrient sinks for their host plants, because nutrients are redirected from ungalled tissues and neighboring plant parts to the gall itself (McCrea *et al.* 1985; Larson & Whitharn, 1991; Marini-Filho & Fernandes, 2012). Furthermore, many studies have shown gall formation to cause reduced flower, fruit, seed, and biomass production, suggesting that galling insects have the potential to reduce fitness of susceptible host plants (Sacchi *et al.*, 1988; Fernandes

& Ribeiro, 1990; Souza *et al.*, 1998). In this sense, cecidogenous insects are considered parasites (Larson, 1998).

Several plants of economic interest are vulnerable to damage by gall-inducing insects, including ornamental (*e.g.*, *Laelia* spp., Orchidaceae), edible (*e.g.*, *Manihot* spp., Euphorbiaceae), medicinal and pesticidal plants (*e.g.*, *Melissa officinalis* L., Lamiaceae, as well as species used in carpentry, cosmetics, cabinet making, and agroforestry (Gagné & Jaschhof, 2017). However, the most well-known galling pests are those associated with edible plants, such as *Apiomyia bergentammii* (Wachtl, 1882) (Diptera, Cecidomyiidae), a pear pest (*Pyrus communis* L.) (Rosaceae) in Europe; *Orseolia oryzae* Wood-Mason, 1889 (Cotes, 1889) (Diptera, Cecidomyiidae), a major endemic rice pest throughout Asia (Kalode & Bentur, 1989); *Lasioptera rubi* Heeger, 1851 (Diptera, Cecidomyiidae), a blackberry, raspberry, dewberry (*Rubus* spp.) (Rosaceae) pest in Europe; and *Ficiomyia perarticulata* Felt, 1922 (Diptera, Cecidomyiidae), a fig pest in the USA (Barnes, 1948). Most galling pests have been reported from the Old World and North America,

whereas few examples are known from South America, such as *latrophobia brasiliensis* (Rübsaamen, 1908a) (Diptera, Cecidomyiidae), a manioc pest (EMBRATER/CIAT, 1982).

Due to the negative impact that gall-inducing insects have on their hosts they can be employed as biological control agents of invasive plants. For example, *Spurgia esulae* Gagné, 1990 (Cecidomyiidae, Diptera), *Kochiomyia stackelbergi* (Mamaev, 1972) (Cecidomyiidae, Diptera), and *Urophora cardui* (Linnaeus, 1758) (Tephritidae, Diptera) have been used to control *Euphorbia esula* L. (Euphorbiaceae) in the USA (Hansen *et al.*, 1997), *Salsola tragus* L. (Chenopodiaceae) in the USA and Uzbekistan (Sobhian *et al.*, 2003), and *Cirsium arvense* (L.) Scop. (Asteraceae) in Canada (Peschken & Harris, 1975), respectively.

Although several cecidogenous species have been associated with plants of economic interest in Brazil, only *latrophobia brasiliensis* Rübsaamen, 1915 has been recognized as a pest (EMBRATER/CIAT, 1982). Nevertheless, there are other examples, such as *Clinodiplosis* sp. on panicles of hog plum, *Anacardium mombium* L. (Anacardiaceae) (Moura *et al.*, 2010), and *Contarinia* sp. on leaves of the cashew tree, *Anacardium occidentale* L. (Anacardiaceae) (Tavares *et al.*, 2011), but these species are still unidentified.

Several insect gall inventories in Brazil have been carried out in areas of restinga (Bregonci *et al.*, 2010; Carvalho-Fernandes *et al.*, 2016; Maia, 2001a, 2013a; Maia & Oliveira, 2010; Maia & Silva, 2016; Maia *et al.*, 2008; Monteiro *et al.*, 1994, 2004; Oliveira & Maia, 2005; Rodrigues *et al.*, 2014), one of the most threatened physiognomies of the Atlantic Forest (MMA, 2018). These inventories have recorded great richness of insect galls, but none have focused on their economic importance. The restinga consists of a mosaic of plant communities (Rizzini, 1997) that grow on sandy and chemically-poor soils, with sea spray as the main source of nutrients (Araújo & Lacerda, 1987). They comprise several plant species of local or widespread economic importance, and so the galls that are associated with them are also of economic significance, since they are generally species-specific and can cause severe damage to their hosts. Since the restinga is the best investigated Brazilian physiognomy with respect to insect galls, it is an ideal ecosystem for studying the economic significance of galling insects.

Thus, the objectives of the present study were to compile a list of gall-inducing insects in restingas that have potential economic significance, and to present the current state of knowledge regarding their biology and geographical distribution.

MATERIAL AND METHODS

A comprehensive list of gall-inducing insects reported for Brazilian restingas was compiled starting with Maia (2013a). More recent data were then obtained by searching the Thompson ISI database for papers pub-

lished from 2013 through 2017 using 'Insect (title) and gall (topic)' as key-words (performed November 2017). Among the resulting papers, those with a focus on restingas were verified. Botanical names were updated using the site "Flora do Brasil, 2020" and a list of synonyms was developed to aid in the search of host plants, while plant common names were obtained from Zanith & Scarano (2004). The potential economic significance of host plant species was acquired from Santos *et al.* (2009) and the site Useful Tropical Plants (2014). Data on galls, gall morphology, natural enemies and geographical distributions were also compiled.

RESULTS AND DISCUSSION

Fifty-eight (58) gall-inducing species were found in association with 29 economically important plant species of 18 families in Brazilian restingas (Table 1). Since these galls are species-specific (Carneiro *et al.*, 2009), they are considered economically important, at least potentially. The inducers found belong to Diptera (n = 56) and Hemiptera (2), with each order being represented by two families: the former by Agromyzidae (n = 1) and Cecidomyiidae (n = 55), and the latter by Eriococcidae and Psyllidae (Table 1).

The economically important Cecidomyiidae found in the present study are distributed among 28 genera, of which seven are wide spread (*Asphondylia* Loew, 1850, *Bruggmanniella* Tavares, 1909, *Clinodiplosis* Kieffer, 1894, *Contarinia* Rondani, 1860, *Dasineura* Rondani, 1840, *Lopesia* Rübsaamen, 1908b, and *Schizomyia* Kieffer, 1889), three occur in more than one zoogeographic region (*Youngomyia* Felt, 1908 in the Nearctic, Neotropical and Oriental regions; and *Neolasioptera* Felt, 1908 and *Stephomyia* Tavares, 1916 in the Nearctic and Neotropical regions) and 18 are restricted to the Neotropics (*Alycaulus* Rübsaamen, 1915, *Arrabiadaeamyia* Maia, 2001b, *Clusiamyia* Maia, 1997, *Cordiomyia* Maia, 1996a, *Costadiplosis*, Viceconte & Maia, 2009, *Dactylodiplosis* Rübsaamen, 1915, *Epihormomyia* Felt, 1915, *Eugeniomyia* Maia, Mendonça & Romanowski, 1997, *Jorgenseniella* Maia, 2005, *Liodiplosis* Gagné, 2001, *Manilkaramyia* Maia, 2001b, *Mayteniella* Maia, 2001b, *Myrciomyia* Maia, 1996c, *Myrciariamyia* Maia, 1995, *Neomitranthella* Maia, 1996c, *Parazalepidota* Maia, 2001b, *Paulliniamyia* Maia, 2001b, and *Perasphondylia* Möhn, 1960). Fourteen have been reported only from Brazil (*Arrabiadaeamyia*, *Clusiamyia*, *Cordiomyia*, *Costadiplosis*, *Dactylodiplosis*, *Eugeniomyia*, *Jorgenseniella*, *Liodiplosis*, *Mayteniella*, *Myrciomyia*, *Myrciariamyia*, *Neomitranthella*, *Manilkaramyia*, and *Parazalepidota*) (Gagné & Jaschhof, 2017), of which eight can be considered endemic to the country because they occur exclusively on Brazilian endemic host plant species (*Clusiamyia*, *Costadiplosis*, *Jorgenseniella*, *Mayteniella*, *Myrciomyia*, *Neomitranthella*, *Manilkaramyia*, and *Parazalepidota*). Six of these are endemic to the Atlantic Forest and, until now, have been exclusively reported for restingas (*Clusiamyia*, *Costadiplosis*, *Jorgenseniella*, *Neomitranthella*, *Manilkaramyia*, and *Parazalepidota*).

Table 1. Gall-inducing species of potential economic interest in Brazilian restingas (Atlantic Forest), including their host plants (scientific and common names), gall morphology (galled organ and gall shape), economic importance, natural enemies, geographic distribution, and references.

Gall-inducing species	Host plant scientific/ common name	Galled organ/ gall shape	Economical importance	Natural enemies	Geographic distribution	References
Diptera, Agromyzidae:						
<i>Japanagomyza inferna</i> Spencer, 1973	<i>Centrosema virginianum</i> (L.) Benth. (Fabaceae)/jetirana	Leaf/globoid	Agroforestry (erosion control)	No data	Bahamas, Guadeloupe, Brazil (RJ: Mangaratiba, Saquarema, Arraial do Cabo, São João da Barra)	Carvalho-Fernandes <i>et al.</i> (2016), Sousa & Couri (2014), Spencer & Stegmaier (1973), Useful Tropical Plants (2014)
Diptera, Cecidomyiidae:						
<i>Alycaulus globulus</i> Gagné, 2001	<i>Mikania glomerata</i> Spreng. (Asteraceae)/guaco	Leaf/lenticular	Edible (as tea), medicinal (treatment of asthma and bronchitis)	Parasitoids: Platygasteridae (Hymenoptera)	Brazil: RJ (Itatiaia, Rio de Janeiro, Silva Jardim)	Gagné <i>et al.</i> (2001), Maia (2013a), Maia & Mascarenhas (2017), Maia <i>et al.</i> (2008), Oliveira & Maia (2005), Skuhrová (1989), Teske & Trentini (1997)
<i>Arrabidaemyia serrata</i> Maia, 2001	<i>Fridericia conjugata</i> (Vell.) Mart. (Bignoniaceae)/Cipó bugi	Leaf/conical	Manufacture of rope	Parasitoids: <i>Eurytoma</i> sp. (Eurytomidae, Hymenoptera)	Brazil: RJ (Araruama, Arraial do Cabo, Carapebus, Mangaratiba, Maricá, Rio de Janeiro)	Carvalho-Fernandes <i>et al.</i> (2016), Maia (2001a, b, 2013a), Maia & Azevedo (2009), Monteiro <i>et al.</i> (1994, 2004), Oliveira & Maia (2005), Rodrigues <i>et al.</i> (2014), Santos <i>et al.</i> (2009)
<i>Asphondylia communis</i> Maia & Couri, 1992	<i>Ximeria americana</i> L. (Olacaceae)/ameixa-do-brasil	Stem/fusiform	Edible, medicinal (laxative, treatment for fevers, colds, toothaches, headaches, angina), agroforestry, manufacture of soap, body lubrication, and hair oil	Parasitoids: Encyrtidae (Hymenoptera)	RJ (Maricá)	Maia (2001a, 2013a), Maia & Azevedo (2009), Maia <i>et al.</i> (1992), Useful Tropical Plants (2014)
<i>Asphondylia cf. cordiae</i> Möhn, 1959	<i>Varronia curassavica</i> Jacq. (Boraginaceae)/erva-baleeira	Flower bud/ovoid	Medicinal (as anti-inflammatory) manufacture of rope, mosquito repellent	Parasitoids: <i>Eurytoma</i> sp. (Eurytomidae), Eulophidae, Elasmidae (Hymenoptera)	El Salvador Brazil: RJ (Araruama, Arraial do Cabo, Cabo Frio, Maricá, Saquarema, São Francisco de Itabapoana, São João da Barra), SP (Bertioga)	Carvalho-Fernandes <i>et al.</i> (2016), Maia (1996a, 2001a, 2013a), Maia & Carvalho-Fernandes (2016), Maia & Azevedo (2009), Maia <i>et al.</i> (2008), Möhn (1959), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Asphondylia mœhni</i> Skuhrová, 1989	<i>Mikania glomerata</i> Spreng. (Asteraceae)/guaco	Stem/globoid	Edible (as tea), medicinal (treatment of asthma and bronchitis)	Parasitoids: Hymenoptera	Brazil: RJ (Itatiaia, Poço das Antas, Silva Jardim, Rio de Janeiro)	Gagné <i>et al.</i> (2001), Maia (2013a), Maia & Mascarenhas (2017), Maia <i>et al.</i> (2008), Oliveira & Maia (2005), Skuhrová (1989), Teske & Trentini (1997)
<i>Bruggmanniella bysoniniae</i> (Maia & Couri, 1992)	<i>Byrsominia sericea</i> DC. (Malvaceae)/murici da praia	Flower bud/ovoid	Edible, ornamental, carpentry, source of tannins, textile dyes	Parasitoids: Encyrtidae (Hymenoptera)	Brazil: RJ (Mangaratiba, Maricá, São Francisco de Itabapoana, São João da Barra)	Carvalho-Fernandes <i>et al.</i> (2016), Maia (2001a, b, 2013a), Maia <i>et al.</i> (1992), Maia & Carvalho-Fernandes (2016), Maia & Azevedo (2009), Monteiro <i>et al.</i> (1994, 2004), Rodrigues <i>et al.</i> (2014), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Bruggmanniella maytenuse</i> (Maia & Couri, 1992)	<i>Maytenus obtusifolia</i> Mart. (Celastraceae)/papagaio, carne de anta, congonha brava de folha miúda, lenha branca	Fruit/ovoid	Edible (as tea), medicinal (treatment of diarrhea, serious ulcers, general inflammations, cancer and external ulcers on the skin), carpentry, rural construction work	Parasitoids: Eulophidae (Hymenoptera)	Brazil: RJ (Maricá, São Francisco de Itabapoana, São João da Barra)	Carvalho-Fernandes <i>et al.</i> (2016), Maia (1999b, 2001a, b, 2013a), Maia & Azevedo (2009), Maia & Carvalho-Fernandes <i>et al.</i> (2016), Maia <i>et al.</i> (1992), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)

Gall-inducing species	Host plant scientific/ common name	Galled organ/ gall shape	Economical importance	Natural enemies	Geographic distribution	References
<i>Climodiplosis conica</i> Oliveira & Maia, 2008	<i>Microstachys comiculata</i> (Vahl) Griseb. (Euphorbiaceae)/falsa-guanxuma, guanxuma-de-chiffe	Bud/conical	Medicinal (treatment of headaches, heart problems, sores, thrush, wounds and cuts)	Parasitoids: Eulophidae (Hymenoptera)	Brazil: RJ (Carapebus, Maricá)	Maia (2001a, 2013a), Maia & Azevedo (2009), Oliveira & Maia (2008). Useful Tropical Plants (2014)
<i>Climodiplosis costata</i> Maia, 2005	<i>Paullinia weinmanniifolia</i> Mart. – (Sapindaceae)/cipó-sangue	Leaf/leaf roll	Ornamental	Predator: <i>Lestodiplosis</i> sp. (Cecidomyiidae, Diptera)	Brazil: RJ (Arraial do Cabo, Carapebus, Maricá)	Maia (2001a, 2005a, 2013a), Monteiro <i>et al.</i> (1994, 2004), Santos <i>et al.</i> (2009)
<i>Climodiplosis profusa</i> Maia, 2001	<i>Eugenia uniflora</i> L. (Myrtaceae)/pitangueira	Leaf/conical	Edible, medicinal (head colds, influenza, chest colds, coughs and fever), carpentry, insect repellent	Parasitoids: <i>Chrysonotomomyia</i> sp., <i>Aprostocetus</i> sp. (Eulophidae, Hymenoptera)	Brazil: RJ (Araruama, Arraial do Cabo, Carapebus, Mangaratiba, Maricá, Rio de Janeiro, São João da Barra, Saquarema), RS (Pelotas)	Bierhals <i>et al.</i> (2012), Diez-Rodriguez <i>et al.</i> (2011), Maia (1993, 2013a), Maia & Azevedo (2009), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Clusiomyia granulosa</i> Maia, 2001	<i>Clusia hilariana</i> Schltdl. (Clusiaceae)/clísia-baiana	Leaf/lenticular	Edible, medicinal (treatment of diarrhea), carpentry, ornamental, fuel source or charcoal manufacture	Parasitoids: Eupelmidae (Hymenoptera)	Brazil: ES (Guaapari), RJ (Arraial do Cabo, Carapebus)	Bregonci <i>et al.</i> (2010), Maia (2001a, b, 2013a), Maia & Azevedo (2009), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Clusiomyia nitida</i> Maia, 1997	<i>Clusia lanceolata</i> Camb. (Clusiaceae)/cebola-da-praia	Leaf/globoid	Used as fuel	Parasitoids: <i>Ercasia</i> sp. (Aphelinidae), <i>Innosterna</i> p. (Platygastridae) (Hymenoptera)	Brazil: RJ (Arraial do Cabo, Maricá)	Maia (1997, 2001a, 2013a), Maia & Azevedo (2009), Useful Tropical Plants (2014)
<i>Contarinia gemmae</i> Maia, 2003	<i>Calophyllum brasiliense</i> Cambess. (Calophyllaceae)/guanandi	Bud/globoid	Carpentry, medicinal (antidiabetic and vermífuge, headache), agroforestry (to stabilize soils and to relieve soil compaction in degraded pastures), general construction	No data	Brazil: AM (Reserva de Desenvolvimento Sustentável Amanã), BA (Parque Estadual da Serra dos Montes Altos), GO (Prenópolis), MG (Januária, São Tomé das Letras), RJ (Carapebus), SP (Bertioga)	Arriola <i>et al.</i> (2016), Madeira <i>et al.</i> (2003), Maia (2013a), Maia <i>et al.</i> (2008), Proença & Maia (2015), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Conidiomyia globosa</i> Maia, 1996	<i>Varronia curassavica</i> Jacq. (Boraginaceae)/erva-baleira	Leaf/globoid	Medicinal (as anti-inflammatory)	Parasitoids: <i>Galeopsomyia</i> sp. and <i>Cirrospilus</i> sp. (Eulophidae), <i>Erytoma</i> sp. (Eurytomidae), <i>Lyrcus</i> sp. (Peromalidae), <i>Synopaeas</i> sp. (Platygastridae), <i>Torymnoidea</i> sp. and <i>Torymus</i> sp. (Torymidae) (Hymenoptera)	Brazil: ES (Guaapari), RJ (Arraial do Cabo, Cabo Frio, Carapebus, Maricá, São Francisco de Itabapoana, São João da Barra), Saquarema), SP (Bertioga)	Bregonci <i>et al.</i> (2010), Carvalho-Fernandes <i>et al.</i> (2016), Maia (1996a, 2001a, 2013a), Maia & Azevedo (2009), Maia & Tavares (2000), Maia <i>et al.</i> (2008), Möhn (1959), Santos <i>et al.</i> (2009) Useful Tropical Plants (2014)
<i>Costadiplosis maricaensis</i> Vreconite & Maia, 2009	<i>Psittacanthus dichroos</i> (Mart.) Mart. (Loranthaceae)/erva-de-passarinho	Leaf/lenticular	Medicinal (treatment of haemorrhoids)	Parasitoids: <i>Aprostocetus</i> sp. (Eulophidae, Hymenoptera)	Brazil: RJ (Maricá)	Maia (2001a, 2013a), Maia & Azevedo (2009), Santos <i>et al.</i> (2009), Vreconite & Maia (2009)
<i>Dactylopiosis heptaphylli</i> Maia, 2004	<i>Protium heptaphyllum</i> (Aubl.) March. (Burseraceae)/almecega	Leaf/globoid	Edible, medicinal (treatment of spine problems, asthma and bronchitis), carpentry, agroforestry (excellent shade used for incense and in varnishes)	No data	Brazil: RJ (Carapebus, São Francisco de Itabapoana, São João da Barra)	Carvalho-Fernandes <i>et al.</i> (2016), Maia (2013a), Maia & Carvalho-Fernandes (2016), Narahara <i>et al.</i> (2004), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Dactylopiosis icaribae</i> Maia, 2002	<i>Protium icariba</i> (DC.) March. (Burseraceae)/insenso	Leaf/conical	Edible, medicinal (treatment of spine and skin problems, antirheumatic, haemostatic, expectorant, and stimulant)	No data	Brazil: RJ (Carapebus, Mangaratiba)	Maia (2001a, 2002, 2013a), Maia <i>et al.</i> (2002), Rodrigues <i>et al.</i> (2014), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Dasineura byrsionimae</i> Maia, 2010	<i>Byrsionima sericea</i> DC. (Malpighiaceae)/murici da praia	Leaf/lenticular	Edible, ornamental, carpentry, source of tannins, textile dyes	Parasitoids: Hymenoptera	Brazil: ES (Guaapari), RJ (Angra dos Reis, Araruama, Arraial do Cabo, Cabo Frio, Carapebus, Mangaratiba, Maricá, Rio de Janeiro, São Francisco de Itabapoana, São João da Barra, Saquarema)	Bregonci <i>et al.</i> (2010), Carvalho-Fernandes <i>et al.</i> (2016), Maia (2001a, 2010, 2013a), Maia & Carvalho-Fernandes (2016), Maia & Oliveira (2010), Oliveira & Maia (2005), Monteiro <i>et al.</i> (2004), Rodrigues <i>et al.</i> (2014), Santos <i>et al.</i> (2009) Useful Tropical Plants (2014)

Gall-inducing species	Host plant scientific/ common name	Galled organ/ gall shape	Economical importance	Natural enemies	Geographic distribution	References
<i>Dasineura gigantea</i> Angelo & Maia, 1999	<i>Psidium cattleianum</i> Sabine (Myrtaceae)/Araçá	Bud/leaf rosette	Edible, flowers used as a garnish, medicinal (treatment for haemorrhages), agroforestry, ornamental, used for lathe work, tool handles and for objects that require much resistance, fuel and charcoal	Parasitoids: <i>Leptacis</i> sp. (Platygastridae, Hymenoptera), <i>Alysiini</i> sp. (Braconidae)	Brazil: PR (Piraquara, Pontal do Paraná), RS (Pelotas), SP (Bertioga), SC (Itapoa)	Angelo (2007), Angelo & Maia (1999), Diez-Rodríguez <i>et al.</i> (2011), Maia (2013a), Maia <i>et al.</i> (2008), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Dasineura globosa</i> Maia, 1996	<i>Eugenia astringens</i> Cambess. (Myrtaceae)/aperta-goela	Leaf/lenticular	Medicinal (treatment of diarrhea)	Parasitoids: Eulophidae (Hymenoptera)	Brazil: RJ (Araruama, Arraial do Cabo, Cabo Frio, Carapebus, Mangaratiba, Maricá, Rio de Janeiro, São João da Barra, Saquarema)	Carvalho-Fernandes <i>et al.</i> (2016), Maia (1996b, 2001a, 2013a), Maia <i>et al.</i> : 2005, Maia & Azevedo (2009), Oliveira & Maia (2005), Santos <i>et al.</i> (2009), Tavares (1920)
<i>Dasineura marginalis</i> Maia, 2005	<i>Eugenia astringens</i> Cambess. (Myrtaceae)/aperta-goela	Leaf/marginal roll	Medicinal (treatment of diarrhea)	Parasitoids: Eulophidae (Hymenoptera) Predator: <i>Lestodiplosis</i> sp. (Cecidomyiidae, Diptera)	Brazil: RJ (Araruama, Arraial do Cabo, Cabo Frio, Carapebus, Mangaratiba, Maricá, Rio de Janeiro, Saquarema)	Carvalho-Fernandes <i>et al.</i> (2016), Maia (1993, 2013a), Maia <i>et al.</i> : 2005, Oliveira & Maia (2005), Rodrigues <i>et al.</i> (2014), Santos <i>et al.</i> (2009), Tavares (1920)
<i>Dasineura myrcianae</i> Maia, 1996	<i>Myrciaria floribunda</i> (H. West ex Willd.) Legrand (Myrtaceae)/cambui-amarelo	Leaf/marginal roll	Edible fruit	Parasitoids: <i>Proacris</i> sp. (Eulophidae), Aphelinidae (Hymenoptera) Predator: <i>Lestodiplosis</i> sp. (Cecidomyiidae, Diptera)	Brazil: ES (Guarapari), RJ (Carapebus, Maricá, Rio de Janeiro, São Francisco de Itabapoana)	Bregonci <i>et al.</i> (2010), Maia (1996b, 1994, 2001a, 2013a), Maia & Azevedo (2009), Maia & Carvalho-Fernandes (2016)
<i>Dasineura ovalifoliae</i> Maia & Carvalho-Fernandes, 2011	<i>Erythroxylum ovalifolium</i> Peyr. (Erythroxylaceae)/fruta-de-pomba	Leaf/triangular	Medicinal (spine problems)	Parasitoids: Eulophidae, Eupelmidae, Mymaridae, Pteromalidae (Hymenoptera) Modifier: Tetrastichinae (Eulophidae, Hymenoptera)	Brazil: RJ (Araruama, Arraial do Cabo, Cabo Frio, Carapebus, Maricá, Rio de Janeiro, Saquarema)	Carvalho-Fernandes <i>et al.</i> (2016), Maia (2001a, 2013a), Maia & Azevedo (2009), Maia & Fernandes (2011), Santos <i>et al.</i> (2009)
<i>Dasineura tavares</i> Maia, 1996 (inquiline or gall maker)	<i>Neomitranthes obscura</i> (DC.) N.J.E. Silveira (Myrtaceae)/cambui-preto	Leaf/marginal roll	Edible fruit, medicinal (treatment of diarrhea), fuel and charcoal	Parasitoids: Encyrtidae (Hymenoptera) Free living phytophagous: <i>Stenomma amosa</i> (Butler, 1877) (Oecophoridae, Lepidoptera)	Brazil: ES (Guarapari), RJ (Araruama, Arraial do Cabo, Cabo Frio, Carapebus, Maricá, Saquarema)	Bregonci <i>et al.</i> (2010), Carvalho-Fernandes <i>et al.</i> (2016), Maia (1996b, 1995, 2001a, 2013a), Maia & Azevedo (2009), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Epithormomyia miconiae</i> Maia, 2001	<i>Miconia cinnamomifolia</i> (DC.) Naudin. (Melastomataceae)	Stem/ovoid	Carpentry, agroforestry (pioneer in reforestation projects), ornamental	Inquiline: <i>Resellella</i> sp. (Cecidomyiidae, Diptera)	Brazil: RJ (Carapebus)	Maia (2001a, b, 2013a), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Eugeniomyia dispar</i> Maia, Mendonça & Romanovski, 1996	<i>Eugenia uniflora</i> L. (Myrtaceae)/Pitangueira	Leaf/globoid	Edible, medicinal (head colds, influenza, chest colds, coughs and fever), carpentry, insect repellent	Parasitoids: <i>Rileyia hegeli</i> Girault, 1916 (Eurytomidae, Hymenoptera) Predator: <i>Pseudomyrmex</i> sp. (Formicidae, Hymenoptera) Phytophagous: <i>Pheidole</i> sp. (Formicidae, Hymenoptera)	Brazil: SP (Bertioga), SC (Florianópolis), RS (Pelotas, Porto Alegre, Juiú, Parque Florestal de Nonoai)	Bierhals <i>et al.</i> (2012), Diez-Rodríguez <i>et al.</i> (2011), Maia (1993, 2013a), Maia <i>et al.</i> (1996, 2008), Maia & Nava (2011), Mendonça & Romanovski, 2002a, b, Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Eugeniomyia triangularis</i> Maia & Nava, 2001	<i>Eugenia uniflora</i> L. (Myrtaceae)/Pitangueira	Leaf/triangular	Edible, medicinal (head colds, influenza, chest colds, coughs and fever), carpentry, insect repellent	No data	Brazil: RJ (Maricá)	Maia (1993, 2013a), Maia <i>et al.</i> (1996), Maia & Nava (2011), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Jorgensenella eugeniae</i> Maia, 2005	<i>Eugenia astringens</i> Cambess. (Myrtaceae)/aperta-goela	Leaf/lenticular	Medicinal (treatment of diarrhea)	No data	Brazil: RJ (Arraial do Cabo, Macaé)	Maia (1993, 2013a), Maia <i>et al.</i> : 2005, Santos <i>et al.</i> (2009), Tavares (1920)
<i>Liodiplosis conica</i> Gagné, 2001	<i>Mikania glomerata</i> Spreng. (Asteraceae)/guaco	Leaf/conical	Edible (as tea), medicinal (treatment of asthma and bronchitis)	No data	Brazil: RJ (Itaitiaia, Rio de Janeiro, Silva Jardim)	Gagné <i>et al.</i> (2001), Maia (2013a), Maia & Mascarenhas (2017), Oliveira & Maia (2005), Skuhraivá (1989), Teske & Trentini (1997)

Gall-inducing species	Host plant scientific/ common name	Galled organ/ gall shape	Economical importance	Natural enemies	Geographic distribution	References
<i>Lioposis cylindrica</i> Gagné, 2001	<i>Mikania glomerata</i> Spreng. (Asteraceae)/guaco	Leaf/cylindrical	Edible (as tea), medicinal (treatment of asthma and bronchitis)	Parasitoids: Hymenoptera	Brazil: RJ (Itatiaia, Paraty, Silva Jardim)	Gagné <i>et al.</i> (2001), Maia (2013a), Maia & Mascarenhas (2017), Skuhrová (1989), Teske & Trentini (1997)
<i>Lioposis spherica</i> Gagné, 2001	<i>Mikania glomerata</i> Spreng. (Asteraceae)/guaco	Leaf/spherical	Edible (as tea), medicinal (treatment of asthma and bronchitis)	Parasitoids: Hymenoptera	Brazil: RJ (Itatiaia, Paraty, Silva Jardim)	Gagné <i>et al.</i> (2001), Maia (2013a), Maia & Mascarenhas (2017), Skuhrová (1989), Teske & Trentini (1997)
<i>Lopesia caulinaris</i> Maia, 2003	<i>Calophyllum brasiliense</i> Cambess. (Calophyllaceae)/guanandi	Stem/globoid	Carpentry, medicinal (antidiabetic and vermifuge, headache), agroforestry (to stabilize soils and to relieve soil compaction in degraded pastures), general construction	Parasitoids: Hymenoptera	Costa Rica (Heredia), Bolívia (Santa Cruz), Brazil: AM (Reserva de Desenvolvimento Sustentável Amanã), Amapá (Oiapoque), BA (Parque Estadual da Serra dos Montes Altos), GO (Pirenópolis), MG (São Januária, São Tomé das Letras), RJ (Carapebus), SP (Bertioga), SC (São Francisco do Sul)	Arriola <i>et al.</i> (2016), Madeira <i>et al.</i> (2003), Maia (2013a, 2013b), Maia <i>et al.</i> (2008), Proença & Maia (2015), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Lopesia conspicua</i> Maia, 2003	<i>Calophyllum brasiliense</i> Cambess. (Calophyllaceae)/guanandi	Leaf/globoid	Carpentry, medicinal (antidiabetic and vermifuge, headache), agroforestry (to stabilize soils and to relieve soil compaction in degraded pastures), general construction	No data	Brazil: AM (Reserva de Desenvolvimento Sustentável Amanã), RN (Canguaretama), BA (Parque Estadual da Serra dos Montes Altos), GO (Pirenópolis), MG (Januária, São Tomé das Letras), RJ (Carapebus), PR (Paranaíba)	Arriola <i>et al.</i> (2016), Madeira <i>et al.</i> (2003), Maia (2013a, 2013b), Proença & Maia (2015), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Lopesia elliptica</i> Maia, 2003	<i>Calophyllum brasiliense</i> Cambess. (Calophyllaceae)/guanandi	Leaf/elliptical	Carpentry, medicinal (antidiabetic and vermifuge, headache remedy), agroforestry (to stabilize soils and to relieve soil compaction in degraded pastures), general construction	Parasitoids: Eurytomidae (Hymenoptera) Inquilines: Coleoptera	México (Coahuila), Guatemala (Santa Helena and San Francisco), Costa Rica (Heredia), Dominican Republic (Dajabón), Cuba, Guyana (Upper Takutu-Upper Essequibo), Bolívia (Santa Cruz), Peru (Loreto), Paraguay (Amambay), Brazil: AP (Oiapoque), AM (Reserva de Desenvolvimento Sustentável Amanã), PA (Mojú), MA (São Luís), RO (Chupinguaia), RN (Canguaretama), BA (Parque Estadual da Serra dos Montes Altos), Pernambuco, GO (Pirenópolis), TO (Formoso do Araguaia), MT (Santa Terezinha), MG (São Tomé das Letras), RJ (Carapebus), SP (Bertioga), PR (Paranaíba), SC (São Francisco do Sul)	Arriola <i>et al.</i> (2016), Madeira <i>et al.</i> (2003), Maia (2013a), Maia <i>et al.</i> (2008), Proença & Maia (2015), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Lopesia erythroxylifera</i> Rodrigues & Maia, 2010	<i>Erythroxylum ovalifolium</i> Peyr. (Erythroxylaceae)/fruta-de-pomba	Bud/conical	Medicinal (spine problems)	Parasitoids: Eulophidae, Eupelmidae, <i>Eurytoma</i> sp. (Eurytomidae), Mymaridae, Pteromalidae (Hymenoptera) Modifier: Tetrastichinae (Eulophidae, Hymenoptera)	Brazil: RJ (Angra dos Reis, Araruama, Arraial do Cabo, Cabo Frio, Carapebus, Maricá, Rio de Janeiro, São João da Barra, Saquarema)	Carvalho-Fernandes <i>et al.</i> (2016), Maia (2013a), Maia & Azevedo (2009), Maia & Carvalho-Fernandes (2011), Maia & Oliveira (2010), Rodrigues & Maia (2010), Santos <i>et al.</i> (2009)
<i>Lopesia linearis</i> Maia, 2003	<i>Calophyllum brasiliense</i> Cambess. (Calophyllaceae)/guanandi	Leaf/lineal	Carpentry, medicinal (antidiabetic and vermifuge, headache remedy), agroforestry (to stabilize soils and to relieve soil compaction in degraded pastures), general construction	No data	Costa Rica (Heredia), Cuba, Trinidad and Tobago (Saint George), México (Coahuila), Bolívia (Santa Cruz), Colômbia (Antioquia), Guyana (Upper Takutu-Upper Essequibo), Peru (Loreto), Paraguay (Amambay), Brazil: AM (Reserva de Desenvolvimento Sustentável Amanã), PA (Mojú), RO (Chupinguaia), RR (Caracará), RN (Canguaretama), MT (Santa Terezinha), MS (Corumbá), MG (São Tomé das Letras), ES (Linhares), RJ (Carapebus), SP (Bertioga), PR (Paranaíba), SC (São Francisco do Sul)	Arriola <i>et al.</i> (2016), Madeira <i>et al.</i> (2003), Maia (2013a and b), Maia <i>et al.</i> (2008), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Lopesia maricaensis</i> Rodrigues & Maia, 2009	<i>Protium brasiliense</i> (Spr.) Engl. (Engl. Calophyllaceae)	Leaf/marginal roll	Edible (oil obtained from seeds), carpentry	Parasitoids: <i>Goniozus</i> sp. (Bethyloidea, Hymenoptera)	Brazil: RJ (Maricá)	Maia (2013a), Maia & Azevedo (2009), Rodrigues & Maia (2010), Useful Tropical Plants (2014)

Gall-inducing species	Host plant scientific/ common name	Galled organ/ gall shape	Economical importance	Natural enemies	Geographic distribution	References
<i>Lopesia similis</i> Maia, 2004	<i>Protium heptaphyllum</i> (Aubl.) March. (Burseraceae)/almecega	Leaf/marginal roll	Edible, medicinal (treatment of spine problems, asthma and bronchitis), carpentry, agroforestry (excellent shade used for incense and in varnishes)	Parasitoids: Eupelmidae (Hymenoptera)	Brazil: MG (Tridentes), MS (Aquidauana), RJ (Arraial do Cabo, Carapebus, São João da Barra)	Carvalho-Fernandes <i>et al.</i> (2016), Maia (2013a), Maia & Azevedo (2009), Maia & Fernandes (2004), Narahara <i>et al.</i> (2004), Santos <i>et al.</i> (2009), Silva & Maia (2014), Useful Tropical Plants (2014)
<i>Lopesia simplex</i> Maia, 2002	<i>Protium icariba</i> DC. March. (Burseraceae)/insenço	Leaf/marginal roll	Edible, medicinal (treatment of spine and skin problems, antirheumatic, haemostatic, expectorant, and stimulant)	Parasitoids: Hymenoptera	Brazil: ES (Guarapari), RJ (Araruama, Carapebus, Mangaratiba, São João da Barra)	Bregonci <i>et al.</i> (2010), Carvalho-Fernandes <i>et al.</i> (2016), Maia (2001a, 2002, 2013a), Maia <i>et al.</i> (2002), Rodrigues <i>et al.</i> (2014), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Lopesia singularis</i> Maia, 2001	<i>Pouteria venosa</i> (Mart.) (Sapotaceae)/Sapota-preta	Leaf/lenticular	Edible fruit, medicinal (treatment for diarrhea and heartburn)	Parasitoids: Tetrastichinae (Eulophidae), Eupelmidae, Torymidae (Hymenoptera)	Brazil: RJ (Maricá, Carapebus), SP (Bertioga)	Maia (2001a, b, 2013a), Maia & Azevedo (2009), Maia <i>et al.</i> (2008), Monteiro <i>et al.</i> (1994), Useful Tropical Plants (2014)
<i>Mamilikaramyia notabilis</i> Maia, 2001	<i>Mamilikara subsericea</i> (Mart.) Dubard. (Sapotaceae)/maçaranduba-da-praia	Bud/ovoid	Edible fruit, carpentry	Parasitoids: Torymidae (Hymenoptera)	Brazil: ES (Guarapari), RJ (Carapebus)	Bregonci <i>et al.</i> (2010), Maia (2001a, b, 2013a), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Mayeniella distincta</i> Maia, 2001	<i>Maytenus obtusifolia</i> Mart. (Celastraceae)/papagalo, carne de anta, congonha brava de folha miúda, lenha branca	Leaf/globoid	Edible (as tea), medicinal (treatment of diarrhea, serious ulcers, general inflammations, cancer and also against external ulcers on the skin), carpentry, rural construction work	Parasitoids: <i>Xanthellum</i> (Eulophidae, Hymenoptera)	Brazil: RJ (Arraial do Cabo, Mangaratiba, Maricá, Rio de Janeiro, São Francisco de Itabapoana, São João da Barra)	Maia & Carvalho-Fernandes (2016), Maia (2001a, b, 2013a), Maia & Azevedo (2009), Maia & Silva (2016), Monteiro <i>et al.</i> (1994), Rodrigues <i>et al.</i> (2014), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Myrciameya maricaensis</i> Maia, 1996	<i>Myrcia amazonica</i> DC. and <i>M. ovata</i> Cambes. (Myrtaceae)/aracá-açaiara	Bud/ovoid	Edible fruit	Parasitoids: <i>Aprostocetus</i> sp. (Eulophidae), Platygastridae, Modifiers: Eulophidae (Hymenoptera)	Brazil: RJ (Arraial do Cabo, Carapebus, Maricá)	Ferraz & Monteiro (2003), Maia (1996c, 2001a, 2013a), Maia & Azevedo (2009), Monteiro <i>et al.</i> (1994), Santos <i>et al.</i> (2009)
<i>Myrciameya bivalva</i> Maia, 1994	<i>Myrciaria floribunda</i> (H. West ex Willd.) Legrand (Myrtaceae)/cambuí-amarelo	Bud/bivalve	Edible fruit	Parasitoids: Eulophidae, Encyrtidae, Pteromalidae (Hymenoptera)	Brazil: RJ (Arraial do Cabo, Carapebus, Maricá)	Maia (1993, 1994, 2001a, 2013a), Maia & Azevedo (2009), Monteiro <i>et al.</i> (1994, 2004)
<i>Neolasioptera acer</i> Rübsaamen, 1905	<i>Hyloceus setaceus</i> (Salm-Dyck) R. Bauer (Cactaceae)/cardeiro-trepador	Stem/fusiform	Edible fruit	No data	Brazil: RJ (Angra dos Reis, Cabo Frio, Carapebus, São João da Barra)	Carvalho-Fernandes <i>et al.</i> (2016), Maia (2013a), Maia & Oliveira (2010), Rübsaamen (1905), Useful Tropical Plants (2014)
<i>Neolasioptera eugeniae</i> Maia, 1993	<i>Eugenia uniflora</i> L. (Myrtaceae)/pitangueira	Leaf/lenticular	Edible, medicinal (head colds, influenza, chest colds, coughs and fever), carpentry, insect repellent	Parasitoids: Eulophidae, Eurytomidae (Hymenoptera)	Brazil: RJ (Angra dos Reis, Arraial do Cabo, Cabo Frio, Maricá, Rio de Janeiro)	Maia (1993, 2001a, 2013a), Maia & Azevedo (2009), Maia & Oliveira (2010), Monteiro <i>et al.</i> (1994), Oliveira & Maia (2005), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Neomitranthella robusta</i> Maia, 1996	<i>Neomitranthes obscura</i> (DC.) N.J.E. Silveira (Myrtaceae)/cambuí-preto	Bud/imbricated	Edible fruit, medicinal (treatment of diarrhea), fuel and charcoal	Inquiline: <i>Stenomamma ammosa</i> (Butler, 1877) (Decaphoridae, Lepidoptera)	Brazil: RJ (Araruama, Cabo Frio, Carapebus, Maricá, Saquarema)	Carvalho-Fernandes <i>et al.</i> (2016), Maia (1996c, 2001a) Maia <i>et al.</i> (2013a), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Parazalepidota clusiae</i> Maia, 2001	<i>Clusia fluminensis</i> Planch. & Triana (Clusiaceae)/abaneiro	Leaf/lenticular	Veterinary, medicinal, ornamental, agroforestry (reforestation near the coast) Fuel or charcoal	Parasitoids: <i>Rileya</i> sp. (Eurytomidae, Hymenoptera)	Brazil: RJ (Maricá, Rio de Janeiro)	Maia (2001a, b, 2013a), Maia & Azevedo (2009), Oliveira & Maia (2005), Useful Tropical Plants (2014)

Gall-inducing species	Host plant scientific/ common name	Galled organ/ gall shape	Economical importance	Natural enemies	Geographic distribution	References
<i>Paulliniomyia ampla</i> Maia, 2001	<i>Paullinia weinmannifolia</i> Mart. (Sapindaceae)/cipó-sangue	Leaf/conical	Ornamental	Parasitoids: Braconidae, Eulophidae, Eurytomidae, <i>Dimeromicrus cecidomyiae</i> Ashmead, 1887 (Torymidae), Platygastriidae (Hymenoptera) Gall modifier: Eulophidae (Hymenoptera)	Brazil: RJ (Araruama, Arraial do Cabo, Cabo Frio, Carapebus, Maricá, São Francisco de Itabapoana, São João da Barra, Saquarema), SP (Bertioga)	Carvalho-Fernandes <i>et al.</i> (2016), Maia (2001a, b, 2013a), Maia & Azevedo (2009), Maia & Carvalho-Fernandes (2016), Maia <i>et al.</i> (2008), Santos <i>et al.</i> (2009)
<i>Perasphondylia mikaniae</i> Gagné, 2001	<i>Mikania glomerata</i> Spreng. (Asteraceae)/guaco	Bud/ovoid	Edible (as tea), medicinal (treatment of asthma and bronchitis)	Parasitoids: Hymenoptera	Brazil: RJ (Itatiaia, Silva Jardim), SP (Bertioga), RS	Gagné <i>et al.</i> (2001), Maia (2013a), Maia <i>et al.</i> (2008), Maia & Mascarenhas (2017), Mendonça <i>et al.</i> (2014), Skuhrová (1989), Teske & Trentini (1997), Useful Tropical Plants (2014)
<i>Schizomyia spherica</i> Maia & Oliveira, 2007	<i>Microstachys comiculata</i> (Vahl) Griseb. (Euphorbiaceae)/falsa-guanxuma, guanxuma-de-chifre	Bud/spherical	Medicinal (treatment of headaches, heart problems, sores, thrush, wounds and cuts)	Parasitoids: Eulophidae (Hymenoptera)	Brazil: RJ (Araruama, Arraial do Cabo, Carapebus, Maricá, Saquarema)	Maia (2013a), Maia & Azevedo (2009), Maia & Oliveira (2007), Useful Tropical Plants (2014)
<i>Stephomyia clavata</i> (Tavares, 1920)	<i>Eugenia astringens</i> Cambess. (Myrtaceae)/aperta-goela	Leaf/claviform	Medicinal (treatment of diarrhea)	Parasitoids: Eulophidae (Hymenoptera)	Brazil: RJ (Carapebus, Rio de Janeiro), BA (Madre de Deus)	Maia (1994, 2001a, 2013a), Maia <i>et al.</i> (2005), Maia & Azevedo (2009), Maia & Silva (2016), Santos <i>et al.</i> (2009), Tavares (1920)
<i>Stephomyia mina</i> Maia, 1994	<i>Neomitranthes obscura</i> (DC.) N.J.E. Silveira (Myrtaceae)/cambul-preto	Leaf/elliptical	Edible fruit, medicinal (treatment of diarrhea), fuel and charcoal	Parasitoids: Braconidae (Hymenoptera)	Brazil: RJ (Araruama, Arraial do Cabo, Carapebus, Maricá)	Carvalho-Fernandes <i>et al.</i> (2016), Maia (1993, 1995, 2001a, 2013a), Maia & Azevedo (2009), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014)
<i>Stephomyia rotundifoliarum</i> Maia, 1994	<i>Eugenia astringens</i> Cambess. (Myrtaceae)/aperta-goela	Bud/cylindrical	Medicinal (treatment of diarrhea)	Parasitoids: <i>Dimeromicrus cecidomyiae</i> (Torymidae), <i>Donackia</i> sp. (Braconidae), <i>Eurytoma</i> sp. and <i>Rileya</i> sp. (Eurytomidae), Eulophidae, Eupelmidae, Platygastriidae, Scelionidae, Torymidae (Hymenoptera) Modifiers: Hymenoptera Predators: <i>Novahorus</i> sp. (Olpitidae, Pseudoscorpiones)	Brazil: RJ (Araruama, Arraial do Cabo, Cabo Frio, Carapebus, Maricá, São João da Barra, Saquarema)	Carvalho-Fernandes <i>et al.</i> (2016), Maia (1994, 2001a, 2013a), Maia <i>et al.</i> (2005), Maia & Azevedo (2009), Monteiro <i>et al.</i> (1994), Santos <i>et al.</i> (2009), Tavares (1920)
<i>Youngomyia pouteriae</i> Maia, 2001	<i>Pouteria caimito</i> (Ruiz & Pav.) Radlk. (Sapotaceae)/abil	Leaf/cylindrical	Edible, carpentry (construction and external work)	Parasitoids: <i>Dimeromicrus cecidomyiae</i> (Torymidae), <i>Xanthellum</i> (Eulophidae), Eupelmidae, Platygastriidae, (Hymenoptera) Inquiline: <i>Troteria quadridentata</i> Maia, 2001c (Cecidomyiidae, Diptera)	Brazil: RJ (Araruama, Arraial do Cabo, Cabo Frio, Maricá, Rio de Janeiro, São João da Barra, Saquarema)	Maia (2001a, c, 2013a), Maia & Azevedo (2009), Monteiro <i>et al.</i> (1994), Oliveira & Maia (2005), Useful Tropical Plants (2014)
Hemiptera, Ericococcidae:						
<i>Nothotria tavaresi</i> (Crawford, 1925) (Psyllidae)	<i>Psidium cattleianum</i>	Leaf/lenticular	Edible, flowers used as a garnish, medicinal (treatment for haemorrhages), agroforestry, ornamental, used for lathe work, tool handles and for objects that require much resistance, fuel and charcoal	No data	Brazil: BA, RS	Burckhardt & Queiroz (2012), Carneiro <i>et al.</i> (2012), Crawford (1925), Mendonça <i>et al.</i> (2014), Tavares (1921, 1922)
<i>Tectococcus ovaris</i> Hempel, 1990	<i>Psidium cattleianum</i> Sabine (Myrtaceae)/araçá	leaf/biconvex	Edible, flowers used as a garnish, medicinal (treatment for haemorrhages), agroforestry, ornamental, used for lathe work, tool handles and for objects that require much resistance, fuel and charcoal	Parasitoids: <i>Metaphycus flavus</i> Howard, 1881 (Encyrtidae), <i>Aprostocetus</i> sp. (Eulophidae) (Hymenoptera) Predator: <i>Hyperaspis delicata</i> Massuti & Vittorino, 1997 (Coccinellidae, Coleoptera)	Brazil: RJ (Angra dos Reis), PR (Curitiba), RS	Angelo & Maia (1999), Hempel (1990), Maia (2013a), Maia & Oliveira (2010), Santos <i>et al.</i> (2009), Useful Tropical Plants (2014), Vittorino <i>et al.</i> (2007)

About 78% of the galling species compiled have been recorded only in Southeast Brazil, and about 64% only in the state of Rio de Janeiro, although their host plant species have broader geographic distributions. These inconsistencies between galler and host distributions are likely the result of a lack of data. For example, until two years ago *Lopesia caulinaris* Maia, 2003, *L. cylindrica* Maia, 2003 and *L. linearis* Maia, 2003 were known only from Southeast Brazil, but as the result of the examination of galled exsiccates deposited in several herbaria, the extent of their known distribution was greatly expanded, even to include other countries of South and Central America (Arriola *et al.*, 2016).

Biological data for the economically important gallers cited in the present study are scarce. To date, the population dynamics of only nine species [*Clusiomyia nitida* Maia, 1997, *Cordiamyia globosa* Maia, 1996a, *Dasineura gigantea* Angelo & Maia, 1999, *Dasineura globosa* Maia, 1993b, *Eugeniomyia dispar* Maia, Mendonça & Romanovski, 1996, *Myrciomyia maricaensis* Maia, 1996c, *Stephomyia rotundifoliorum* Maia, 1994, *Nothotrioza tavaresi* (Crawford, 1925), and *Tectococcus ovatus* Hempel, 1990] have been studied (Maia, 1999a; Maia & Tavares, 2000; Angelo, 2007, 2008; Mendonça & Romanowski, 2002a; Ferraz & Monteiro, 2003; Butignol & Pedrosa-Macedo, 2003; Vitorino *et al.*, 2000).

Data on natural enemies are also deficient, since most records have been published at the taxonomic levels of family and genus. Natural enemies include mainly parasitoids, which are associated with 43 galling species (about 74%) and are represented exclusively by Hymenoptera of 13 families: Aphelinidae, Bethyridae, Braconidae, Elasmidae, Encyrtidae, Eulophidae, Eupelmidae, Eurytomidae, Mymaridae, Platygasteridae, Pteromalidae, Scelionidae, and Torymidae. Most records of natural enemies are at the taxonomic levels of family ($n = 43$; about 51%) and genus ($n = 26$; about 32%), with other categories totaling about 16% (order $n = 7$, tribe $n = 1$, subtribe $n = 1$ and species $n = 5$).

At least 16 cecidogenous species are attacked by more than one species of parasitoid. Among these, *Stephomyia rotundifoliorum*, *Cordiamyia globosa*, *Lopesia erythroxylis* Rodrigues & Maia, 2010, *Paulliniomyia ampla* Maia, 2001b, *Dasineura ovalifoliae* Maia & Carvalho-Fernandes, 2011 and *Youngomyia pouteria* e Maia, 2001c, host the greatest richness of parasitoids (9, 7, 5, 5, 4, and 4 species, respectively). However, the role of each of these species in controlling the population of their host remains unknown, as does their level of specificity, which are essential data for evaluating their potential effectiveness as biological control agents.

Predators, inquiline and even free-living larvae that feed on galls have also been reported. The first, represented by cecidomyiids, pseudoscorpions, beetles, and ants, have been associated with five galling species and include records mainly at the genus level ($n = 5$), but also at the family ($n = 1$) and species ($n = 1$) levels. Although the last two guilds are phytophagous, they can cause death to the galler by interfering with gall growth, changing its morphology (in this case, they are

referred to as gall modifiers), destroying the gall or competing with the inducer for internal space and food. The inquiline are represented by cecidomyiids and beetles. The former have been associated with three gall-inducing species and their records are at the genus ($n = 2$) and species ($n = 1$) levels, while the latter were found in a single gall. Finally, the free living species, recorded only in galls of *Neomitranthes obscura* (DC.) N.J.E. Silveira, are represented by a single species of moth, *Stenoma annosa* (Butler, 1877) (Oecophoridae, Lepidoptera). This moth is a facultative inquiline, which feeds on leaves of species of Myrtaceae in restingas as well as on galled tissues. Since the caterpillar is highly voracious, it can destroy numerous galls, which always results in the galler's death (*personal observation*).

The impacts of all of these guilds on galler populations is poorly known since few studies on their biology have been published. In fact, data on parasitoidism and predation rates are known only for the nine cecidogenous species: *Clusiomyia nitida*, *Cordiamyia globosa*, *Dasineura gigantea*, *Dasineura globosa*, *Eugeniomyia dispar*, *Myrciomyia maricaensis*, *Stephomyia rotundifoliorum*, *Nothotrioza tavaresi*, and *Tectococcus ovatus* (Maia, 1999a; Maia & Tavares, 2000; Mendonça & Romanowski, 2002b; Bierhals *et al.*, 2012; Ferraz & Monteiro, 2003; Butignol & Pedrosa-Macedo, 2003; Vitorino *et al.*, 2000).

Most of the gallers found in the present study are associated with edible and/or medicinal plant species (about 59% and 68%, respectively), but there are also records of associations with plants used in agroforestry and carpentry, or as ornamentals and repellents, among others (Table 2). Some host plants have several uses, which explains the association percentages summing to more than 100%.

Although 58 gall-inducing species were identified in the present study, the number of insect galls associated with plants of economic interests in restinga environ-

Table 2. Number species host plants of economic importance and their gall-inducing insect species of Brazilian restingas (Atlantic Forest).

Host plant	Number of gall-inducing species	
	Use	Number species
Carpentry	24	19
Edible	34	34
Medicinal	33	39
Agroforestry	14	13
Fuel and charcoal	13	8
Ornamental/garnish	12	11
Cabinet making	9	0
Used to make ropes/whips	4	3
Ceramic	3	—
Pesticidal/repellent	3	4
Hair/body oil	2	1
Source of tannins	2	2
Aromatic baths	1	—
Soap manufacture and lubrication	1	1
Veterinary medicine	1	1
Weaving baskets	1	—
Textile dyes	1	2

ments is about three times greater, since a total of 186 gall morphotypes have been reported. Thus, 128 morphotypes were not included since the gall-inducing species remain undetermined. These data reveal the present deficiency of the taxonomical knowledge regarding gall-inducing taxa. Reasons for this scenario include: 1) a yet poorly understood Brazilian fauna; 2) few taxonomists focused on cecidogenous insects; 3) rearing gallers is difficult due to high mortality rates and little biological data, so there is often insufficient material for study; and 4) several studies have focused on ecological patterns, and do not consider galler identification.

This is the first list of cecidogenous insects of potential economic importance in Brazil, or in the Neotropical Region for that matter. Other areas of the world are better represented by such efforts, such as those of Barnes, 1946a, b, 1948, 1949, 1951, for the Nearctic Region, and Nijveldt, 1969; Harris, 1966; Jiang, 1994; and Skuhřavá, 1997 in the Palearctic Region. However, it must be remembered that as far as the Neotropical Region and Brazil are concerned, the present list is partial, as it focuses only on restinga environments. Nonetheless, this work represents a starting point for future compilations.

CONCLUSIONS

Restinga environments harbor a great number of gall-inducing species of potential economic importance. Knowledge of their geographic distributions, biology and impacts on host plants remains deficient, and their role as pests needs to be evaluated. There remains a large number of undetermined cecidogenous species, so an effort to identify them, perhaps through rearing, is essential. Since this list is restricted to restingas, studies of other phytophysiognomies will certainly amplify knowledge regarding the gall-inducing insects of potential economic importance in Brazil.

REFERENCES

- Angelo, A.C. 2007. Biologia de *Dasineura gigantea* Angelo & Maia, 1999 (Diptera, Cecidomyiidae), indutor de galhas sobre *P. cattleianum* e seus efeitos sobre a planta. In: Pedrosa-Macedo, J.H.; Dal Molin, A. & Smith, C.W. (Org.). O araçazeiro: ecologia e controle biológico. Curitiba, UFPEF do Paraná. p. 131-143.
- Angelo, A.C. 2008. Ciclo de vida de *Dasineura gigantea* Angelo & Maia, 1999 (Diptera, Cecidomyiidae). *Floresta*, 38: 23-32.
- Angelo, A.C. & Maia, V.C. 1999. *Dasineura gigantea* sp. n. (Diptera, Cecidomyiidae) associada a *Psidium cattleianum* Sabine (Myrtaceae) no Brasil. *Revista Brasileira de Zoologia*, 16: 191-195.
- Araújo, D.S.D. & Lacerda, L.D. 1987. A Natureza das Restingas. *Ciência Hoje*, 6: 42-48.
- Arriola, I.A.; Melo-Júnior, J.C.F. de; Mougá, D.M.D. da S.; Isaias, R.M. dos S. & Costa, E.C. 2016. Where host plant goes, galls go too: new records of the Neotropical galling Cecidomyiidae (Diptera) associated with *Calophyllum brasiliense* Cambess. (Calophyllaceae). *Check List*, 12(4): 1-8.
- Barnes, H.F. 1946a. *Gall Midges of Economic Importance*. I: Gall midges of root and vegetable crops. London, Crosby Lockwood & Son, Ltd. v. 1, 104p., 10 pls.
- Barnes, H.F. 1946b. *Gall Midges of Economic Importance*. II. Gall midges of fodder crops. London, Crosby Lockwood & Son, Ltd. v. 2, 160p., 3 pls.
- Barnes, H.F. 1948. *Gall midges of economic importance*. III. Gall midges of fruits. London, Crosby Lockwood & Son Ltd. v. 3, 184p.
- Barnes, H.F. 1949. *Gall Midges of Economic Importance*. VI. Miscellaneous Crops. London, Crosby Lockwood & Son, Ltd. v. 6, 229p., 14 pls.
- Barnes, H.F. 1951. *Gall Midges of Economic Importance*. V. Gall midges of trees. London, Crosby Lockwood & Son, Ltd. v. 5, 270p., 8 pls.
- Bierhals, A.N.; Nava, D.E.; Costa, V.A.; Maia, V.C. & Diez-Rodríguez, G.I. 2012. *Eugeniomyia dispar* em pitangueira: parasitoides associados, dinâmica populacional e distribuição de galhas na planta. *Revista Brasileira de Fruticultura*, 34: 109-115.
- Bregonci, J.M., Polycarpo, P.V. & Maia, V.C. 2010. Galhas de insetos do Parque Estadual Paulo César Vinha (Guarapari, ES, Brasil). *Biota Neotropica*, 10: 265-274.
- Burckhardt, D. & Queiroz, D.L. 2012. Checklist and comments on the jumping plant-lice (Hemiptera: Psylloidea) from Brazil. *Zootaxa*, 3571: 26-48.
- Butignol, C.A. & Pedrosa-Macedo, J.H. 2003. Biologia de *Neotrioza tavaresi* Crawford, 1925 (Hemiptera, Psyllidae), galhador da folha do araçazeiro (*Psidium cattleianum*). *Revista Brasileira de Entomologia*, 47: 1-7.
- Butler, A.G. 1877. Descriptions of new species of *Cryptolechia* from the Amazonas. *Cistula Entomologica*, 2: 187-193.
- Carneiro, M.A.A.; Branco, C.S.A.; Braga, C.E.D.; Almada, E.D.; Costa, M.B.M. & Maia, V.C. 2009. Are gall midge species (Diptera, Cecidomyiidae) host plant specialists? *Revista Brasileira de Entomologia*, 53: 365-378.
- Carneiro, R.G.S.; Burckhardt, D. & Isaias, R.M. 2012. Biology and Systematics of gall-inducing trizoids (Hemiptera: Psylloidea) associated with *Psidium* sp. (Myrtaceae). *Zootaxa*, 3620: 129-146.
- Carvalho-Fernandes, S.P.; Ascendino, S.; Maia, V.C. & Couri, M.S. 2016. Diversity of insect galls associated with coastal shrub vegetation in Rio de Janeiro, Brazil. *Anais da Academia Brasileira de Ciências*, 88: 1407-1418.
- Cotes, E.C. Entomology notes. *Indian Museum Notes*, 1: 82-124, pls. V-VII.
- Crawford, D.L. 1925. Psyllidae of South America. *Brotéria, Série Zoológica*, 22: 56-74.
- Diez-Rodríguez, G.I.; Nava, D.E.; Antunes, L.E.C.; Maia, V.C. & Hübner, L.K. 2011. *Insetos Galhadores da Pitangueira e do Araçazeiro no Sul do Rio Grande do Sul*. Embrapa Clima Temperado, Pelotas, RS, 29p. (Documento 337).
- Empresa Brasileira de Assistência Técnica e Extensão Rural, Centro Internacional de Agricultura Tropical (EMBRATER/CIAT). 1982. *Descrição das pragas que atacam a mandioca (Manihot esculenta Crantz) e características de seus prejuízos*. Brasília, DF, EMBRATER/DIRET. 47p.
- Felt, E.P. 1908. Appendix D. In: his 23d report of the State Entomologist on injurious and other insects of the State of New York, 1907. *New York State Museum Bulletin*, 124: 286-422; 489-510.
- Felt, E.P. 1915. New South American gall midges. *Psyche*, 22: 152-157.
- Felt, E.P. 1922. A new and remarkable fig midge. *The Florida Entomologist*, 6: 5-6.
- Fernandes, G.W. & Price, P.W. 1992. The adaptative significance of insect gall distribution: survivorship of species in xeric and mesic habitats. *Oecologia*, 9: 14-20.
- Fernandes, G.W. & Ribeiro, S.P. 1990. Plant response to herbivory: two examples from the neotropics. *Ecotropicos*, 3: 77-86.
- Ferraz, F.F.F. & Monteiro, R.F. 2003. Complex interactions involving a gall midge *Myrciomyia maricaensis* Maia (Diptera, Cecidomyiidae), phytophagous modifiers and parasitoids *Revista Brasileira de Zoologia*, 20: 433-437.
- Flora do Brasil. 2020. *Jardim Botânico do Rio de Janeiro*. [under construction]. Available at: <http://floradobrasil.jbrj.gov.br>. Access in: 24/11/2017.

- Florentine, S.K.; Raman, A. & Dhileepan, K. 2005. Effects of gall induction by *Epiblema strenuana* on gas exchange, nutrients, and energetics in *Parthenium hysterophorus*. *Biocontrol*, 50: 787-801.
- Gagné, R.J. 1990. Gall midge complex (Diptera: Cecidomyiidae) in bud galls of Palearctic *Euphorbia* (Euphorbiaceae). *Annals of the Entomological Society of America*, 83: 335-345.
- Gagné, R.J. & Jaschhof, M. 2017. *A Catalog of the Cecidomyiidae (Diptera) of the World*. 4.ed. Digital. 762p. Disponível em: https://www.ars.usda.gov/ARSystemFiles/80420580/Gagne_2017_World_Cat_4th_ed.pdf.
- Gagné, R.J.; Oda, R.A.M. & Monteiro, R.E. The gall midges (Diptera: Cecidomyiidae) of *Mikania glomerata* (Asteraceae) in Southeastern Brazil. *Proceedings of the Entomological Society of Washington*, 103: 110-134.
- Girault, A.A. 1916. Descriptions of and observations on some chalcidoid Hymenoptera II. *The Canadian Entomologist*, 48: 337-344.
- Guimarães, A.L.A.; Cruz, S.M.S. & Vieira, A.C.M. 2014. Structure of floral galls of *Byrsonima sericea* (Malpighiaceae) induced by *Bruggmanniella byrsonimae* (Cecidomyiidae, Diptera) and their effects on host plants. *Plant Biology*, 16: 467-475.
- Hansen, R.W., Richard, R.D., Parker, P.E. & Wendel, E.L. 1997. Distribution of biological control agents of Leafy Spurge (*Euphorbia esula* L.) in the United States: 1988-1996. *Biological control*, 10: 129-142.
- Harris, K.M. 1966. Gall midge genera of economic importance (Diptera, Cecidomyiidae). Part 1: Introduction and subfamily Cecidomyiinae; supertribe Cecidomyiidi. *Transactions of the Royal Entomological Society of London*, 118: 313-358.
- Heeger, E. 1851. Beiträge zur Naturgeschichte der Insekten. Als Beiträge zur Fauna Oesterreichs. *Sitzungsberichte der mathematisch-naturwissenschaftlichen Classe der kaiserlichen Akademie der Wissenschaften*, 7: 203-217, pls. II-VII.
- Hempel, A. 1990. As coccidas brasileiras. *Revista do Museu Paulista*, 4: 365-537.
- Howard, L.O. 1881. Report on the parasites of Coccidae in the collection of this department. *Report of the United States Department of Agriculture, Entomology*, (1881): 350-372.
- Jiang, X. 1994. A checklist of gall midges of economic importance from Sichuan Province, China. *Entomological Knowledge*, 31: 245-247.
- Kalode, M.B. & Bentur, J.S. 1989. Characterization of Indian biotypes of the rice gall midge, *Orseolia oryzae* (Wood-Mason) (Diptera: Cecidomyiidae). *International Journal of Tropical Insect Science*, 10: 219-224.
- Kieffer, J.J. 1894. Description de quelques larves de Cécidomyes (suite). *Feuille des Jeunes Naturalistes*, 24: 119-121.
- Larson, K.C. 1998. The impact of two gall-forming arthropods on the photosynthetic rates of their hosts. *Oecologia*, 115: 161-166.
- Larson, K.C. & Whitham, T.G. 1991. Manipulation of food resources by a gall-forming aphid: the physiology of sink-source interactions. *Oecologia*, 88: 15-21.
- Linnaeus, C. 1758. *Systema Naturae per Regna Tria Naturae*. Edition 10, Vol. I. Stockholm. 824p.
- Loew, H. 1850. *Dipterologische Beiträge*. IV. Posen. 40p.
- Madeira, J.A.; Maia, V.C. & Monteiro, R.F. 2003. Gall makers (Cecidomyiidae, Diptera) on *Calophyllum brasiliense* Camb. (Clusiaceae): descriptions and biology. *Arquivos do Museu Nacional*, 61: 31-48.
- Maia, V.C. 1993. Descrição de duas espécies novas de Cecidomyiidae (Diptera) associadas a *Eugenia* spp. (Myrtaceae). *Revista Brasileira de Entomologia*, 37: 717-721.
- Maia, V.C. 1994. Considerações sobre *Stephomyia* Tavares (Diptera, Cecidomyiidae, Asphondyliidi), com descrição de quatro espécies novas associadas com *Eugenia* L. e *Neomithranthes obscura* (DC.) Legr. (Myrtaceae). *Revista Brasileira de Zoologia*, 10: 521-530.
- Maia, V.C. 1995. *Myrciariamyia bivalve*, gen. n. e sp. n. (Diptera, Cecidomyiidae, Oligotrophini), associado com *Myrciaria floribunda* (Camb.) Legr. (Myrtaceae) no Brasil. *Revista Brasileira de Zoologia*, 11: 635-638.
- Maia, V.C. 1996a. *Cordiamyia globosa* gen. n. e sp. n. (Diptera, Cecidomyiidae) associado com *Cordia verbenacea* DC. (Boraginaceae) no Brasil. *Revista Brasileira de Zoologia*, 13: 579-583.
- Maia, V.C. 1996b. Três espécies novas de *Dasineura* Rondani (Diptera, Cecidomyiidae) associadas a Myrtaceae, na restinga de Barra de Maricá, Rio de Janeiro. *Revista Brasileira de Zoologia*, 12: 1001-1008.
- Maia, V.C. 1996c. Dois gêneros novos de Cecidomyiidae (Diptera) associados a Myrtaceae, na restinga de Barra de Maricá, Rio de Janeiro, Brasil. *Revista Brasileira de Zoologia*, 12: 567-574.
- Maia, V.C. 1997. *Clusiamyia nitida* gen. n. e sp. n. (Diptera, Cecidomyiidae, Cecidomyiidi) associada com *Clusia lanceolata* Camb. (Clusiaceae) no Brasil. *Revista Brasileira de Zoologia*, 13: 829-832.
- Maia, V.C. 1999a. Artrópodes associados às galhas de Cecidomyiidae (Diptera) em *Eugenia rotundifolia* (Myrtaceae) e *Clusia lanceolata* (Clusiaceae) em uma restinga do Rio de Janeiro, Brasil. *Iheringia, Série Zoologia*, 87: 75-79.
- Maia, V.C. 1999b. Descrição de imaturos de quatro espécies de Asphondyliini neotropicais e nota taxonômica sobre *Asphondylia maytenuse* Maia & Couri (Diptera, Cecidomyiidae). *Revista Brasileira de Zoologia*, 16: 775-778.
- Maia, V.C. 2001a. The gall midges (Diptera, Cecidomyiidae) from three restingas of Rio de Janeiro State, Brazil. *Revista Brasileira de Zoologia*, 18: 583-629.
- Maia, V.C. 2001b. New genera and species of gall midges (Diptera, Cecidomyiidae) from three restingas of Rio de Janeiro State, Brazil. *Revista Brasileira de Zoologia*, 18(Supl. 1): 1-32.
- Maia, V.C. 2001c. Two new species of gall midges (Diptera, Cecidomyiidae) associated with *Pouteria caimito* var. *laurifolia* (Sapotaceae) in Brazil. *Studia Dipterologica*, 8: 103-110.
- Maia, V.C. 2005. *Clinodiplosis costai*, a new galler species (Diptera, Cecidomyiidae) associated with *Paullinia weinmanniaefolia* Mart (Sapindaceae). *Revista Brasileira de Zoologia*, 22: 676-679.
- Maia, V.C. 2010. A new species of *Dasineura* Rondani, 1840 (Diptera, Cecidomyiidae) associated with *Byrsonima sericea* (Malpighiaceae). *Revista brasileira de Biociências*, 8: 377-380.
- Maia, V.C. 2013. Galhas de insetos em restingas da região sudeste do Brasil com novos registros. *Biota Neotropica*, 13: 183-209.
- Maia, V.C. & Azevedo, M.A.P. 2009. Micro-himenópteros associados com galhas de Cecidomyiidae (Diptera) em Restingas do Estado do Rio de Janeiro (Brasil). *Biota Neotropica*, 9(2): 151-164.
- Maia, V.C. & Carvalho-Fernandes, S.P.C. 2011. Two new species of gall midges (Diptera, Cecidomyiidae) associated with *Erythroxylum ovalifolium* Peyr. (Erythroxylaceae) from the Barra de Maricá restinga, Maricá, Rio de Janeiro, Brasil. *Brazilian Journal of Biology*, 71: 521-526.
- Maia, V.C.; Constantino, P. de A.L. & Monteiro, R.F. New gall midges (Diptera, Cecidomyiidae) associated with two species of *Eugenia* (Myrtaceae). *Revista Brasileira de Entomologia*, 49: 347-352.
- Maia, V.C.; Couri, M.S. & Monteiro R.F. 1992. Sobre seis espécies de *Asphondylia* Loew, 1850 do Brasil (Diptera, Cecidomyiidae). *Revista Brasileira de Entomologia*, 36: 653-661.
- Maia, V.C. & Fernandes, G.W. 2004. Insect Galls from Serra de São José (Tiradentes, MG, Brazil). *Brazilian Journal of Biology*, 64: 423-445.
- Maia, V.C. & Fernandes, S.P.C. 2016. Insect galls of a protected remnant of the Atlantic Forest tableland from Rio de Janeiro State (Brazil). *Revista Brasileira de Entomologia*, 60: 40-56.
- Maia, V.C. & Mascarenhas, B. 2017. Insect Galls of the Parque Nacional do Itatiaia (Southeast Region, Brazil). *Anais da Academia Brasileira de Ciências*, 89(Supl. 1): 505-575.

- Maia, V.C. & Nava, D.E. 2011. New gall midges (Diptera, Cecidomyiidae) associated with *Eugenia uniflora* and *Psidium cattleianum* (Myrtaceae). *Iheringia, Série Zoológica*, 101: 69-74.
- Maia, V.C. & Oliveira, J.C. 2010. Galhas de insetos da Reserva Biológica Estadual da Praia do Sul (Ilha Grande, Angra dos Reis, RJ). *Biota Neotropica*, 10: 227-238.
- Maia, V.C. & Oliveira, U.P. 2007. Uma nova espécie de Cecidomyiidae (Diptera) associada com *Sebastiania glandulosa* (Euphorbiaceae). *Iheringia, Série Zoológica*, 97: 97-101.
- Maia, V.C. & Silva, L.O. 2016. Insect galls of Restinga de Marambaia (Barra de Guaratiba, Rio de Janeiro, RJ). *Brazilian Journal of Biology*, 76: 787-795.
- Maia, V.C. & Tavares, M. 2000. *Cordiamyia globosa* Maia (Diptera, Cecidomyiidae), flutuação populacional e parasitoides (Hymenoptera) associados. *Revista brasileira de Zoologia*, 17(3): 589-593.
- Maia, V.C.; Magenta, M.A.G. & Martins, S.E. 2008. Ocorrência e caracterização de galhas de insetos em áreas de restinga de Bertioiga (São Paulo, Brasil). *Biota Neotropica*, 8: 167-197.
- Maia, V.C.; Mendonça, M. de S. Jr. & Romanowski, H.P. 1996. *Eugeniomyia dispar* gen. n. e sp. n. (Diptera, Cecidomyiidae, Lasiopteridi) associated with *Eugenia uniflora* L. (Myrtaceae) in Brazil. *Revista Brasileira de Zoologia*, 13: 1087-1090.
- Maia, V.C.; Monteiro, R.F. & Narahara, K.L. 2002. Two new species of gall midges (Diptera, Cecidomyiidae) associated with *Protium icicaria* (Burseraceae) in Brazil. *Studia Dipterologica*, 9: 171-178.
- Mamaev, B.M. 1972. [New gall midges (Diptera, Cecidomyiidae) of the desert zone. *Entomologicheskoe Obozrenie*, 51: 886-899. Translated in *Entomological Review*, (1974) 51(4): 526-534.
- Mamaev, B.M. & Krivosheina, N.P. 1993. *The larvae of the gall midges (Diptera, Cecidomyiidae): comparative morphology, biology, keys*. Rotterdam, Balkema. 293p.
- Marini-Filho, O.J. & Fernandes, G.W. 2012. Stem galls drain nutrients and decrease shoot performance in *Diplusodon orbicularis* (Lythraceae). *Arthropod-plant interactions*, 6: 121-128.
- Massuti, L. & Vittorino, M.D. 1997. A new species of *Hyperaspis redtenbacher* (Coleoptera: Coccinellidae) and notes about the life habits. *The Coleopterists Bulletin*, 51: 213-216.
- McCrea, K.D.; Abrahamson, W.G. & Weis, A.E. 1985. Goldenrod ball gall effects on *Solidago altissima*: ¹⁴C translocation and growth. *Ecology*, 66: 1902-1907.
- Mendonça, M.S. & Romanowsky, H.P. 2002a. Life history of the gall-maker *Eugeniomyia dispar* Maia, Mendonça-Jr. & Romanowski, 1996 (Diptera, Cecidomyiidae). *Brazilian Journal of Biology*, 62: 277-283.
- Mendonça, M.S. & Romanowsky, H.P. 2002b. Natural enemies of the gall-maker *Eugeniomyia dispar* (Diptera, Cecidomyiidae): predatory ants and parasitoids. *Brazilian Journal of Biology*, 62: 269-275.
- Mendonça, M.S.; Toma, T.S.P. & Silva, J.S. 2014. Galls and Gallling Arthropods of Southern Brazil. In: Fernandes, G.W. & Santos, J.C. (Eds.). *Neotropical insect Galls*. New York, Springer. p. 221-256.
- MMA (Ministério do Meio Ambiente). 2018. *Mata Atlântica*. Disponível em: <http://www.mma.gov.br/biomas/mata-atlantica>.
- Möhn, E. 1959. Gallmücken (Diptera, Itonididae) aus El Salvador. 1. Teil. *Senckenbergiana Biologica*, 40: 297-368.
- Möhn, E. 1960. Gallmücken (Diptera, Itonididae) aus El Salvador. 2. Teil. *Senckenbergiana Biologica*, 41: 197-240.
- Monteiro, R.F.; Oda, R.A.M.; Narahara, K.L. & Constantino, A.L. 2004. Galhas: diversidade, especificidade e distribuição. In: Rocha, C.F.D.; Esteves, F.A. & Scarano, F.R. *Pesquisa de longa duração na Restinga de Jurubatiba: ecologia, história natural e conservação*. Macaé, NUPEM/UFRJ. p. 127-141.
- Monteiro, R.F.; Ferraz, F.F.F.; Maia, V.C. & Azevedo, M.A.P. 1994. Galhas entomógenas em restingas: uma abordagem preliminar. *ACIESP*, 3: 210-220.
- Moura, J.Z.; Borges, S.A.; Silva, P.R.R.; Maia, V.C. & Borges, T.I.A. 2010. Primeiro registro de galhas em panículas de cajazeira causadas por *Clinodiplosis* (Diptera: Cecidomyiidae) e avaliação de genótipos no estado do Piauí. *Revista Brasileira de Fruticultura*, 32: 921-924.
- Narahara, K.L.; Maia, V.C. & Monteiro, R.F. 2004. Two new species of gall midges (Diptera, Cecidomyiidae) associated with *Protium heptaphyllum* (Aubl.) Marchand (Burseraceae) in Brazil. *Revista Brasileira de Entomologia*, 48: 485-490.
- Nijveldt, W. 1969. *Gall midges of economic importance*. VIII: Miscellaneous. London, Crosby Lockwood & Son, Ltd. v. 8, 221p., 15 pls.
- Oliveira, J.C. & Maia, V.C. 2005. Ocorrência e caracterização de galhas de insetos na restinga de Grumari (Rio de Janeiro, RJ, Brasil). *Arquivos do Museu Nacional*, 63: 669-676.
- Oliveira, U.P. de & Maia, V.C. 2008. A new species of gall midge (Diptera, Cecidomyiidae) associated with *Sebastiania glandulosa* (Euphorbiaceae). *Arquivos do Museu Nacional*, 66: 355-358.
- Peschken, D. & Harris, P. 1975. Host specificity and biology of *Urophora cardui* (Diptera, Tephritidae). A biocontrol agent for Canada thistle (*Cirsium arvense*). *The Canadian Entomologist*, 107: 1101-1110.
- Price, P.W.; Fernandes, G.W. & Waring, G.L. 1987. Adaptive Nature of Insect galls. *Forum: Environmental Entomology*, 16: 15-24.
- Price, P.W.; Waring, G.L. & Fernandes, G.W. 1986. Hypotheses on the adaptive nature of galls. *Proceedings of the Entomological Society of Washington*, 88: 361-363.
- Proença, B. & Maia, V.C. 2015. New state record of gall midge species (Diptera, Cecidomyiidae) associated with *Calophyllum brasiliense* Cambess (Calophyllaceae). *Check List*, 10: 1557-559.
- Rizzini, C.T. 1997. *Tratado e fitogeografia do Brasil: Aspectos ecológicos, sociológicos e florísticos*. Rio de Janeiro, Âmbito Cultural Edições. 747p.
- Rodrigues, A.R. & Maia, V.C. 2010. Duas novas espécies de *Lopesia* Rübsaamen (Diptera, Cecidomyiidae) do Brasil, com chave para as espécies. *Biota Neotropica*, 10: 85-99.
- Rodrigues, A.R.; Maia, V.C. & Couri, M.S. 2014. Insect galls of restinga areas of Ilha da Marambaia, Rio de Janeiro, Brazil. *Revista Brasileira de Entomologia*, 58: 173-197.
- Rondani, C. 1840. *Sopra alcuni nuovi generi di insetti ditteri. Memoria seconda per servire Allá ditteologia italiana*. Parma. 27p., 1 pl.
- Rondani, C. 1860. *Stirpis cecidomyarum*. Genera revisa. Nota undecima, pro dipterologia italiana. *Atti della Società Italiana di Scienze Naturali*, 2: 286-294.
- Rübsaamen, E.H. 1905. Beitrage zur Kenntnis aussereuropaischer Zooecidien. II. Beitrag: Gallen aus Brasilien und Peru. (Vorlaufige Mitteilung.). *Marcellia*, 4: 65-85.
- Rübsaamen, E.H. 1908a. Beitrage zur Kenntnis aussereuropaischer Zooecidien. III. Beitrag: Gallen aus Brasilien und Peru. *Marcellia*, 6: 110-173.
- Rübsaamen, E.H. 1908b. Beitrage zur Kenntnis aussereuropaischer Zooecidien. III. Beitrag: Gallen aus Brasilien und Peru. *Marcellia*, 7: 15-79.
- Rübsaamen, E.H. 1915. Beitrag zur Kenntnis aussereuropaischer Gallmücken. *Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin*, 1915: 431-481.
- Sacchi, C.F.; Price, P.W.; Crai T.P. & Itami, J.K. 1988. Impact of the shoot galler attack on sexual reproduction in the arroyo willow. *Ecology*, 69: 2021-2030.
- Santos, M.G.; Fevereiro, P.C.A.; Reis, G.L.; Barcelos, J.I. & Ney, F.M.M.A. 2009. *Plantas da restinga: potencial econômico*. Rio de Janeiro, Technical Books Editora. 139p.
- Silva, S.H.A. & Maia, V.C. 2014. New state record of *Lopesia similis* Maia (Insecta, Diptera, Cecidomyiidae). *Check List*, 10: 1555-1556.

- Skuhravá, M. 1989. Taxonomic changes and records in Palaearctic Cecidomyiidae (Diptera). *Acta Entomologica Bohemoslovaca*, 86: 202-233.
- Skuhravá, M. 1997. Family Cecidomyiidae. In: Papp, L. & Darvas, B. (Eds.). *Contributions to a Manual of Palaearctic Diptera (with special reference to flies of economic importance)*. Vol. 2: Nematocera and Lower Brachycera. Budapest, Science Herald. p. 71-204.
- Sobhian, R.; Ryan, F.J.; Khamraev, A.; Pitcairn, M.J. & Bellb, D.E. 2003. DNA phenotyping to find a natural enemy in Uzbekistan for California biotypes of *Salsola tragus* L. *Biological Control*, 28: 222-228.
- Sousa, V.R. & Couri, M.S. 2014. Redescription of *Japanagromyza inferna* Spencer, first recorded from Brazil, and a key to the Neotropical species of *Japanagromyza* Sasakawa (Diptera, Agromyzidae). *Zookeys*, 374: 45-55.
- Souza, A.L.T.; Fernandes, G.W.; Figueira, J.E.C. & Tanaka, M.O. 1998. Natural history of a gall-inducing weevil *Collabismus clitellae* (Coleoptera: Curculionidae) and some effects on its host plant *Solanum lycocarpum* (Solanaceae) in southeastern Brazil. *Annals of the Entomological Society of America*, 91: 404-409.
- Spencer, K.A. & Stegmaier, C.E. Jr. 1973. Agromyzidae of Florida with a supplement on species from the Caribbean. In: *Arthropods of Florida and neighboring land areas*. Florida, Florida Department of Agriculture and Consumer Services. 205p.
- Tavares, J.S. 1909. Contributio prima ad cognitionem cecidologiae braziliae. *Brotéria, Série Zoológica*, 8: 5-28, pls. I-VIII.
- Tavares, J.S. 1916. Cecidomyias novas do Brazil. *Brotéria, Série Zoológica*, 14: 36-57.
- Tavares, J.S. 1920. Cecidologia brasileira: cecidias que se criam em plantas das familias das Leguminosae, Sapotaceae, Lauraceae, Myrtaceae, Punicaceae, Aurantiaceae, Malpighiaceae, Sapindaceae, Umbelliferae, Lorantheaceae, Apocynaceae, Urticaceae, Salicaceae e Gramineae. *Brotéria, Série Zoológica*, 18: 97-125.
- Tavares, J.S. 1921. Cecidologia brasileira. Cecídeas que se criam em plantas das familias das Leguminosae, Sapotaceae, Lauraceae, Myrtaceae, Punicaceae, Aurantiaceae, Malpighiaceae, Sapindaceae, Umbelliferae, Lorantheaceae, Apocynaceae, Urticaceae, Salicaceae e Gramineae. *Brotéria, Série Zoológica*, 19: 7 6-112, pl. I.
- Tavares, J.S. 1922. Cecidologia brasileira: as restantes famílias. *Brotéria, Série Zoológica*, 20: 5-48, pls. XI-XIX.
- Tavares, T.M.; Siebeneichler, S.C.; Cavalcanti, J.J.V.; Aférri, F.S.; Souza, C.M. & Nunes, T.V. 2011. Desempenho fenológico de progênies de meio irmãos de cajueiro anão precoce na região central do Tocantins no primeiro ano de plantio. *Revista Brasileira de Ciências Agrárias*, 1981-1160.
- Teske, M. & Trentini, A.M. 1997. *Compêndio de Fitoterapia Herbarium*. 3.ed. Curitiba, Ed. Herbarium Laboratório Botânico.
- Useful Tropical Plants. 2014. Disponível em: <http://tropical.theferns.info>. Access in: 24/11/2017.
- Viceconte, K.S. de M. & Maia, V.C. 2009. Novo gênero e nova espécie de Diptera, Cecidomyiidae, associada com *Psittacanthus dichrous* (Loranthaceae). *Arquivos do Museu Nacional*, 67: 35-40.
- Vitorino, D.V.; Pedrosa-Macedo, J.H. & Smith, C.W. 2000. The biology of *Tectococcus ovatus* Hempel (Heteroptera: Eriococcidae) and its potencial as a biocontrol agent of *Psidium cattleianum* (Myrtaceae). In: Spencer, N.R. (Eds.). *International Symposium on Biological Control of Weeds*, 10^o. Proceedings. Montana, Montana State University Bozeman. p. 651-657.
- Wachtl, F.A. 1882. Beitrage zur Kenntniss der gallenerzeugenden Insekten Europas. *Wiener Entomologische Zeitung*, 1: 289-293.
- Zanith, L.R. & Scarano, F.R. 2004. Produção de mudas de espécies das Restingas do município do Rio de Janeiro, RJ, Brasil. *Acta Botanica Brasílica*, 18(1): 161-17.