

Pilocarpus spp.: A survey of its chemical constituents and biological activities

Ana Paula Santos, Paulo Roberto Hrihorowitsch Moreno*

Laboratório de Química de Produtos Naturais, Departamento de Química Orgânica, Instituto de Química, Universidade de São Paulo

Pilocarpus species have been exploited as the only source of the imidazole alkaloid pilocarpine (used in glaucoma treatment), since its isolation up to the present. Almost all Pilocarpus species are native from Brazil. Because of the medicinal importance of pilocarpine, several of them are in the path of extinction. Other secondary metabolites, such as coumarins, flavonoids and terpenes, were described for Pilocarpus species. In this review the secondary metabolites, other than pilocarpine, isolated from Pilocarpus species and their biological activities were compiled. Although the variety of structures and the importance of the biological activities described in literature for Pilocarpus species this is an unexploited field of research in Natural Products and Pharmacology.

Uniterms

- Rutaceae
- *Pilocarpus* spp.
- Secondary metabolites
- Biological activities

*Correspondência:

P. R. H. Moreno
 Laboratório de Química de Produtos Naturais
 Instituto de Química, USP
 Av. Prof. Lineu Prestes, 748. Bloco 11 T
 Cidade Universitária
 05508-000 São Paulo – SP
 E-mail: prmoreno@iq.usp.br

INTRODUCTION

Pilocarpus Vahl is one of the 150 genus that compose the Rutaceae family and it was first known in Europe from the reports of the early scientific expeditions made to Brazil (Gabriel Soares de Souza, in 1570, Guilhaume Piso, in 1637, and Georg Marcgrave, in 1638).

Several *Pilocarpus* species are popularly known by the name of “jaborandi” which was derived from the tupi-Guarani language, *ya-mbor-endi* (the one who causes mouth-dripping) (Holmsted *et al.*, 1975). Several Rutaceae species are important due to their economical value as food, ornamental and medicinal uses. Some of the most common examples are listed in Table I.

TABLE I - Examples of the several uses of Rutaceae species

Species	Usage	References
<i>Citrus</i> spp.	Horticulture, volatile oils	Albuquerque, 1968
<i>Pilocarpus</i> spp.	Medicinal (pilocarpine)	Albuquerque, 1968
<i>Zanthoxylum</i> spp.	Carpentry	Corrêa, 1975
<i>Euxilophora paraensis</i> Huber	Carpentry	Loureiro, Silva, 1968
<i>Esenbeckia febrifuga</i> St. Hil.	Medicinal (febrifuge)	Albuquerque, 1968 Corrêa, 1969

The *Pilocarpus* genus was defined in 1797 with the descriptions of the species *P. racemosus*, however the position of the new taxon was not indicated (Skorupa, 1996). The initial conflicts in the systematic studies of the Rutaceae family imposed difficulties to classify this genus (Pirani, 1999; Albuquerque, 1985).

The most complete study about this genus was performed by Kaastra (1982) that kept the classification of *Pilocarpus* accordingly to Engler and recognized the genus *Raulinoa* Cowan as part of the subtribe Pilocarpineae (Scheme 1).

The *Pilocarpus* species are distributed in a large region of the American continent from Southern Central-America (Mexico) until the Southern part of the South-American continent (Argentina, Paraguay). There are 16 species described for this genus accordingly to Skorupa (1996). From these 13 can be found in Brazil, and 11 occur exclusively in the Brazilian territory (Table II).

These plants started to be studied concerning their pharmacological properties only in 1873, when it was confirmed that “jaborandi” was a potent stimulant of the secretory system, showing sialagogue, diuretic and diaphoretic activities (Corrêa, 1969). Since then, infusions of “jaborandi” leaves have been used for the treatment of fevers, stomatitis, bronchitis, gout, psoriasis, kidney diseases and several others illnesses (Holmsted *et al.*, 1975).

After the isolation of the imidazole alkaloid pilocarpine (Figure 1, compound **1**), in 1875 (Gerrard, 1875; Hardy, 1875), and its employment in ophthalmology (Weber, 1876), the “jaborandi” reached the status of one of the most important plants in the Brazilian flora (Pinheiro, 1997).

Besides alkaloids, a number of secondary metabolites have been reported in *Pilocarpus* species such

as coumarins (Amaro-Luis *et al.*, 1990), flavonoids (Bertrand *et al.*, 2001), hydrocarbons (Skorupa *et al.*, 1998; Negri *et al.*, 1998), triterpenes (Andrade Neto *et al.*, 1994) and volatile oils (Craveiro *et al.*, 1979).

The main goal of this review is to point out the importance of the *Pilocarpus* spp. as a source of secondary metabolites in Rutaceae with a potential pharmacological activity.

SECONDARY METABOLITES OF *PILOCARPUS* SPP.

Alkaloids

The “jaborandi” alkaloids are mainly of the imidazole group and it is thought that *L*-histidine is the biosynthetic precursor of the imidazole moiety (Battersby, Openshaw, 1953; Brochmann-Hanssen *et al.*, 1975; Cordell, 1981; Dewick, 2000).

Pilocarpine and other related alkaloids were isolated from several *Pilocarpus* species (Table III). Although the structural resemblance, only pilocarpine posses CNS activity, as a direct cholinergic which stimulates the parasympathetic system (bladder, tear ducts, sudoriferous and salivary glands). This alkaloid is the elected drug in the glaucoma treatment (Korolkovas, 1996) and the sialagogue property of pilocarpine has been exploited to treat the xerostomy (dry mouth) caused by the radio- or chemotherapy of throat cancer or those in the head area (Miller, 1993; Valdez *et al.*, 1993).

Others alkaloids derived from *L*-tryptophan were also isolated from *Pilocarpus* species. Some examples are *N,N*-dimethyl-5-methoxy-triptamine (**10**) and *N,N*-dimethyl-triptamine (**11**) (known as DMT) obtained from

Family	Subfamily	Tribe	Subtribe	Genus
Rutaceae	Rutoideae	Galipeae	Pilocarpineae	<i>Pilocarpus</i>
		(Cusparieae)		<i>Essenbeckia</i>
	Dictyolomatoideae	Zanthoxyleae	<i>Metrodorea</i>	
	Spathelioideae	Ruteae	<i>Raulinoa</i>	
	Toddalioideae	Boroneae	<i>Galipea</i> *	
	Flindersioideae	Diosmeae	<i>Raputia</i> *	
	Citroideae			<i>Neoraputia</i> *

* cited by PIRANI (1999)

SCHEME 1 - Systematic classification of *Pilocarpus* genus in Rutaceae.

TABLE II - Distribution of *Pilocarpus* species (Skorupa, 1996)

Species	Localization
<i>Pilocarpus alatus</i> C. J. Joseph ex Skorupa	Brazil (Pará and Mato Grosso)
<i>Pilocarpus carajaensis</i> Skorupa	Brazil (Pará)
<i>Pilocarpus demerarae</i> Sandwith	Guiana Inglesa
<i>Pilocarpus giganteus</i> Engler	Brazil (Bahia, Espírito Santo, Minas Gerais, Rio de Janeiro and São Paulo)
<i>Pilocarpus grandiflorus</i> Engler	Brazil (Alagoas, Bahia and Espírito Santo)
<i>Pilocarpus jaborandi</i> Holmes	Brazil (Ceará and Pernambuco)
<i>Pilocarpus manuensis</i> Skorupa	Peru (Manu National Park)
<i>Pilocarpus microphyllus</i> Stapf ex Wardleworth	Brazil (Maranhão, Mato Grosso, Pará and Piauí), Surinam
<i>Pilocarpus pauciflorus</i> St. Hilaire	Brazil (Paraná, Santa Catarina, Rio de Janeiro and São Paulo) (subSpecies <i>pauciflorus</i>) Brazil (Bahia) (subspecies <i>clavatus</i>) Brazil (subSpecies <i>organensis</i>)
<i>Pilocarpus pennatifolius</i> Lemmaire	Brazil (Bahia, Goiás, Mato Grosso and Mato Grosso do Sul, Minas Gerais, Paraná, Pernambuco, Rio Grande do Sul, Santa Catarina and São Paulo), Paraguay (Caazapá, Paraguari), Argentina (Misiones)
<i>Pilocarpus peruvianus</i> (Macbride) Kaastra	Bolivia, Brazil (Acre, Rodônia e Mato Grosso) and Peru
<i>Pilocarpus racemosus</i> Vahl	Mexico, Antilles, Colombia, Venezuela and British Guyana (subspecies <i>racemosus</i>) Venezuela and Colombia (subspecies <i>goudotianus</i>) Costa Rica and El Salvador (subspecies <i>viridulus</i>)
<i>Pilocarpus riedelianus</i> Engler	Brazil (Alagoas, Bahia, Espírito Santo, Paraíba and Pernambuco)
<i>Pilocarpus spicatus</i> St. Hilaire	Brazil (Bahia, Espírito Santo, Rio de Janeiro and São Paulo) (subspecies <i>spicatus</i>) Brazil (Bahia) (subspecies <i>longeracemosus</i>) Brazil (Ceará, Paraíba and Pernambuco) (subspecies <i>aracatensis</i>)
<i>Pilocarpus sulcatus</i> Skorupa	Brazil (Bahia and Minas Gerais)
<i>Pilocarpus trachyllophus</i> Holmes	Brazil (Bahia, Ceará, Minas Gerais and Piauí)

P. organensis leaves; both have a CNS activity causing hallucinations (Balsam, Voigtländer, 1978).

Dictamine (**14**) a quinoline alkaloid obtained from *P. grandiflorus* posses a structural relationship with coumarins and, probably because of this resemblance, this substance exhibited a photosensitizing activity (Ashwood *et al.*, 1982).

Coumarins

Coumarins are secondary metabolites biosynthesized from *p*-coumaric acid (a phenylalanine derived). Besides the simple coumarins, like scopoletin (**28**), in *Pilocarpus* species are also found pyrano- or furanocoumarins either with linear — xantyletin (**13**), chalepin (**32**) and bergapten (**20**) — or with an angular

structure — allo-xantoxillettol (**17**) and 3-methoxy-angelicin (**24**). Some coumarins found in *Pilocarpus* species may also be *O*-prenylated — donatin (**18**), imperatorin (**23**) — or *C*-prenylated — osthole (**30**).

Although studies on the isolation of secondary products from *Pilocarpus* sp. have been performed for a long time, the isolation of coumarins has only been reported just over a decade ago from the leaves of *P. goudotianus* (Amaro-Luis *et al.*, 1990). The efficacy of the jaborandi infusions in the treatment of skin diseases such as psoriasis and vitiligo might be related with this class of natural products. Nowadays, several coumarins (particularly the 8-methoxy-psoralen (**21**)) have been used for skin repigmentation (Manderfeld *et al.*, 1997).

Some coumarins also showed good results as germination inhibitors (MACÍAS *et al.*, 1993), insecticide

TABLE III - Alkaloids derived from *Pilocarpus* species and some related biological activities

Alkaloid	Species	Activities	References
pilocarpine (1)	<i>P. jaborandi</i>	Sialagogue	Gerrard, 1875
	<i>P. microphyllus</i>	Diuretic	Hardy, 1875
	<i>P. pennatifolius</i>	Febrifuge	Jowett, 1900
	<i>P. racemosus</i>		Mester, 1973
	<i>P. trachyllophus</i>		Payo Hill <i>et al.</i> , 1995
	<i>P. riedelianus</i>		Andrade Neto <i>et al.</i> , 1996 Guerreiro <i>et al.</i> , 2003
isopilocarpine (2)	<i>P. jaborandi</i>	-	Gerrard, 1875
	<i>P. microphyllus</i>		Hardy, 1875
	<i>P. pennatifolius</i>		Mester, 1973
pilocarpidine (3)	<i>P. jaborandi</i>	-	Jowett, 1900
isopilocarpidine (4)	<i>P. jaborandi</i>	-	Jowett, 1900
pilosine (5)	<i>P. jaborandi</i>	-	Pyman, 1912
	<i>P. microphyllus</i>		Voightländer, Rosenberg, 1959 Tedeschi <i>et al.</i> , 1973
isopilosine (6)	<i>P. jaborandi</i>	-	Voightländer, Rosenberg, 1959
epiisopilosine (7)	<i>P. jaborandi</i>	-	Tedeschi <i>et al.</i> , 1973
epiisopiloturine (8)	<i>P. jaborandi</i>	-	Tedeschi <i>et al.</i> , 1973
13-nor-7(11)-dehidro-pilocarpine (9)	<i>P. trachyllophus</i>	-	Andrade Neto <i>et al.</i> , 1996
<i>N,N</i> -dimethyl-5-methoxy-triptamine (10)	<i>P. organensis</i>	Hallucinogen	Balsam, Voightländer, 1978
			Riba <i>et al.</i> , 2002
<i>N,N</i> -dimethyl-triptamine (11)	<i>P. organensis</i>	Hallucinogen	Balsam, Voightländer, 1978
			Riba <i>et al.</i> , 2002
			De Souza <i>et al.</i> , 2003
plastydesmine (12)	<i>P. grandiflorus</i>	-	De Souza <i>et al.</i> , 2003
(1 <i>H</i>)-4-methoxy-2-quinolone (13)	<i>P. grandiflorus</i>	-	De Souza <i>et al.</i> , 2003
dictamine (14)	<i>P. grandiflorus</i>	Antifungic	Asswood-Smith <i>et al.</i> , 1982
		Vasorelaxing	Yu <i>et al.</i> , 1992
		Photosensitizing	De Souza <i>et al.</i> , 2002
		Anticoagulant	Su <i>et al.</i> , 1998

(Calcagno *et al.*, 2002; Ngwedson *et al.*, 2003; Manderfeld *et al.*, 1997), antioxidant (Shaw *et al.*, 2003; Hoult, Paya, 1996), and against Chagas' disease (Pavão *et al.*, 2002; Mafezoli *et al.*, 2000). Synthetic coumarins derived from seselin (35) exhibit potential activity against HIV (Huang *et al.*, 1994).

The known coumarins isolated from *Pilocarpus* species and their respective structures are listed in Table IV and Figure 2.

Flavonoids

Flavonoids represent one of the most important groups of phenolic natural products and, like coumarins, they are also biosynthesized from *L*-phenylalanine. These metabolites are obtained from *Citrus* species (Rutaceae)

from seeds, fruits, flowers and leaves (Middleton, 1984). Among others, the suggested activities of flavonoids are: analgesic (Thirugnanasambanthan *et al.*, 1990), sedative (Medina *et al.*, 1997; Marder & Paladini, 2002), anti-inflammatory (Pathak *et al.*, 1991).

The antioxidant activity of these substances is due to the ability of reducing free radical formation and to scavenge free radicals. Accordingly to Makersby (1997), the neurodegenerative Alzheimer's disease manifestation is related to the production of these species in the organism. In this sense, flavonoids can be considered as promising drugs in the treatment of this disease (Kim *et al.*, 2002).

Hesperidin (46) was isolated from *P. trachyllophus* (Bertrand *et al.*, 2001). This substance is currently isolated from other Rutaceae, such as *Citrus* and *Zanthoxylum*




FIGURE 1 – Chemical structures of the alkaloids isolated from *Pilocarpus* species.

species (Kanes *et al.*, 1993; Horowitz, Gentili, 1963; Arthur *et al.*, 1956). This flavonoid has various pharmacological activities and some of them are anti-inflammatory, bactericidal, fungicidal, antiviral, abortive, analgesic and antipyretic (Garg *et al.*, 2001). Recently, hesperidin was obtained from *Valeriana* species (“the Valium of 19th century”) and also exhibited sedative and sleep-enhancer properties (Marder *et al.*, 2003).

The following flavonoids, listed in Table V, were isolated from *P. trachyllophus* leaves.

Terpenes

Several reports dealt with the terpenes present in *Pilocarpus* species, mostly in volatile oils. Terpenes are biosynthesized from IPP (isopentenyl diphosphate),

TABLE IV - Coumarins isolated from *Pilocarpus* species and some related biological activities

Coumarin	Species	Activity	References
xantyletin (15)	<i>P. goudotianus</i>	Antifungic	Amaro-Luis <i>et al.</i> , 1990
	<i>P. racemosus</i>	Anti-PAF	Teng <i>et al.</i> , 1992
	<i>P. riedelianus</i>	Allelochemical	Macías <i>et al.</i> , 1993
xanthoxyletin (16)	<i>P. goudotianus</i>	Insecticide	Calcagno <i>et al.</i> , 2002
		Anti-PAF	Amaro-Luis <i>et al.</i> , 1990
		Allelochemical	Teng <i>et al.</i> , 1992
		Insecticide	Macías <i>et al.</i> , 1993
luvangetin (17)	<i>P. goudotianus</i> <i>P. racemosus</i>	Allelochemical	Amaro-Luis <i>et al.</i> , 1990
		Insecticide	Compagnone, Rodriguez, 1993
			Macías <i>et al.</i> , 1993
donatin (18)	<i>P. goudotianus</i>	Allelochemical	Calcagno <i>et al.</i> , 2002
		Insecticide	Amaro-Luis <i>et al.</i> , 1990
			Macías <i>et al.</i> , 1993
allo-xanthoxillettol (19)	<i>P. goudotianus</i>	Allelochemical	Calcagno <i>et al.</i> , 2002
		Insecticide	Amaro-Luis <i>et al.</i> , 1990
			Macías <i>et al.</i> , 1993
bergapten (20)	<i>P. goudotianus</i> <i>P. racemosus</i>	Allelochemical	Calcagno <i>et al.</i> , 2002
		Photosensitizing	Amaro-Luis <i>et al.</i> , 1990
		Bactericide	Compagnone, Rodriguez, 1993
		Insecticide	Macías <i>et al.</i> , 1993
8-methoxy-psoralen (21)	<i>P. goudotianus</i> <i>P. racemosus</i>	Allelochemical	Manderfeld <i>et al.</i> , 1997
		Insecticide	Calcagno <i>et al.</i> , 2002
		Photosensitizing	Amaro-Luis <i>et al.</i> , 1990
		Vasorelaxing	Compagnone, Rodriguez, 1993
		Bactericide	Macías <i>et al.</i> , 1993
		Allelochemical	Manderfeld <i>et al.</i> , 1997
isopimpinellin (22)	<i>P. goudotianus</i>	Insecticide	Chiou <i>et al.</i> , 2001
			Calcagno <i>et al.</i> , 2002
		Vasorelaxing	Amaro-Luis <i>et al.</i> , 1990
		Bactericide	Macías <i>et al.</i> , 1993
		Allelochemical	Chiou <i>et al.</i> , 2001
		Insecticide	Calcagno <i>et al.</i> , 2002
imperatorin (23)	<i>P. goudotianus</i>		Ngwendson <i>et al.</i> , 2003
		Vasorelaxing	Amaro-Luis <i>et al.</i> , 1990
		Allelochemical	Su <i>et al.</i> , 1998
tricocline (24)	<i>P. goudotianus</i>		Chiou <i>et al.</i> , 2001
			Ngwendson <i>et al.</i> , 2003
(+)-elisin (25)	<i>P. goudotianus</i>	-	Amaro-Luis <i>et al.</i> , 1990
3,6-dimethoxy-angelicin (26)	<i>P. grandiflorus</i> <i>P. riedelianus</i>	-	Amaro-Luis <i>et al.</i> , 1990
		-	De Souza <i>et al.</i> , 2003
		-	Müller <i>et al.</i> , 1993
3-methoxy-angelicin (27)	<i>P. riedelianus</i>	-	Müller <i>et al.</i> , 1993
scopoletin (28)	<i>P. riedelianus</i> <i>P. grandiflorus</i>	Antioxidant	Müller <i>et al.</i> , 1993
			De Souza <i>et al.</i> , 2002
			Shaw <i>et al.</i> , 2003
pimpinellin (29)	<i>P. riedelianus</i>	-	Müller <i>et al.</i> , 1993
osthole (30)	<i>P. riedelianus</i>	Anticoagulant	Müller <i>et al.</i> , 1993
		Vasorelaxing	Chiou <i>et al.</i> , 2001
meranzin hidrate (31)	<i>P. riedelianus</i>	-	Müller <i>et al.</i> , 1993
sphondin (32)	<i>P. riedelianus</i>	-	Müller <i>et al.</i> , 1993

TABLE IV - Coumarins isolated from *Pilocarpus* species and some related biological activities (continuation)

Coumarin	Species	Activity	References
auraptene (33)	<i>P. riedelianus</i>	Antitumoral	Müller <i>et al.</i> , 1993 Murakami <i>et al.</i> , 1997
chalepin (34)	<i>P. spicatus</i>	Anti-Chagas' disease	Pavão <i>et al.</i> , 2002
seselin (35)	<i>P. riedelianus</i>	Anti-HIV	Guerreiro <i>et al.</i> , 2001 Huang <i>et al.</i> , 1994
umbeliferone (36)	<i>P. riedelianus</i>	Antioxidant	Hoult, Paya, 1996 Guerreiro <i>et al.</i> , 2001
13,13-dimethyl-6 methoxy-2 <i>H</i> , 8 <i>H</i> -pyrano- (2,3)-cromen-2-one (37)	<i>P. riedelianus</i>	-	Guerreiro <i>et al.</i> , 2001
7-hydroxy-3-(1',1'-dimethyl-allyl)- 8-methoxy-coumarin (38)	<i>P. riedelianus</i>	Bactericide Antifungic	Guerreiro <i>et al.</i> , 2001
3'-hydroxy-dipetalactone (39)	<i>P. grandiflorus</i>	-	De Souza <i>et al.</i> , 2001
3',4'-dihydro-3'-hydroxy- dipetalactone (40)	<i>P. grandiflorus</i>	Antifungic	De Souza <i>et al.</i> , 2003
3-(1',1'-dimethyl-allyl)-scopoletin (41)	<i>P. grandiflorus</i>	Antifungic	De Souza <i>et al.</i> , 2003




FIGURE 2 - Chemical structures of some coumarins isolated from de *Pilocarpus* species.




FIGURE 2 – Chemical structures of some coumarins isolated from de *Pilocarpus* species (continuation).

TABLE V - Flavonoids isolated from *Pilocarpus* species and some related biological activities

Flavonoid	Species	Activity	References
3,4',5,7-tetrahydroxy-3',6-dimethoxy-flavone-3- <i>O</i> - β -rutinoside (42)	<i>P. trachyllophus</i>		Bertrand <i>et al.</i> , 2001
3,4',5,7-tetrahydroxy-6-methoxy-flavone-3- <i>O</i> - β -rutinoside (43)	<i>P. trachyllophus</i>		Bertrand <i>et al.</i> , 2001
patulentine-3- <i>O</i> - β -rutinoside (44)	<i>P. trachyllophus</i>	Antioxidant (aglicone)	Bertrand <i>et al.</i> , 2001
axilarin (45)	<i>P. trachyllophus</i>	Antioxidant	Bertrand <i>et al.</i> , 2001 Kim <i>et al.</i> , 2002
hesperidin (46)	<i>P. trachyllophus</i>	Antioxidant Sedative	Andrade Neto <i>et al.</i> , 1996 Kim <i>et al.</i> , 2002




FIGURE 3 - Chemical structures of flavonoids isolated from *Pilocarpus* species.

through the mevalonate pathway that gives rise to the mono- (C10), sesqui- (C15), di- (C20) and triterpenes (C30).

There are many references about the importance of terpenes in plant metabolism, mainly as essential oils. Although it is of general sense that terpenes represent a fundamental tool in pollinator's attraction, guaranteeing a species reproduction. They also displayed a variety of biologic activities such as insecticide, antimicrobial and antiseptic (Figueiredo, 1992). The essential oils of *P. spicatus*, *P. grandiflorus* and *P. pauciflorus* species exhibited bactericidal activity against *Bacillus subtilis*, *Salmonella typhimurium*, *Escherichia coli* and *Pseudomonas aeruginosa* (Santos *et al.*, 1997).

Although the occurrence of triterpenes and steroids in plants is quite common, and these compounds were only reported in three *Pilocarpus* species (Table VI and Figure 4). For *P. grandiflorus* the triterpenes b-amyrin (**109**), occotilone (**111**) and germanicol (**114**) were described. The b-amyrin acetate (**110**) obtained from *P. trachyllophus* showed a high cytostatic activity against cells of HEp-2 (Gomez *et al.*, 1997).

Additional pharmacological activities, such as angiogenic, antihelminthic, antimicrobial and antitumoral,

were described only for the steroid β-sitosterol (**86**) (Moon *et al.*, 1999; Villasenor *et al.*, 2002; Singh, Singh, 2003; Awad *et al.*, 2003).

Other secondary metabolites isolated from *Pilocarpus* species

Besides alkaloids, coumarins, flavonoids and terpenes there are minor compounds that can occur in some *Pilocarpus* species. Among these compounds are lignans, hydrocarbons and its alcohols, amides, carboxylic acids, esters and ketones derivatives (Table VII and Figure 5).

Lignans (also *L*-phenylalanine derivatives) were obtained from two different *Pilocarpus* species (Table VII). Lignans play an important role in the plant sustentation as cell wall constituents. The lignan liriioresinol B (**124**), isolated from *P. grandiflorus* (De Souza *et al.*, 2003), showed antichagasic activity (Sauvain *et al.*, 1996).

The amide pellitorine (**125**) was found in *P. trachyllophus* (Andrade Neto *et al.*, 1995). This substance is frequently isolated from *Piper* species and it is widely known as an insecticide (He *et al.*, 2002).

TABLE VI - Terpenoids isolated from *Pilocarpus* species

Terpene (class)	Species	References
<i>Monoterpenes</i>		
3,7,7-trimethylbicyclo[3.1.1]- 2-heptene (47)	<i>P. microphyllus</i> , <i>P. affinis</i>	Craveiro <i>et al.</i> , 1979
α -pinene (48)	<i>P. jaborandi</i> , <i>P. microphyllus</i> , <i>P. affinis</i> , <i>P. spicatus</i> , <i>P. grandiflorus</i>	Craveiro <i>et al.</i> , 1979 Santos <i>et al.</i> , 1997 Andrade Neto <i>et al.</i> , 2002
β -pinene (49)	<i>P. spicatus</i> , <i>P. grandiflorus</i>	Santos <i>et al.</i> , 1997 Andrade Neto <i>et al.</i> , 2002
canfene (50)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
borneol (51)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
linalool (52)	<i>P. spicatus</i> , <i>P. grandiflorus</i>	Santos <i>et al.</i> , 1997 Andrade Neto <i>et al.</i> , 2002
mircene (53)	<i>P. jaborandi</i> , <i>P. affinis</i> , <i>P. spicatus</i> , <i>P. grandiflorus</i>	Craveiro <i>et al.</i> , 1979 Santos <i>et al.</i> , 1997 Andrade Neto <i>et al.</i> , 2002
<i>E</i> - β -ocimene (54)	<i>P. spicatus</i> , <i>P. pennatifolius</i>	Andrade Neto <i>et al.</i> , 2002 Santos <i>et al.</i> , 2004
<i>Z</i> - β -ocimene (55)	<i>P. pennatifolius</i>	Santos <i>et al.</i> , 2004
Limonene (56)	<i>P. jaborandi</i> , <i>microphyllus</i> , <i>P. affinis</i> , <i>P. trachyllophus</i> , <i>P. spicatus</i> , <i>P. grandiflorus</i>	Craveiro <i>et al.</i> , 1979 Andrade Neto <i>et al.</i> , 1995 Santos <i>et al.</i> , 1997 Andrade Neto <i>et al.</i> , 2000, 2002
Sabinene (57)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
δ -3-carene (58)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
α -terpinene (59)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
γ -terpinene (60)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
α -terpineol (61)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
terpinen-4-ol (62)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
<i>p</i> -cimene (63)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
β -felandrene (64)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
1,8-cineole (65)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
<i>Sesquiterpenes</i>		
<i>Z,Z</i> -farnesol (66)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
<i>E</i> -nerolidol (67)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
β -bisabolene (68)	<i>P. trachyllophus</i> , <i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 1995 Santos <i>et al.</i> , 1997 Andrade Neto <i>et al.</i> , 2000, 2002
δ -cadinene (69)	<i>P. microphyllus</i> , <i>P. trachyllophus</i> , <i>P. spicatus</i> , <i>P. pennatifolius</i>	Craveiro <i>et al.</i> , 1979 Andrade Neto <i>et al.</i> , 1995 Andrade Neto <i>et al.</i> , 2000, 2002 Santos <i>et al.</i> , 2004
γ -cadinene (70)	<i>P. trachyllophus</i> , <i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 1995 Santos <i>et al.</i> , 1997 Andrade Neto <i>et al.</i> , 2000, 2002
germacrene D (71)	<i>P. trachyllophus</i> , <i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 1995 Andrade Neto <i>et al.</i> , 2002

TABLE VI - Terpenoids isolated from *Pilocarpus* species (continuation)

Terpene (class)	Species	References
<i>Sesquiterpenes</i>		
β -cubebene (72)	<i>P. spicatus</i> , <i>P. pennatifolius</i>	Andrade Neto <i>et al.</i> , 1995 Andrade Neto <i>et al.</i> , 2002 Santos <i>et al.</i> , 2004
γ -muurolene (73)	<i>P. spicatus</i> , <i>P. pennatifolius</i>	Andrade Neto <i>et al.</i> , 1995 Andrade Neto <i>et al.</i> , 2002 Santos <i>et al.</i> , 2004
α -muurolene (74)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
germacrene-4-D-ol (75)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
α -cadinol (76)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
cubenol (77)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
T-muurolol (78)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
calacorene (79)	<i>P. riedelianus</i>	Guerreiro <i>et al.</i> , 2001
α -calacorene (80)	<i>P. riedelianus</i>	Guerreiro <i>et al.</i> , 2001
γ -calacorene (81)	<i>P. riedelianus</i>	Guerreiro <i>et al.</i> , 2001
cadalene (82)	<i>P. riedelianus</i>	Guerreiro <i>et al.</i> , 2001
sesquichamaenol (83)	<i>P. riedelianus</i>	Guerreiro <i>et al.</i> , 2001
sesquichamaenol A (84)	<i>P. riedelianus</i>	Guerreiro <i>et al.</i> , 2001
δ -elemene (85)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
germacrene A (86)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
germacrene B (87)	<i>P. pennatifolius</i>	Santos <i>et al.</i> , 2004
selinene (88)	<i>P. affinis</i> , <i>P. pennatifolius</i>	Craveiro <i>et al.</i> , 1979 Santos <i>et al.</i> , 2004
β -elemene (89)	<i>P. spicatus</i> , <i>P. pennatifolius</i>	Andrade Neto <i>et al.</i> , 1995 Andrade Neto <i>et al.</i> , 2002 Santos <i>et al.</i> , 2004
bicyclogermacrene (90)	<i>P. spicatus</i> , <i>P. pennatifolius</i>	Andrade Neto <i>et al.</i> , 2002 Santos <i>et al.</i> , 2004
elemol (91)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
β -eudesmol (92)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
cyclosativene (93)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
caryophyllene oxide (94)	<i>P. trachyllophus</i> , <i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 1995 Santos <i>et al.</i> , 1997 Andrade Neto <i>et al.</i> , 2000, 2002
α -humulene (95)	<i>P. microphyllus</i> , <i>P. trachyllophus</i> , <i>P. spicatus</i> , <i>P. grandiflorus</i> , <i>P. pennatifolius</i>	Craveiro <i>et al.</i> , 1979 Andrade Neto <i>et al.</i> , 1995 Santos <i>et al.</i> , 1997 Andrade Neto <i>et al.</i> , 2000, 2002 Santos <i>et al.</i> , 2004
α -copaene (96)	<i>P. microphyllus</i> , <i>P. spicatus</i> , <i>P. pennatifolius</i>	Craveiro <i>et al.</i> , 1979 Andrade Neto <i>et al.</i> , 2002 Santos <i>et al.</i> , 2004
β -borbonene (97)	<i>P. trachyllophus</i> , <i>P. pennatifolius</i>	Andrade Neto <i>et al.</i> , 1995 Andrade Neto <i>et al.</i> , 2002 Santos <i>et al.</i> , 2004
β -caryophyllene (98)	<i>P. microphyllus</i> , <i>P. trachyllophus</i> , <i>P. spicatus</i> , <i>P. pennatifolius</i>	Craveiro <i>et al.</i> , 1979 Andrade Neto <i>et al.</i> , 1995 Santos <i>et al.</i> , 1997 Andrade Neto <i>et al.</i> , 2000, 2002 Santos <i>et al.</i> , 2004

TABLE VI - Terpenoids isolated from *Pilocarpus* species (continuation)

Terpene (class)	Species	References
<i>Sesquiterpenes</i>		
α -guaiaene (99)	<i>P. microphyllus</i>	Craveiro <i>et al.</i> , 1979
spathulenol (100)	<i>P. trachyllophus</i> , <i>P. pennatifolius</i>	Andrade Neto <i>et al.</i> , 1995 Andrade Neto <i>et al.</i> , 2000 Santos <i>et al.</i> , 2004
α -cubebene (101)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
allo-aromadendrene (102)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
globulol (103)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
viridiflorol (104)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
guaiol (105)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
bulnesol (106)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 2002
<i>Diterpenes</i>		
sandaracopimaradiene (107)	<i>P. jaborandi</i>	Craveiro <i>et al.</i> , 1979
angustifolin (108)	<i>P. riedelianus</i>	Guerreiro <i>et al.</i> , 2001
<i>Triterpenes</i>		
β -amyrin (109)	<i>P. grandiflorus</i>	De Souza <i>et al.</i> , 2003
β -amyrin acetate (110)	<i>P. trachyllophus</i>	Bertrand <i>et al.</i> , 2001
ocotillone (111)	<i>P. grandiflorus</i>	De Souza <i>et al.</i> , 2003
3- β -acetoxy-24-methyl-25-ethyl-20,24-epoxydammarane (112)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 1994
3-oxo-24-methyl-25-ethyl-20,24-epoxydammarane (113)	<i>P. spicatus</i>	Andrade Neto <i>et al.</i> , 1994
germanicol (114)	<i>P. grandiflorus</i>	De Souza <i>et al.</i> , 2003
<i>Steroids</i>		
β -sitosterol (115)	<i>P. grandiflorus</i>	De Souza <i>et al.</i> , 2003
β -sitosterone (116)	<i>P. grandiflorus</i>	De Souza <i>et al.</i> , 2003
7-oxo- β -sitosterol (117)	<i>P. grandiflorus</i>	De Souza <i>et al.</i> , 2003
stigmast-3,7-dione (118)	<i>P. grandiflorus</i>	De Souza <i>et al.</i> , 2003
7- β -hydroxy-stigmast-4-en-3-one (119)	<i>P. grandiflorus</i>	De Souza <i>et al.</i> , 2003
<i>Polyprenols</i>		
12-prenol (120)	<i>P. trachyllophus</i>	Bertrand <i>et al.</i> , 2001
[3 <i>E</i> , 9 <i>Z</i>]-13-prenol (121)	<i>P. trachyllophus</i>	Bertrand <i>et al.</i> , 2001
[3 <i>E</i> , 10 <i>Z</i>]-14-prenol (122)	<i>P. trachyllophus</i>	Bertrand <i>et al.</i> , 2001

The pattern of *n*-alkanes is considered an important taxonomic marker in plant classification accordingly to Eglinton *et al.* (1962). The *n*-alkanes identified in leaf epicuticular waxes of different *Pilocarpus* species with a carbon chain ranging from C₁₄ to C₃₃ (Skorupa *et al.*, 1998). An aromatic hydrocarbon (C₂₀) was detected only in *P. jaborandi* and this substance was considered a marker for the species (Negri *et al.*, 1998).

CONCLUDING REMARKS

In this review, information concerning the occurrence and pharmacological activities of secondary metabolites isolated from *Pilocarpus* spp were collected. Alkaloids, coumarins, flavonoids and terpenoids are current compounds to this genus (Guerreiro *et al.*, 2001). Compounds derived from *L*-phenylalanine are predominant in this genus.




FIGURE 4 – Chemical structures of some terpenoids isolated from *Pilocarpus* species.

Although the medicinal value of pilocarpine is widely recognized, but the importance of other secondary metabolites obtained from *Pilocarpus* species was not yet well described. The summary of the main bioactive compounds, other than pilocarpine, described for *Pilocarpus* species are listed below:

1. From the 16 known *Pilocarpus* species, alkaloids have only been described in eight species (almost 1 alkaloid/species) (Table III), and for the majority of them the bioactivity was not yet determined;

2. There are a few reports over the presence of coumarins in *Pilocarpus* spp., however the pharmacological activities of this class of compounds are numerous;

3. Flavonoids have been increasingly described as antioxidant drugs but these compounds were only found in *P. trachyllophus* (Table V);

4. Only two lignans were isolated from two *Pilocarpus* species and one of them has been described as an antichagasic agent (Table VII).

The use of plants in folk medicine has lead scientists




FIGURE 4 – Chemical structures of some terpenoids isolated from *Pilocarpus* species (continuation).




FIGURE 4 – Chemical structures of some terpenoids isolated from *Pilocarpus* species (continuation).




FIGURE 4 – Chemical structures of some terpenoids isolated from *Pilocarpus* species (continuation).

TABLE VII – Miscellaneous secondary metabolites isolated from *Pilocarpus* species and their respective biological activities

Compound	Species	Activity	References
(-)-episesamin (123)	<i>P. trachyllophus</i>	-	Andrade Neto <i>et al.</i> , 1995
lirioresinol B (124)	<i>P. grandiflorus</i>	Antichagásica	De Souza <i>et al.</i> , 2003 Sauvain <i>et al.</i> , 1996
pellitorine (125)	<i>P. trachyllophus</i>	Inseticida	Andrade Neto <i>et al.</i> , 1995 He <i>et al.</i> , 2002
eugenol (126)	<i>P. pennatifolius</i>	Inseticida Anestésica Antioxidante Antifúngica	Santos <i>et al.</i> , 2004 Park <i>et al.</i> , 2000 Ogata <i>et al.</i> , 2000 Priyadarsini <i>et al.</i> , 1998 Meepagala <i>et al.</i> , 2002
<i>N,N'</i> -dimethyl antranilic acid (127)	<i>P. grandiflorus</i>	-	De Souza <i>et al.</i> , 2003
vanilic acid (128)	<i>P. grandiflorus</i>	-	De Souza <i>et al.</i> , 2003
siringaldehyde (129)	<i>P. grandiflorus</i>	-	De Souza <i>et al.</i> , 2003
2-undecanone (130)	<i>P. spicatus</i> <i>P. trachyllophus</i> <i>P. jaborandi</i>	-	Santos <i>et al.</i> , 1997 Andrade Neto <i>et al.</i> , 1995 Andrade Neto <i>et al.</i> , 2000, 2002
2-tridecanone (131)	<i>P. spicatus</i> <i>P. trachyllophus</i> <i>P. jaborandi</i>	-	Santos <i>et al.</i> , 1997 Andrade Neto <i>et al.</i> , 1995 Andrade Neto <i>et al.</i> , 2000, 2002
1-hydroxy-2-tridecanone (132)	<i>P. trachyllophus</i>	-	Andrade Neto <i>et al.</i> , 1996
2-hydroxy-ethyl-glutarate (133)	<i>P. racemosus</i>	-	Compagnone & Rodriguez, 1993
methyl hexadecanoate (134)	<i>P. pennatifolius</i>	-	Santos <i>et al.</i> , 2004
hexanol (135)	<i>P. pennatifolius</i>	-	Santos <i>et al.</i> , 2004
undecanol (136)	<i>P. pennatifolius</i>	-	Santos <i>et al.</i> , 2004
1-phenyl-5-vinyl-5,9-dimethyl-decane (137)	<i>P. jaborandi</i>	-	Negri <i>et al.</i> , 1998 Skorupa <i>et al.</i> , 1998
tridecane (138)	<i>P. pennatifolius</i>	-	Santos <i>et al.</i> , 2004
pentadecane (139)	<i>P. pennatifolius</i>	-	Santos <i>et al.</i> , 2004
<i>n</i> -alcanes (140)	<i>P. alatus</i> , <i>P. carajaensis</i> , <i>P. giganteus</i> , <i>P. sulcatus</i> , <i>P. pauciflorus</i> , <i>P. jaborandi</i> , <i>P. grandiflorus</i> , <i>P. riedelianus</i> , <i>P. microphyllus</i>	-	Skorupa <i>et al.</i> , 1998

to find a wide variety of active substances of invaluable importance to mankind. Considering the variety of metabolites isolated from *Pilocarpus*, these species represent a vast research field to be exploited.

RESUMO

Pilocarpus spp.: Revisão sobre sua constituição química e atividades biológicas

Espécies de Pilocarpus têm sido exploradas como única

fonte do alcalóide imidazólico pilocarpina (utilizado no tratamento do glaucoma) desde o isolamento dessa substância até os dias atuais. A maioria das espécies de Pilocarpus conhecida é nativa do Brasil e, devido à importância medicinal que a pilocarpina possui e ao desmatamento, várias se encontram em risco de extinção. Outros metabólitos secundários entre os quais cumarinas, flavonóides e terpenos foram descritos em espécies desse gênero. Nesta revisão foram relacionados os metabólitos secundários isolados em diversas espécies de Pilocarpus bem como suas atividades biológicas. Apesar da varieda-




FIGURE 5 – Chemical structures of miscellaneous secondary metabolites isolated from *Pilocarpus* species.

de de estruturas e as importantes atividades biológicas já descritas na literatura para as outras classes de metabólitos secundários, ainda há um vasto campo de estudo para as espécies de Pilocarpus.

UNITERMOS: Rutaceae. Pilocarpus spp. Metabólitos secundários. Atividades biológicas.

REFERENCES

- ALBUQUERQUE, B.W.P. Rutaceae do Estado da Guanabara. *An. Acad. Bras. Ciênc.*, v. 40, n. 4, p. 500, 1968. Apud: ALBUQUERQUE, B.W.P. Rutaceae. In.: RIZZO J.A., org. Flora do Estado de Goiás. Goiânia: Editora da UFG, 1985. v. 6. 35 p.
- ALBUQUERQUE, B.W.P. Rutaceae. In: RIZZO J.A., org. Flora do Estado de Goiás. Goiânia: Editora da UFG, 1985. v. 6. 35 p.
- AMARO-LUIS, J.M.; MASSANET, G.M.; PANDO, E.; RODRIGUEZ LUIZ, F. New coumarins from *Pilocarpus goudotianus*. *Planta Med.*, v. 56, p. 304-306, 1990.
- ANDRADE NETO, M.; SILVEIRA, E.R.; BRAZ-FILHO, R.; GAMBARDELA, M.T.P.; SANTOS, R.H.A. 24-methyl-25-ethyl-dammarane derivatives from *Pilocarpus spicatus*. *Phytochemistry*, v. 35, n. 3, p. 739-743, 1994.
- ANDRADE NETO, M.; MAFEZOLI, J.; CUNHA, U.A.; SILVEIRA, S.C.; SILVEIRA, E.R. Volatile constituents of *Pilocarpus trachyllophus* Holmes. *J. Essent. Oil Res.*, v. 7, p. 671-674, 1995.
- ANDRADE NETO, M.; MENDES, P.H.; SILVEIRA, E.R. An imidazole alkaloid and other constituents from *Pilocarpus trachyllophus*. *Phytochemistry*, v. 42, n. 3, 885-887, 1996.
- ANDRADE NETO, M.; CUNHA, U.A.; MAFEZOLI, J.; SILVEIRA, E.R. Volatile constituents of *Pilocarpus trachyllophus* Holmes and *Pilocarpus jaborandi* Holmes (Rutaceae). *J. Essent. Oil Res.*, v. 12, p. 769-774, 2000.
- ANDRADE NETO, M.; CUNHA, U.A.; MAFEZOLI, J.; SILVEIRA, E.R. Volatile constituents of different populations of *Pilocarpus spicatus* Saint Hill. (Rutaceae) from the Northeast of Brazil. *J. Essent. Oil Res.*, v. 14, p. 319-324, 2002.
- ARTHUR, H.R.; WUI, W.H.; MA, C.N. An examination of the Rutaceae of Hong Kong (Part I), Flavonoid glycosides from *Zanthoxylum* species and occurrence of optically active hesperidin. *J. Chem. Soc.*, p. 632-635, 1956.
- ASHWOOD-SMITH, M.J.; TOWERS, G.H.; ABRAMOWSKI, Z.; POULTON, G.A.; LIU, M. Photobiological studies with dictamnine, a furoquinoline alkaloid. *Mutat. Res.*, v. 102, n. 4, p. 401-412, 1982.
- AWAD, A.B.; ROY, R.; FINK, C.S. b-sitosterol, a plant sterol induces apoptosis and activates key caspases in MDA-MB-231 human breast cancer cells. *Oncol. Rep.*, v. 10, n. 2, p. 497- 500, 2003.
- BALSAM, G.; VOIGTLÄNDER, H.W. A psychotropic alkaloid from *Pilocarpus organensis*. *Archiv. Pharm.*, v. 311, n. 78, p. 1016-1018, 1978.
- BATTERSBY, A.R.; OPENSHAW, H.T. *The imidazole alkaloids*. In: MANSKE, R.F.H.; HOLMES, H.L., eds. *The alkaloids*. San Diego: Academic Press, 1953. v. 20, p. 221-246.
- BERTRAND, C.; FABRE, N.; MOULIS, C. Constituents of *Pilocarpus trachyllophus*. *Fitoterapia*, v. 72, p. 844-847, 2001.
- BROCHMANN-HANSEN, E.; NUNES, M.A.; OLAH, C.K. On the biosynthesis of pilocarpine. *Planta Med.*, v. 28, p. 1-5, 1975.
- CALCAGNO, M.P.; COLL, J.; LLORIA, J.; FAINI, F.; ALONSO-AMELOT, M.E. Evaluation of synergism in feeding deterrence of some furanocoumarins on *Spodoptera littoralis*. *J. Chem. Ecol.*, v. 28, n. 1, 175-191, 2002.
- CHIOU, W.F.; HUANG, Y.L.; CHEN, C.F.; CHEN, C.C. Vasorelaxing effect of coumarins from *Cnidium monnieri* on rabbit corpus cavernosum. *Planta Med.*, v. 67, n. 3, p. 282-284, 2001.
- COMPAGNONE, R.; RODRIGUEZ, M.C. Coumarins from *Pilocarpus racemosus*. *Fitoterapia*, v. 64, n. 6, p. 557, 1993.
- CORDELL, G. *Introduction to alkaloids - a biogenic approach*. New York: John-Wiley & Sons, 1981. 1055 p.

- CORRÊA, M.P. *Dicionário de Plantas úteis do Brasil e das exóticas cultivadas*. Rio de Janeiro: Imprensa Nacional, 1969. v. 4. 374 p.
- CORRÊA, M.P. *Dicionário de Plantas úteis do Brasil e das exóticas cultivadas*. Rio de Janeiro: Imprensa Nacional, 1974. v. 5. 256 p.
- CRAVEIRO, A.A.; ANDRADE, C.H.S.; MATOS, F.J.A.; ALENCAR, J.W. Essential oils from Brazilian Rutaceae. I Genus *Pilocarpus*. *J. Nat. Prod.*, v. 42, n. 6, p. 669-671, 1979.
- DEWICK, P.M. *Medicinal natural products: a biosynthetic approach*. New York: John-Wiley & Sons, 2000. 466 p.
- EGLINTON, G.; HAMILTON, R.J.; RAPHAEL, R.A.; GONZALEZ, A.G. Hydrocarbon constituents of the wax coatings of plants leaves: a taxonomic survey. *Phytochemistry*, v. 1, p. 89-102, 1962.
- FIGUEIREDO, A.C.S. *Achillea millefolium* ssp. *millefolium*: produção de metabólitos secundários *in vivo* e *in vitro*. 1992. 208 p. [Tese de Doutorado, Universidade de Lisboa].
- GARG, A.; GARG, S.; ZANEVELD, L.J.D.; SINGLA, A.K. Chemistry and pharmacology of the *Citrus* bioflavonoid Hesperidin. *Phytother. Res.*, v. 15, p. 655-669, 2001.
- GERRARD, A.W. Alkaloid and active principle of jaborandi. *Pharmaceutical Journal*, v. 5, p. 865, 1875. Apud: HOLMSTED, B.; WASSEN, S.H.; SCHULTES, R.E. Jaborandi: An interdisciplinary appraisal. *J. Ethnopharmacol.*, v. 1, p. 3-21, 1979.
- GUERREIRO, G.; FERNANDES, J.B.; VIEIRA, P.C.; DA SILVA, M.F.G.F.; RODRIGUES FILHO, E.; VICTOR, S.R.; MAGALHÃES, C.; BUENO, O.C.; HEBLING, M.J.A.; PAGNOCCA, F.C. Sesquiterpenos e cumarinas bioativos de *Pilocarpus riedelianus*. In: REUNIÃO ANUAL DA SOCIEDADE BRASILEIRA DE QUÍMICA, 24., Poços de Caldas, 2001. *Livro de Resumos*. São Paulo: SBQ, 2001. PN-008.
- GOMEZ, M.A.; SÁENZ, M.T.; GARCIA, M.D.; AHUMADA, M.C.; DE LA PUERTA, R. Cytostatic activity against HEP-2 cells of methanolic extracts from *Viscum cruciatum* Sieber parasitic on *Crataegus monogyna* Jacq. and two isolated principles. *Phytother. Res.*, v. 11, p. 240-242, 1997.
- HE, W.; van PUYVELDE, L.; de KIMPE, N.; VERBRUGGEN, L.; ANTHONISSEN, K.; van der FLAAS, M.; BOSSLAERS, J.; MATHENGE, S.G.; MUDIDA, F.P. Chemical constituents and biological activities of *Zanthoxylum usambarense*. *Phytother. Res.*, v. 16, p. 66-70, 2002.
- HARDY, M.E., Sur le jaborandi (*Pilocarpus pinnatus*). *Bulletin Societe Chimie Paris*, v. 24, p. 497-500, 1875. Apud: HOLMSTED, B.; WASSEN, S.H.; SCHULTES, R.E. Jaborandi: An interdisciplinary appraisal. *J. Ethnopharmacol.*, v. 1, p. 3-21, 1979.
- HOLMSTED, B.; WASSEN, S.H.; SCHULTES, R.E. Jaborandi: An interdisciplinary appraisal. *J. Ethnopharmacol.*, v. 1, p. 3-21, 1979.
- HOROWITZ, R.M.; GENTILI, B. Flavonoids of *Citrus*. *Tetrahedron*, v. 19, p. 773-782, 1963.
- HOULT, J.R.; PAYA, M. Pharmacological and biochemical actions of simple coumarins: natural products with therapeutic potential. *Gen. Pharmacol.*, v. 27, n. 4, 713-722, 1996.
- HUANG, L.; KASHIWADA, Y.; COSENTINO, L.M.; FAN, S.; CHEN, C.H.; MCPHAIL, A.T.; FUJIOKA, T.; MIHASHI, K.; LEE, K.H. Anti-Aids agents 15. Synthesis and anti-HIV activity of dihydroseselins and related analogs. *J. Med. Chem.*, v. 37, n. 23, p. 3947- 3955, 1994.
- JOWETT, H.A.D. Pilocarpine and the alkaloids of Jaborandi leaves. *J. Chem. Soc.*, v. 77, p. 473-499, 1900.
- KAASTRA, R.C. New taxa and combinations in Rutaceae. *Acta Botanica Neerlandica*, v. 26, n. 6, p. 471-488, 1977. Apud: SKORUPA, L.A.; SALATINO, M.L.F.; SALATINO, A. Hydrocarbons of leaf epicuticular waxes of *Pilocarpus* (Rutaceae): Taxonomic meaning. *Biochem. Syst. Ecol.*, v. 26, p. 655-662, 1998.
- KANES, K.; TISSERAT, B.; BERHOW, M.; VANDERCOOK, C. Phenolic composition of various tissues of Rutaceae species. *Phytochemistry*, v. 324, p. 967-974, 1993.
- KIM, S.R.; PARK, M.J.; LEE, M.K.; SUNG, S.H.; PARK, E.J.; KIM, J.; KIM, S.Y.; OH, T.H.; MARKELONIS, G.J.; KIM, Y.C. Flavonoids of *Inula britannica* protect cultured cortical cells from necrotic cell death induced by glutamate. *Free Radical Biol. Med.*, v. 32, n. 7, p. 596-604, 2002.

- KOROLKOVAS, A. *Dicionário Terapêutico Guanabara*. Rio de Janeiro: Guanabara-Koogan, 1995/1996. p. 20.2-20.3.
- LOUREIRO, A.A.; SILVA, M.F. *Catálogo das madeiras da Amazônia*. Belém: SUDAM, 1968. v. 2, p. 261-266.
- MACÍAS, F.A.; GALINDO, J.C.C.; MASSANET, G.M.; RODRIGUEZ-LUIZ, F.; ZUBIA, E. Allelochemicals from *Pilocarpus goudotianus* leaves. *J. Chem. Ecol.*, v. 19, n. 7, p. 1371-1379, 1993.
- MAFEZOLI, J.; VIERA, P.C.; FERNANDES, J.B.; da SILVA, M.F.G.F.; DE ALBUQUERQUE, S. *In vitro* activity of Rutaceae species against the trypomastigote form of *Trypanosoma cruzi*. *J. Ethnopharmacol.*, v. 73, p. 335-340, 2000.
- MANDERFELD, M.M.; SCHAFER, H.W.; DAVIDSON, P.M.; ZOLLOTA, E.A. Isolation and identification of antimicrobial furocoumarins from parsley. *J. Food Prot.*, v. 60, n. 2, p. 72-77, 1997.
- MARDER, M.; PALADINI, A.C. GABA_A-receptor ligands of flavonoid structure. *Curr. Top. Med. Chem.*, v. 2, p. 853-863, 2002.
- MARDER, M.; VIOLA, H.; WASOWSKI, C.; FERNANDEZ, S.; MEDINA, J.H.; PALADINI, A.C. 6-Methylapigenin and hesperidin: new valeriana flavonoids with activity on the CNS. *Pharmacol. Biochem. Behavior*, v. 75, n. 3, p. 537-545, 2003.
- MAKERSBY, W.R. Oxidative stress hypothesis in Alzheimer's disease. *Free Radical Biol. Med.*, v. 23, p. 134-147, 1997.
- MEEPAGALA, K.M.; STURTZ, G.; WEDGE, D.E. Antifungal constituents of the essential oil fraction of *Artemisia dracunculul* L. var *dracunculul*. *J. Agric. Food Chem.*, v. 50, n. 24, p. 6989-6992, 2002.
- MEDINA, J.H.; VIOLA, H.; WOLFMAN, C.; MARDER, M.; WASOWSKI, C.; CALVO, D. Flavonoids: a new family of benzodiazepine receptor ligands. *Neurochem. Res.*, v. 22, n. 4, p. 419-425, 1997.
- MESTER, I. The occurrence of the alkaloids in Rutaceae. *Fitoterapia*, v. 44, p.123-152, 1973.
- MIDDLETON, E.J. The flavonoids. *Trends Pharmacol. Sci.*, v. 8, p.335-338, 1984.
- MILLER, L.J. Oral pilocarpine for radiation-induced xerostomia. *Cancer Bull.(Houston)*, v. 45, n. 6, p. 549-550, 1993.
- MOON, E.J.; LEE, Y.M.; LEE, O.H.; LEE, M.J.; LEE, S.K.; CHUNG, M.H.; PARK, Y.I.; SUNG, C.K.; CHOI, J.S.; KIM, K.W. A novel angiogenic factor derived from *Aloe vera* gel: b-sitosterol, a plant sterol. *Angiogenesis*, v.3, n. 2, p. 117- 123, 1999.
- MÜLLER, A.H.; DEGASPARI, L.R.O.; VIEIRA, P.C.; da SILVA, M.F.G.; FERNANDES, J.B.; PIRANI, J.R. 3-methoxyfuranocoumarins from *Pilocarpus riedelianus*. *Phytochemistry*, v. 34, n. 2, p. 585-586, 1993.
- MURAKAMI, A.; KUKI, W.; TAKAHASHI, Y.; YONEI, H.; NAKAMURA, Y.; OHTO, Y.; OHIGASHI, H.; KOSHIMIZU, K. Auraptene, a citrus coumarin, inhibits 12-O-tetradecanoylphorbol-13-acetate-induced tumor promotion in ICR mouse skin, possibly through suppression of superoxide generation in leukocytes. *Jan. J. Cancer Res.*, v. 88, n. 5, p. 443-452, 1997.
- NEGRI, G.; SALATINO, M.L.F.; SALATINO, A.; SKORUPA, L.A. An aromatic hydrocarbon from the foliar epicuticular wax of *Pilocarpus jaborandi*. *Phytochemistry*, v. 49, n. 1, p. 127-129, 1998.
- NGWENDSON, J.N.; BEDIR, E.; EFANGE, S.M.; OKUNJI, C.O.; IWU, M.M.; SCHUSTER, B.G.; KHAN, I.A. Constituents of *Peucedanum zenkeri* seeds and their antimicrobial effects. *Pharmazie*, v. 58, n. 8, p. 587-589, 2003.
- OGATA, M.; HOSHI, M.; URANO, S.; ENDO, T. Antioxidant activity of eugenol and related monomeric and dimeric compounds. *Chem. Pharm. Bull.*, v. 48, n. 10, p.1467-1469, 2000.
- PARK, I.K.; LEE, H.S.; LEE, S.G.; PARK, J.D.; AHN, Y.J. Insecticidal and fumigant activities of *Connamomun cassia* bark-derived materials against *Mechoris ursulus* (Coleoptera: attelabidae). *J. Agric. Food Chem.*, v. 48, n. 6, p. 2528-2531, 2000.
- PATHAK, D.; PATHAK, K.; SINGLA, A. Flavonoids as medical agents - recent advances. *Fitoterapia*, v. LXII, p. 317-389, 1991.

- PAVÃO F.; CASTILHO M.S.; PUPO M.T.; DIAS R.L.A.; CORREA A.G.; FERNANDES J.B.; SILVA M.F.G.F.; MAFEZOLI J.; VIEIRA P.C.; OLIVA G. Structure of *Trypanosoma cruzi* glycosomal glyceraldehyde-3-phosphate dehydrogenase complexed with chalepin, a natural product inhibitor, at 1.95 Å resolution. *FEBS Lett.*, v. 520, p. 13-17, 2002.
- PAYO HILL, A.; DOMINICIS, M.E.; OQUENDO, M.; SARDUY, R. Obtención de pilocarpina a partir de *Pilocarpus racemosus* Vahl. *Rev. Cubana Farm.*, 1995. Disponível em: www.infomed.sld.cu/revistas/far/vol29_2_95/far12295.htm. Acesso em: 22 abr. 2001.
- PINHEIRO, C.U.B. Jaborandi (*Pilocarpus* sp., Rutaceae): a wild species and its rapid transformation into a crop. *Econ. Bot.*, v. 51, n. 1, p.49-58, 1997.
- PIRANI, J.R. *Estudos taxonômicos em Rutaceae: Revisão de Helietta e Balfourodendron (Pteleinae). Análise cladística de Pteleinae. Sinópsese de Rutaceae do Brasil.* São Paulo. 1999. p. irreg. [Tese de Livre docência, Instituto de Biologia, Universidade de São Paulo, São Paulo].
- PYMAN, F.L. Pilocarpine: a new alkaloid from *Pilocarpus microphyllus*. *J. Chem. Soc.*, v. 101, p. 2206-2271, 1912.
- PRIYADARSINI, K.I.; GUHA, S.N.; RAO, M.N. Physico-chemical properties and antioxidant activities of methoxy phenols. *Free Radical Biol. Med.*, v. 24, n. 6, p. 933-941, 1998.
- RIBA, J.; RODRIGUEZ-FORNELLS, A.; BARBANOJ, M.J. Effects of ayahuasca on sensory and sensorimotor gating in humans as measured by P50 suppression and prepulse inhibition of the startle reflex, respectively. *Psychopharmacol.*, v. 165, n. 1, p. 18-28, 2002.
- SANTOS, F.A.; CUNHA, G.M.A.; VIANE, G.S.B.; RAO, V.S.N.; MANOEL, A.N.; SILVEIRA, E.R. Antibacterial activity of essential oils from *Psidium* and *Pilocarpus* species of plants. *Phytother. Res.*, v. 11, p. 67-69, 1997.
- SANTOS, A.P.; CORNÉLIO, M.L.; LIMBERGER, R.P.; APEL, M.A.; HENRIQUES, A.T.; MORENO, P.R.H. Analysis of the Volatile Oil from *Pilocarpus pennatifolius* Lemmaire (Rutaceae) Leaves by GC/MS. *Flavour Fragrance J.*, v. 19, n. 4, p. 325-326, 2004.
- SAUVAIN, M.; KUNESH, N.; POISSON, J.; GANTIER, J.C.; GAYRAL, P.; DEDET, J.P. Isolation of leishmanicidal triterpenes and lignans from the Amazonian liana *Dolicarpus dentatus* (Dilleniaceae). *Phytother. Res.*, v. 10, p. 1-4, 1996.
- SINGH, B.; SINGH, S. Antimicrobial activity of terpenoids from *Trichodesma amplexicaule* Roth. *Phytother. Res.*, v. 17, n. 7, p. 814- 816, 2003.
- SHAW, C.Y.; CHEN, C.H.; HSU, C.C.; CHEN, C.C.; TSAI, Y.C. Antioxidant properties of scopoletin isolated from *Sinomonium acutum*. *Phytother. Res.*, v. 17, n. 7, p. 823-825, 2003.
- SKORUPA, L.A. *Revisão taxonômica de Pilocarpus Vahl* (Rutaceae). São Paulo. 1996. 426 p. [Tese de doutoramento, Instituto de Biologia, Universidade de São Paulo].
- SKORUPA, L.A.; SALATINO, M.L.F.; SALATINO, A. Hydrocarbons of leaf epicuticular waxes of *Pilocarpus* (Rutaceae): Taxonomic meaning. *Biochem. Systemat. Ecol.*, v. 26, p. 655-662, 1998.
- De SOUZA, R.C.; FERNANDES, J.B.; VIEIRA, P.C.; da SILVA, M.F.G.F.; RODRIGUES FILHO, E.; PIRANI, J.R. Constituintes químicos de *Pilocarpus grandiflorus* (Rutaceae). In: REUNIÃO ANUAL DA SOCIEDADE BRASILEIRA DE QUÍMICA, 26., Poços de Caldas, 2003. *Livro de Resumos*. São Paulo: SBQ, 2003, PN-203.
- De SOUZA, R.C., FERNANDES, J.B., VIEIRA, P.C., Da SILVA, M.F.G.F., RODRIGUES FILHO, E., BUENO, O.C., HEBLING, M.J.A., ZANCUL, M.S.. Alcalóide furoquinolínico de *Pilocarpus grandiflorus* e sua ação sobre o fungo simbionte se formigas cortadeiras. In: REUNIÃO ANUAL DA SOCIEDADE BRASILEIRA DE QUÍMICA, 25., Poços de Caldas, 2002. *Livro de Resumos*. São Paulo: SBQ, 2002, PN-130.
- De SOUZA, R.C., FERNANDES, J.B., VIEIRA, P.C., Da SILVA, M.F.G.F., RODRIGUES FILHO, E., BUENO, O.C., HEBLING, M.J.A., ZANCUL, M.S. Estudo fitoquímico de *Pilocarpus grandiflorus* e sua ação sobre o fungo simbionte de formigas cortadeiras. In: REUNIÃO ANUAL DA SOCIEDADE BRASILEIRA DE QUÍMICA, 24., Poços de Caldas, 2001. *Livro de Resumos*. São Paulo: SBQ, 2001, PN-155.

- SU, T.L.; LIN, F.W.; TENG, C.M.; CHEN, K.T.; WU, T.S. Antiplatelet aggregation principles from the stem and root bark of *Melicope triphylla*. *Phytother. Res.*, v. 12, p. S74-S76, 1998.
- TEDESCHI, E.; KAMIONSKY, J.; FACKLER, S.; SAREL, S. The isolation of pure pilosine and epi-isopilosine from leaves of *Pilocarpus jaborandi*. *Israel J. Chem.*, v. 11, n. 5, p. 731-733, 1973.
- TENG, C.M.; LI, H.L.; YU, S.M.; WU, T.S.; HUANG, S.C.; PENG, I.S.; HUANG, T.F.A.S.I.A. Ca²⁺ channel blocking effects of 3 coumarin compounds, isolated from *Citrus grandis*, in rat thoracic aorta. *Pac. J. Pharmacol.*, v. 7, n. 2, p. 115-120, 1992.
- THIRUGNANASAMBANTHAN, P.; VISWANATHAN, S.; MYTHIRAYEE, C.; KRISHNAMURTY, V.; RAMACHANDRAN, S.; KAMESWARAN, L. Analgesic activity of certain flavone derivatives: a structure-activity study. *J. Ethnopharmacol.*, v. 28, p. 207-214, 1990.
- VALDEZ, I.H.; WOLFF, A.; ATKINSON, J.C.; MACYNSKY, A.A.; FOX, P.C. Use of pilocarpine during head and neck radiation therapy to reduce xerostomia and salivary dysfunction. *Cancer*, v. 71, n. 5, p.1848-1851, 1993.
- VILLASENOR, I.M.; ANGELADA, J.; CANLAS, A.P.; ECHEGOYEN, R. Bioactivity studies on b-sitosterol and its glucoside. *Phytother. Res.*, v. 16, n. 5, p. 417- 421, 2002.
- VOIGTLÄNDER, H.W.; ROSENBERG, W. Über der isopilosin und pilosin. *Arch. Pharm.*, v. 292, p. 579-585, 1959.
- WEBER, U. Ueber die Wirkung des Pilocarpium muriaticum. *Central. Med. Wiss.*, v. 44, p. 409-422, 1876. Apud: HOLMSTED, B.; WASSEN, S.H.; SCHULTES, R.E. Jaborandi: An interdisciplinary appraisal. *J. Ethnopharmacol.*, v. 1, p. 3-21, 1979.
- YU, S.M.; KO, F.N.; SU, M.J.; WU, T.S.; WANG, M.L.; HUANG, T.F.; TENG, C.M. Vasorelaxing effect in rat thoracic aorta caused by fraxinellone and dictamine isolated from the Chinese herb *Dictamnus dasycarpus* Turcz: comparison with cromakalim and Ca²⁺ channel blockers. *Naunyn Schmiedebergs Arch. Pharmacol.*, v. 345, n. 3, p. 349-555, 1992.

Recebido para publicação em 01 de março de 2004.