

Capgras syndrome in a first-episode, late-onset and super-refractory schizophrenia case

BRUNO T. JUNHO¹, ROBERTA R. PENNA¹, JOÃO VINÍCIUS SALGADO¹, LUCAS M. MANTOVANI¹

¹ Instituto Raul Soares, Belo Horizonte, MG, Brazil.

Institution where the study was conducted: Instituto Raul Soares. Avenida do Contorno, 3017 – 30110-080 – Santa Efigênia, Belo Horizonte, MG, Brazil.

Received: 04/21/2018 – Accepted: 04/27/2018

DOI: 10.1590/0101-60830000000162

Junho BT et al. / Arch Clin Psychiatry. 2018;45(3):77

Dear Editor,

Capgras syndrome (CS), late-onset schizophrenia (LOS) and refractory schizophrenia are relatively common conditions in psychiatric practice. However, the combination of these three conditions in a single case is a very rare event and so far seldom reported¹. We report a case of a patient with LOS presenting with CS and super-refractoriness since its first crisis.

A 53-year-old woman was admitted to a psychiatric hospital in an acute psychotic episode. On admission, she was convicted that her family had been enrolled in a plot in which they were replaced by lookalikes. This was the first time she presented psychotic symptoms and no treatment had yet been instituted. Physical examination, laboratory tests and brain MRI presented no abnormalities. She was diagnosed with late-onset schizophrenia (LOS) and Capgras Syndrome (CS). Haloperidol, risperidone and olanzapine, all in monotherapy, were tried at optimal doses for a period of four weeks each, with unsatisfactory response. Then, clozapine was started in a dose up to 425 mg daily leading to serum levels of 587 ng/mL (reference value: 50 to 700 ng/mL), with no clinical signs of response after 33 weeks. Subsequently, electroconvulsive therapy was prescribed as adjuvant therapy to clozapine. After 17 bitemporal sessions, psychosis with Capgras delusion persisted, maintaining high risk of aggression against her family and the need to remain hospitalized.

Schizophrenia is a relatively common disorder in clinical practice. Predominantly, it onsets in adolescence or early adult years, however 23,5% of these patients have this condition triggered after 40 years and are classified as LOS². This subtype schizophrenia have a preponderance of cases among women and the presence of schizoid and paranoid personality traits are frequent².

Several hypotheses have been proposed to understand schizophrenia, especially cases that occur with CS, a condition characterized by delusions that a close subject, usually parents or spouse, has been replaced by a lookalike. In CS, some studies

indicate the affection lack when in their relatives' presence. Delusion development would be an attempt to explain it³.

No complementary test is necessary for the schizophrenia diagnosis. In LOS, laboratory exams and brain images are advisable, especially in the elderly. Particularly, in this population, there is a higher incidence of potentially serious neurological conditions, such as stroke or dementia, which could also present with psychotic symptoms and be misdiagnosed as schizophrenia at first sight⁴.

Given the unsatisfactory response to two different antipsychotics, clozapine is an important option to be considered, despite its potential side effects. Nonresponders patients are classified as super-refractory. However, the literature lacks well-designed studies with large enough samples to draw subsequent guidelines. Also of note, electroconvulsive therapy proved to be effective in association to clozapine⁵.

This case highlights the unusual combination of three different conditions that is rarely reported in the literature¹, the occurrence of CS in a case of LOS and super-refractory schizophrenia since its first crisis. In addition, it is important to emphasize the need for more well-designed studies to shed light on the treatment of refractory schizophrenia.

References

1. Ain MK, Rosdinom R, Raynuha M. Capgras syndrome in a very late onset, treatment resistant schizophrenia. *Int Psychogeriatr*. 2015;27(9):1573-5.
2. Harris MJ, Jeste DV. Late-onset schizophrenia: an overview. *Schizophr Bull*. 1988;14(1):39-55.
3. Corlett PR, Taylor JR, Wang XJ, Fletcher PC, Krystal JH. Toward a neurobiology of delusions. *Prog Neurobiol*. 2010;92(3):345-69.
4. Reinhardt MM, Cohen CI. Late-life psychosis: diagnosis and treatment. *Curr Psychiatry Rep*. 2015;17(2):1.
5. Elkis H, Buckley PF. Treatment-Resistant Schizophrenia. *Psychiatr Clin North Am*. 2016;39(2):239-65.

