

Rev. Inst. Med. Trop. Sao Paulo
53(5):283-289, September-October, 2011
doi: 10.1590/S0036-46652011000500008

USE OF ELISA EMPLOYING HOMOLOGOUS AND HETEROLOGOUS ANTIGENS FOR THE DETECTION OF IgG AND SUBCLASSES (IgG1 AND IgG2) IN THE DIAGNOSIS OF CANINE VISCERAL LEISHMANIASIS

Flávia Coelho RIBEIRO(1), Armando de O. SCHUBACH(1), Eliame MOUTA-CONFORT(1), Tânia M.V. PACHECO(2), Maria de Fátima MADEIRA(1), Luiz Cláudio de Souza ABBOD(3), Carla de Oliveira HONSE(2), Andreia Silva ALVES(1) & Mauro C.A. MARZOCHI(1)

SUMMARY

Indirect immunofluorescence is the method recommended for the diagnosis of visceral leishmaniasis in dogs, however, the accuracy of this technique is low and its use on a large scale is limited. Since ELISA does not present these limitations, this technique might be an option for the detection of IgG or specific IgG1 and IgG2 subclasses. Canine ehrlichiosis is an important differential diagnosis of American Visceral Leishmaniasis (AVL). The present study compared ELISA using *Leishmania chagasi* and *Leishmania braziliensis* antigen for the detection of anti-*Leishmania* IgG and subclasses in serum samples from 37 dogs naturally infected with *L. chagasi* (AVL) and in samples from four dogs co-infected with *L. braziliensis* and *L. chagasi* (CI). The occurrence of cross-reactivity was investigated in control serum samples of 17 healthy dogs (HC) and 35 infected with *Ehrlichia canis* (EC). The mean optical density obtained for the detection of IgG was significantly higher when *L. chagasi* antigen was used, and was also higher in subgroup VLs (symptomatic) compared to subgroup VIa (asymptomatic). The correlation between IgG and IgG1 was low. The present results suggest that IgG ELISA using homologous antigen yields the best results, permitting the diagnosis of asymptomatic *L. chagasi* infection and the discrimination between cases of AVL and ehrlichiosis in dogs.

KEYWORDS: American visceral leishmaniasis; Dog; IgG subclasses; *L. (V.) braziliensis*, *L. (L.) chagasi*.

INTRODUCTION

Leishmania (Leishmania) chagasi is a viscerotropic parasite of lymphoid organs and the causative agent of American visceral leishmaniasis (AVL), also known as Kala-azar¹. AVL is found in various American countries. In Brazil, AVL presents a high incidence in urban and periurban areas³⁹. In the municipality of Rio de Janeiro, AVL emerged at the end of the 1970s in periurban areas of the northern zone of the city, especially at the hillside of the Pedra Branca massif³². The domestic dog (*Canis familiaris*) is the main reservoir of AVL due to intense parasitism of healthy skin and viscera since the onset of infection^{15,28,32,41,45}. It is believed that most infected dogs do not develop clinical signs and remain asymptomatic for variable periods of time^{19,32}. In Brazil, AVL requires obligatory notification and control programs involve the treatment of human cases, control of the insect vector, and euthanasia of seropositive infected dogs. The Brazilian Ministry of Health recommends serological testing by indirect immunofluorescence (IIF), with antibody titers of 1:40 or higher being used as a criterion for the culling of dogs³⁹.

The control measures adopted in Brazil are a matter of discussion since new AVL foci continue to appear in the country¹³. The relative

inefficiency of these measures has contributed to the emergence of AVL, especially because of the underestimated seroprevalence of canine infection provided by the IIF test⁵¹. Some authors suggest that a highly sensitive test, combined with rapid culling of the dogs, is necessary for a significant reduction in the transmission of the disease and effective control^{11,16}. According to BRAGA *et al.*⁸, only 35.4% of infected dogs identified by enzyme-linked immunosorbent assay (ELISA) would be detected by IIF. As a consequence, some infected dogs would not be identified during control actions and would remain in endemic areas, a fact facilitating propagation of the infection. In addition to its lower sensitivity and specificity when compared to ELISA³⁰, IIF is more time consuming and requires trained professionals^{5,17}. The rapidity and possibility of automation of ELISA are additional advantages that favor its use on a large scale.

Once clinical signals are installed, AVL should be differentiated from ehrlichiosis, lymphosarcoma, myeloma, seborrhea, pemphigus and systemic fungal infections, as well as from other diseases causing proliferation of the reticuloendothelial system²³. Ehrlichiosis is a disease caused by the rickettsia *Ehrlichia canis*, which primarily parasitizes cytoplasmic vacuoles of monocytes and granulocytes^{47,55}, causing a

(1) Laboratório de Vigilância em Leishmanioses, Instituto de Pesquisa Clínica Evandro Chagas, Fundação Oswaldo Cruz (FIOCRUZ), Rio de Janeiro, RJ, Brasil.

(2) Laboratório de Pesquisa Clínica em Dermatoozoonoses em Animais Domésticos, Instituto de Pesquisa Clínica Evandro Chagas, Fundação Oswaldo Cruz (FIOCRUZ), Rio de Janeiro, RJ, Brasil.

(3) Instituto Municipal de Medicina Veterinária Jorge Vaitsman.

Correspondence to: Flávia Coelho Ribeiro, Laboratório de Vigilância em Leishmanioses, Instituto de Pesquisa Clínica Evandro Chagas, Fundação Oswaldo Cruz, Av. Brasil 4365, 21045-900 Rio de Janeiro, RJ, Brasil. Tel:+55 21 3865 9594; +55 21 3865 9541 or +55 21 3865 9785. E-mail: fcribeiro@fiocruz.br

potentially fatal multisystemic infection that is characterized by clinical and hematological anomalies similar to those observed in AVL, including fever, lymphadenopathy, anorexia, lethargy and thrombocytopenia^{31,36,42}. *Ehrlichia canis* presents a wide geographic distribution and mainly occurs in tropical and subtropical areas. In Rio de Janeiro, ehrlichiosis is one of the most common causes of visits to veterinary clinicians, with a mean prevalence of the disease of 26.8% in thrombocytopenic dogs²⁷. Despite the clinical and epidemiological similarity to AVL, there is no indication for euthanasia of dogs as a public health measure. Thus, the test used for the serodiagnosis of canine AVL should be able to differentiate the two infections⁴⁸.

The IgG subclasses, whose expression depends on the nature of the antigen and antigenic stimulus, are involved in different biological activities. In dogs, four IgG subclasses have been defined^{33,34,35}, which can be applied to the diagnosis of infectious diseases, including leishmaniasis^{24,52}. The study of IgG subclasses in canine AVL might have a practical application both to the diagnosis and to the development of control strategies⁵⁰.

In the present study, we compared the results of ELISA using *L. (L.) chagasi* (homologous) and *L. (V.) braziliensis* (heterologous) antigen for the detection of anti-*Leishmania* IgG, IgG1 and IgG2 antibodies for the diagnosis of symptomatic and asymptomatic infection of dogs with *L. (L.) chagasi* and for the possible discrimination with cases of canine ehrlichiosis.

MATERIAL AND METHODS

Population and biological samples: Serum samples obtained from 93 domestic dogs were tested for leishmaniasis by total IgG and subclass IgG1 and IgG2 ELISA. The animals were classified into three well-defined groups: group VL (n = 37) included animals infected with *L. (L.) chagasi*. This group was subdivided into two subgroups: asymptomatic animals (VLa, n = 12), and symptomatic animals (VLs, n = 25) that presented two or more clinical signs of AVL: skin ulcerations, desquamation, adenitis, weight loss, alopecia, hepatosplenomegaly, onychogryphosis, anemia, fever, apathy, and keratoconjunctivitis. Group CI (n = 4) consisted of dogs co-infected with *L. (L.) chagasi* and *L. (V.) braziliensis*. All dogs included in groups VL and CI were from AVL-endemic areas in the municipality of Rio de Janeiro, tested seropositive by IIF, and were euthanized according to the municipal zoonosis control program. In all animals of the two groups, the diagnosis of infection was established by isolation of *Leishmania* from cultured viscera or skin and species characterization by isoenzyme electrophoresis²⁹.

The control group consisted of 52 dogs subdivided into two subgroups: healthy controls (HC, n = 17) and *E. canis* infected controls (EC, n = 35). Group HC included healthy dogs with no clinical alterations and normal blood count and biochemical parameters (protein, globulin, glutamate-oxaloacetate transaminase/aspartate aminotransferase (GOT/AST), glutamate-pyruvate transaminase/alanine aminotransferase (GPT/ALT), albumin, blood urea and creatinine) that lived outside AVL endemic areas and had a negative parasitological diagnosis of canine ehrlichiosis. Group EC included dogs with two or more clinical signs compatible with AVL but that lived outside AVL endemic areas and in which *E. canis* was visualized by direct parasitological examination of blood smears.

Indirect immunofluorescence: IIF for the detection of anti-*Leishmania* IgG was performed according to manufacturer instructions (Biomanguinhos/FIOCRUZ kit, Rio de Janeiro, Brazil). Titers $\geq 1:40$ were considered to be positive. Serum samples of dogs from the VL and CI groups (tested within the leishmaniasis control program as inclusion criterion for this study) were retested and serum samples of dogs from the HC and EC groups were tested for the first time.

Antigens used for ELISA: Partially soluble fractions of promastigote forms of *L. (V.) braziliensis* (MHOM/BR/75/M2903) and *L. (L.) chagasi* (MHOM/BR/74/PP75) obtained during the stationary phase of growth according to the protocol of RIBEIRO *et al.*⁴⁶ were used as antigens for ELISA.

ELISA: ELISA for the detection of anti-*Leishmania* IgG, IgG1 and IgG2 was carried out according to the method of VOLLER *et al.*⁵⁴ using *L. (V.) braziliensis* or *L. (L.) chagasi* antigen. The cut-off point was determined as described by RIBEIRO *et al.*⁴⁶. Briefly, the antigens were diluted as follows: *L. (V.) braziliensis* at concentrations of 5 $\mu\text{g}/\text{mL}$ for the detection of IgG and 7.5 $\mu\text{g}/\text{mL}$ for the detection of IgG1 and IgG2, and *L. (L.) chagasi* at concentrations of 10 $\mu\text{g}/\text{mL}$ for the detection of IgG and IgG1 and 2.5 $\mu\text{g}/\text{mL}$ for the detection of IgG2. Next, the serum samples were diluted 1:20 for *L. (V.) braziliensis* and 1:100 for *L. (L.) chagasi* antigen in a solution of 1% skim milk (Molico-Nestlé®) in PBS/0.05% Tween 20 (v/v). The plates were then incubated with peroxidase-conjugated anti-dog IgG (Sigma Chemical Co., St. Louis, MO, USA) and anti-dog IgG1 and IgG2 immunoglobulins (Bethyl Laboratories, Inc., Montgomery, TX, USA) at dilutions of 1:40,000, 1:10,000 and 1:30,000 for reactions with *L. (V.) braziliensis* antigen and at dilutions of 1:20,000, 1:5,000 and 1:30,000 for reactions with *L. (L.) chagasi* antigen, respectively.

Different concentrations of antigen and conjugates used were set independently for each class and subclass of IgG, using criteria for choosing the lowest concentration of antigen and greater dilution of the conjugate, which maximized the parameters of sensitivity and specificity of the test.

Statistical analysis: The cut-off was determined from the ROC curves calculated with the MedCalc statistical program (version 8.2.0.2). The nonparametric Spearman test was used for the comparison of the position and variation in optical density (OD) and their eventual correlation since the data did not follow a normal distribution. Mean OD values were compared by the paired Student t-test using the SPSS version 11.0 statistical software package. Differences at $p < 0.05$ were considered to be significant.

RESULTS

Indirect immunofluorescence: All sera from dogs of group VL reacted at different dilutions: four (10.8%) at a titer of 1:80, seven (18.9%) at titers ranging from 1:160 to 1:320, and 26 (70.3%) at a titer $\geq 1:640$. In the mixed infection group (CI), one serum sample reacted at a titer of 1:160 and three at a titer of 1:320. The sera of one dog in the HC group (5.9%) and two dogs (5.7%) in the EC group reacted at a titer of 1:40.

ELISA

Correlation between ODs in the different reactions: In all groups, high and significant Spearman correlations were observed between the

IgG and IgG2 values for all sera analyzed when *L. (V.) braziliensis* ($r = 0.86, p < 0.01$) or *L. (L.) chagasi* antigen ($r = 0.95, p < 0.01$) was used, respectively. In contrast, the correlation between IgG and IgG1 was low but significant when *L. (V.) braziliensis* ($r = 0.49, p < 0.01$) and *L. (L.) chagasi* antigen ($r = 0.45, p < 0.01$) was used.

In group VL, no significant correlation for the detection of IgG1 was observed between symptomatic and asymptomatic animals with either *L. (V.) braziliensis* ($r = 0.12, p = .35$) or *L. (L.) chagasi* ($r = 0.09, p = 0.46$) antigen.

Determination of the cut-off: The determination of the cut-off was established by ROC curve, considering all sera from infected (LVs, VLa, IC) and uninfected animals (HC and EC), choosing the point on the curve that maximizes sensitivity and specificity provided by the program MedCalc (version 8.2.0.2).

The cut-off value that permitted the best discrimination between the infected groups (VL and CI) and control animals (HC and EC) was determined from the respective ROC curves (Fig. 1 and 2) plotted for each ELISA reaction (IgG, IgG1 and IgG2) obtained with the *L. (V.) braziliensis* and *L. (L.) chagasi* antigens. The following cut-off values were obtained for the IgG, IgG1 and IgG2 reactions: 0.212, 0.081 and 0.347 for *L. (V.) braziliensis* antigen, and 0.32, 0.085 and 0.15 for *L. (L.) chagasi* antigen, respectively.

Comparison of the area under the ROC curve: No significant differences were observed when comparing the areas under the curve (AUC) for the detection of IgG and IgG2 using either *L. (V.) braziliensis* or *L. (L.) chagasi* as antigen ($p = 0.85$) (Table 1). Significantly different AUCs ($p < 0.05$) were observed when comparing IgG and IgG2 with IgG1 for both *L. (V.) braziliensis* and *L. (L.) chagasi* antigen, respectively.

OD values: The following mean OD values and their standard deviations were obtained for the VL group: 0.811 ± 0.142 for IgG, 0.124 ± 0.061 for IgG1 and 1.429 ± 0.426 for IgG2 when *L. (V.) braziliensis* antigen was used, respectively, and 1.728 ± 0.426 , 0.239 ± 0.253 and




Fig. 1 - Distribution of OD values in ELISA reactions (IgG) using *L. (V.) braziliensis* antigen obtained for the visceral leishmaniasis group (VL) (symptomatic subgroup, VLs, and asymptomatic subgroup, VLa), group co-infected with *L. (L.) chagasi* and *L. (V.) braziliensis* (CI), and control group (healthy subgroup, HC, and ehrlichiosis subgroup, EC).




Fig. 2 - Distribution of OD values in ELISA reactions (IgG) using *L. (L.) chagasi* antigen obtained for the visceral leishmaniasis group (VL) (symptomatic subgroup, VLs, and asymptomatic subgroup, VLa), group co-infected with *L. (L.) chagasi* and *L. (V.) braziliensis* (CI), and control group (healthy subgroup, HC, and ehrlichiosis subgroup, EC).

Table 1
Area under the curve values for IgG, IgG1 and IgG2 for antigens of *L. braziliensis* and *L. chagasi*.

<i>L. braziliensis</i>	Area under the curve	<i>L. chagasi</i>	Area under the curve
IgG	1,000	IgG	1,000
IgG1	0.913	IgG1	0.858
IgG2	1,000	IgG2	1,000

1.177 ± 0.337 when *L. (L.) chagasi* antigen was used. The mean OD values obtained for the detection of IgG (Fig.1 and 2) and IgG1 were significantly higher when *L. (L.) chagasi* was used as antigen ($p < 0.01$).

The seroprevalence of IgG and IgG2 was 100% in both symptomatic and asymptomatic dogs, irrespective of the antigen used. When using *L. (L.) chagasi* as antigen to compare symptomatic and asymptomatic dogs, no significant difference was found in the mean OD values obtained for the detection of IgG (1.778 ± 0.387 and $1.622 \pm 0.466, p = 0.20$, respectively) (Fig. 1 and 2), IgG1 (0.252 ± 0.287 and $0.212 \pm 0.166, p = 0.65$) or IgG2 (1.244 ± 0.281 and $1.037 \pm 0.410, p = 0.08$).

The seroprevalence of IgG1 did not differ significantly between symptomatic and asymptomatic dogs when *L. (V.) braziliensis* (84% versus 75%, $p = 0.84$) or *L. (L.) chagasi* antigen (88% versus 83.3%, $p = 0.90$) was used.

In the CI group, mean OD values and their standard deviations for the detection of IgG, IgG1 and IgG2 were, respectively: 0.886 ± 0.342 , 0.134 ± 0.048 and 0.897 ± 0.134 when *L. (V.) braziliensis* antigen was used, and 0.957 ± 0.423 , 0.137 ± 0.076 and 0.418 ± 0.204 when *L. (L.) chagasi* antigen was used. No significant difference in mean OD values for the detection of IgG was observed between antigens ($p = 0.790$) (Fig. 1 and 2).




Fig. 3 - Distribution of OD values in ELISA reactions obtained for the visceral leishmaniasis group (VL) (symptomatic subgroup, VLs, and asymptomatic subgroup, VLs), group co-infected with *L. (L.) chagasi* and *L. (V.) braziliensis* (CI), and control group (healthy subgroup, HC, and ehrlichiosis subgroup, EC): **A**- IgG1; **B**- IgG2 using *L. (V.) braziliensis* antigen and **C**- IgG1 and **D**- IgG2 using *L. (L.) chagasi* antigen.

In the HC and EC control groups, low mean OD values and standard deviations were obtained for the detection of IgG (Fig. 1 and 2), IgG1 and IgG2 when *L. (V.) braziliensis* (0.126 ± 0.031 , 0.076 ± 0.023 , and 0.158 ± 0.088 , respectively) and *L. (L.) chagasi* antigens (0.111 ± 0.076 , 0.073 ± 0.039 , and 0.069 ± 0.024) were used. In the EC group, IgG1 was detected in 29.0% and 14.3% of serum samples tested with *L. (V.) braziliensis* and *L. (L.) chagasi* antigen, respectively.

The distribution of OD values in ELISA reactions obtained for IgG1 and IgG2 for all groups are shown in Fig. 3.

DISCUSSION

In the present study, we evaluated the detection of IgG and its subclasses by ELISA comparing *L. (L.) chagasi* (homologous) and *L. (V.) braziliensis* (heterologous) antigens in order to improve the performance of the assay for the diagnosis of AVL in symptomatic or asymptomatic naturally infected dogs.

Higher mean OD values were obtained for the detection of IgG, IgG1 and IgG2 in sera from dogs with AVL when *L. (L.) chagasi* antigen was employed. The use of homologous antigen for the diagnosis of leishmaniasis has been reported to provide the best results for both canine⁴ and human^{2,21} AVL, as well for canine⁴⁶ and human⁶ American

cutaneous leishmaniasis³. BADARÓ *et al.*³ demonstrated that the use of antigens from *Leishmania* sp. isolated from the same area where the cases occurred increased the accuracy of the assay. DA COSTA *et al.*¹² observed no significant differences in the sensitivity of IIF reactions for the diagnosis of canine AVL when different *Leishmania* species were used as antigens.

Overlapping transmission of AVL caused by *L. (L.) chagasi* and American cutaneous leishmaniasis caused by *L. (V.) braziliensis* has been observed in various regions of the Municipality of Rio de Janeiro^{39,40}. In the group of dogs co-infected with *L. (V.) braziliensis* and *L. (L.) chagasi*, similar mean OD values for the detection of IgG were obtained with the two antigens, a finding suggesting a balance in the humoral immune response. Since euthanasia is not indicated for dogs infected with *L. (V.) braziliensis* from areas with overlapping transmission of the two species, ELISA employing both antigens could be used to select dogs for parasitological investigation⁴⁶.

Some investigators have reported an association between the immune response (Th1/Th2) and IgG subclasses in dogs infected with *L. (L.) infantum*^{7,15}. However, other authors do not agree that the IgG2 or IgG1 subclass is a marker of a Th1 or Th2 immune response in dogs, respectively^{14,33}. This discrepancy might be explained by the different antisera used.

AVL is frequently associated with a marked specific humoral response and anti-*Leishmania* IgG subclass levels have been suggested to be markers of disease susceptibility and resistance, with IgG1 being associated with the development of the disease and IgG2 with asymptomatic infection^{7,9,15,25,43,49}. However, this association was not observed by QUINELL *et al.* (2003) when using monoclonal antisera.

In the present study, mean OD values obtained for the IgG, IgG1 and IgG2 reactions were higher in symptomatic (group VLs) than in asymptomatic dogs (group VL_a), a finding also reported by INIESTA *et al.*²². IgG2 was the most seroprevalent subclass in both subgroups, with a seroprevalence close to that of IgG, a fact that did not increase the performance of the diagnostic test. These results agree with similar studies involving dogs infected with *L. infantum*^{7,10,20,38,53}.

The seroprevalence of IgG1 was relatively low in infected animals but slightly higher in symptomatic (group VLs) dogs when compared to asymptomatic animals (group VL_a). This low seroprevalence of IgG1 indicates its limited use for the diagnosis of AVL. The limitation of IgG1 in the diagnosis of canine AVL has been suggested previously³⁸. Other investigators proposed that this subclass predominates in dogs infected with *L. (L.) chagasi* or *L. (L.) infantum*^{37,44}, or that IgG1 levels are only elevated in the presence of signs of visceral leishmaniasis⁴⁹.

In the present study, the correlation between OD values for the detection of IgG and IgG2 was high and significant. However, the correlation of OD values for IgG or IgG2 with IgG1 was low. LEANDRO *et al.*²⁵, studying symptomatic infected dogs, also observed high levels of IgG2, with a strong correlation between IgG and IgG2. In agreement with these findings, there was no significant difference between the AUCs of IgG and IgG2, whereas a significant difference was observed when comparing each marker to the AUC of IgG1.

Canine ehrlichiosis is an important differential diagnosis of AVL due to the similar clinical findings and high prevalence of dogs infected with *E. canis* in Rio de Janeiro²⁷. Serology has been used to differentiate the two infections⁴⁸. LLERA *et al.*²⁶ observed no cross-reaction between sera of dogs infected with *E. canis* and *L. infantum*. However, we observed cross-reactivity in IIF and in ELISA (IgG1). Similar findings have been reported by FERREIRA *et al.*¹⁸.

In conclusion, the present study suggests that IgG ELISA using *L. (L.) chagasi* antigen yielded the best results for the diagnosis of AVL, permitting the discrimination between cases of AVL and ehrlichiosis in dogs. However, the 100% positivity of IgG observed for serum samples from dogs of the VL group might be related to the selection criterion adopted in this study, i.e., dogs testing seropositive by IIF. We suggest a comparative study of IIF and IgG ELISA using *L. (L.) chagasi* antigen for the evaluation of dogs from AVL-endemic areas under field conditions.

RESUMO

A utilização do ELISA empregando antígenos homólogos e heterólogos para a detecção de IgG e subclasses (IgG1 e IgG2) no diagnóstico de Leishmaniose Visceral Canina

A imunofluorescência indireta é o método recomendado para o diagnóstico de leishmaniose visceral em cães, entretanto, a acurácia

dessa técnica é baixa e seu uso em grande escala é limitado. Uma vez que o ELISA não apresenta essas limitações, essa técnica poderia ser uma opção para a detecção de IgG ou subclasses IgG1 e IgG2 específicas. A ehrlichiose canina é um importante diagnóstico diferencial de Leishmaniose Visceral Americana (LVA). O presente estudo comparou o ELISA usando antígenos de *Leishmania chagasi* e *Leishmania braziliensis* para a detecção de IgG e subclasses anti-*Leishmania* em amostras de soro de 37 cães naturalmente infectados com *L. chagasi* (LVA) e em amostras de quatro cães co-infectados (CI). A ocorrência de reatividade cruzada foi investigada em amostras de soro controle de 17 animais saudáveis (HC) e 35 de infectados por *Ehrlichia canis* (EC). A média de densidade óptica obtida para a detecção de IgG foi significativamente maior quando o antígeno de *L. chagasi* foi usado e também mais elevada no subgrupo LVs (sintomático) quando comparado ao subgrupo LV_a (assintomático). A correlação entre IgG e IgG1 foi baixa. O presente resultado sugere que ELISA IgG empregando antígeno homólogo, produz os melhores resultados, permitindo o diagnóstico de infecção assintomática por *L. chagasi* e a discriminação entre casos de LVA e ehrlichiose em cães.

ACKNOWLEDGMENTS

We thank the Instituto Municipal de Medicina Veterinária Jorge Vaitsman and the veterinarian Bianca Pellegrinetti Junqueira de Andrade for help with the collection of the control sera, and Centro de Controle de Zoonoses Paulo Darcoso Filho and Programa de Controle das Leishmanioses da Secretaria Municipal de Saúde do Rio de Janeiro for help with the identification of the animals studied. The study was conducted according to ethical guidelines and was approved by the Ethics Committee for the Use of Animals (CEUA/FIOCRUZ/P-0276-05).

This study was partially supported by Programa de Apoio a Pesquisa Estratégica em Saúde (PAPES IV). A. Schubach and T. Schubach are the recipients of fellowships from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

REFERENCES

1. Alvar J, Cañavate C, Molina R, Moreno J, Nieto J. Canine leishmaniasis. *Adv Parasitol.* 2004;57:1-88.
2. Badaró R, Reed SG, Carvalho EM. Immunofluorescent antibody test in American visceral leishmaniasis: sensitivity and specificity of different morphological forms of two *Leishmania* species. *Am J Trop Med Hyg.* 1983;32:480-4.
3. Badaró R, Reed SG, Barral A, Orge G, Jones TC. Evaluation of the micro enzyme-linked immunosorbent assay (ELISA) for antibodies in American visceral leishmaniasis: antigen selection for detection of infection-specific responses. *Am J Trop Med Hyg.* 1986;35:72-8.
4. Baleeiro CO, Paranhos-Silva M, dos Santos JC, Oliveira GG, Nascimento EG, de Carvalho LP, *et al.* Montenegro's skin reactions and antibodies against different *Leishmania* species in dogs from a visceral leishmaniasis endemic area. *Vet Parasitol.* 2006;139:21-8.
5. Barbosa-De-Deus R, Dos Mares-Guia ML, Nunes AZ, Costa KM, Junqueira RG, Mayrink W, *et al.* *Leishmania major*-like antigen for specific and sensitive serodiagnosis of human and canine visceral leishmaniasis. *Clin Diagn Lab Immunol.* 2002;9:1361-6.
6. Barroso-Freitas A, Passos SR, Mouta-Confort E, Madeira MF, Schubach AO, Santos GP, *et al.* Accuracy of an ELISA and indirect immunofluorescence for the laboratory diagnosis of American tegumentary leishmaniasis. *Trans R Soc Trop Med Hyg.* 2009;103:383-9.

7. Bourdoiseau G, Bonnefont C, Hoareau E, Boehringer C, Stolle T, Chabanne L. Specific IgG1 and IgG2 antibody and lymphocyte subset levels in naturally *Leishmania infantum*-infected treated and untreated dogs. *Vet Immunol Immunopathol*. 1997;59:21-30.
8. Braga MD, Coelho IC, Pompeu MM, Evans TG, MacAulliffe IT, Teixeira MJ, *et al*. Controle do calazar canino: comparação dos resultados de um programa de eliminação rápida de cães sororreagentes por ensaio imuno-enzimático com outro de eliminação tardia de cães sororreagentes por teste de imunofluorescência indireta de eluato de papel filtro. *Rev Soc Bras Med Trop*. 1998;31:419-24.
9. Cardoso L, Schallig HD, Cordeiro-da-Silva A, Cabral M, Alunda JM, Rodrigues M. Anti-*Leishmania* humoral and cellular immune responses in naturally infected symptomatic and asymptomatic dogs. *Vet Immunol Immunopathol*. 2007;117:35-41.
10. Cordeiro-da-Silva A, Cardoso L, Araújo N, Castro H, Tomás A, Rodrigues M, *et al*. Identification of antibodies to *Leishmania* silent information regulatory 2 (SIR2) protein homologue during canine natural infections: pathological implications. *Immunol Lett*. 2003;86:155-62.
11. Courtenay O, Quinnell RJ, Garcez LM, Shaw JJ, Dye C. Infectiousness in a cohort of Brazilian dogs: why culling fails to control visceral leishmaniasis in areas of high transmission. *J Infect Dis*. 2002;186:1314-20.
12. Da Costa CA, Genaro O, de Lana M, Magalhães PA, Dias M, Michalick MS, *et al*. Leishmaniose visceral canina: avaliação da metodologia sorológica utilizada em inquéritos epidemiológicos. *Rev Soc Bras Med Trop*. 1991;24:21-5.
13. Da Costa-Val AP, Cavalcanti RR, de Figueiredo Gontijo N, Michalick MS, Alexander B, Williams P, *et al*. Canine visceral leishmaniasis: relationships between clinical status, humoral immune response, haematology and *Lutzomyia (Lutzomyia) longipalpis* infectivity. *Vet J*. 2007;174:636-43.
14. Day MJ. Immunoglobulin G subclass distribution in canine leishmaniasis: a review and analysis of pitfalls in interpretation. *Vet Parasitol*. 2007;147: 2-8.
15. Deplazes P, Smith NC, Arnold P, Lutz H, Eckert J. Specific IgG1 and IgG2 antibody responses of dogs to *Leishmania infantum* and other parasites. *Parasite Immunol*. 1995;17:451-8.
16. Dye C. The logic of visceral leishmaniasis control. *Am J Trop Med Hyg*. 1996;55:125-30.
17. Edrissian GH, Darabian P. A comparison of enzyme-linked immunosorbent assay and indirect fluorescent antibody test in the serodiagnosis of cutaneous and visceral leishmaniasis in Iran. *Trans R Soc Trop Med Hyg*. 1979;73:289-92.
18. Ferreira EC, De Lana M., Carneiro M, Reis AB, Paes DV, Da Silva ES, *et al*. Comparison of serological assays for the diagnosis of canine visceral leishmaniasis in animals presenting different clinical manifestations. *Vet. Parasitol*. 2007;146:235-41.
19. Fisa R, Gállego M, Castillejo S, Aisa MJ, Serra T, Riera C, *et al*. Epidemiology of canine leishmaniasis in Catalonia (Spain) the example of the Priorat focus. *Vet Parasitol*. 1999;83:87-97.
20. Harrus S, Waner T, Strauss-Ayali D, Bark H, Jongejan F, Hecht G, *et al*. Dynamics of IgG1 and IgG2 subclass response in dogs naturally and experimentally infected with *Ehrlichia canis*. *Vet Parasitol*. 2001;99:63-71.
21. Hommel M, Peters W, Ranque J, Quilici M, Lannotte G. The micro-ELISA technique in the serodiagnosis of visceral leishmaniasis. *Ann Trop Med Parasitol*. 1978;72:213-18.
22. Iniesta L, Gállego M, Portús M. Immunoglobulin G and E responses in various stages of canine leishmaniasis. *Vet Immunol Immunopathol*. 2005;103:77-81.
23. Jones TC, Hunt RD, King NW. Moléstias causadas por protozoários In: Jones TC, Hunt RD, King NW. *Patologia veterinária*. 6ª ed. São Paulo: Manole; 2000. p. 559-610.
24. Junqueira Pedras M, Orsini M, Castro M, Passos VM, Rabello A. Antibody subclass profile against *Leishmania braziliensis* and *Leishmania amazonensis* in the diagnosis and follow-up of mucosal leishmaniasis. *Diagn Microbiol Infect Dis*. 2003;47:477-85.
25. Leandro C, Santos-Gomes GM, Campino L, Romão P, Cortes S, Rolão N, *et al*. Cell mediated immunity and specific IgG1 and IgG2 antibody response in natural and experimental canine leishmaniasis. *Vet Immunol Immunopathol*. 2001;79:273-84.
26. Llera JL, López García ML, Martín Reinoso E, De Vivar González R. Differential serological testing by simultaneous indirect immunofluorescent antibody test in canine leishmaniasis and ehrlichiosis. *Vet Parasitol*. 2002;109:185-90.
27. Macieira DB, Messick JB, Cerqueira AD, Freire IM, Linhares GF, Almeida NK, *et al*. Prevalence of *Ehrlichia canis* infection in thrombocytopenic dogs from Rio de Janeiro, Brazil. *Vet Clin Pathol*. 2005;34:44-48.
28. Madeira M F, Schubach AO, Schubach TMP, Leal CA, Marzochi CA. Identification of *Leishmania (Leishmania) chagasi* isolated from health skin of symptomatic and asymptomatic dogs seropositive for Leishmaniasis in the Municipality of Rio de Janeiro, Brazil. *Braz J Infect Dis*. 2004;8:440-4.
29. Madeira MF, Schubach AO, Schubach TM, Pereira SA, Figueiredo FB, Baptista C, *et al*. Post mortem parasitological evaluation of dogs seroreactive for *Leishmania* from Rio de Janeiro, Brazil. *Vet Parasitol*. 2006;138:366-70.
30. Mancianti F, Falcone ML, Gianneli C, Poli A. Comparison between an enzyme-linked immunosorbent assay using a detergent-soluble *Leishmania infantum* antigen and indirect immunofluorescence for the diagnosis of canine leishmaniasis. *Vet. Parasitol*. 1995;59:13-21.
31. Manzillo VF, Cappiello S, Oliva G. Tick-transmitted diseases in dogs: clinic pathological findings. *Parassitologia*. 2006;48:135-6.
32. Marzochi MC, Sabroza PC, Toledo LM, Marzochi KB, Tramontano NC, Rangel Filho FB. Leishmaniose visceral na cidade do Rio de Janeiro-Brasil. *Cad Saúde Públ*. 1985;1: 5-17.
33. Mazza G, Duffus WP, Elson CJ, Stokes CR, Wilson AD, Whiting AH. The separation and identification by monoclonal antibodies of dog IgG fractions. *J Immunol Methods*. 1993;161:193-203.
34. Mazza G, Whiting AH, Day MJ, Duffus WPH. Development of an enzyme-linked immunosorbent assay for the detection of IgG subclasses in the serum of normal and diseased dogs. *Res Vet Sci*. 1994;57:133-9.
35. Mazza G, Whiting AH, Day MJ, Duffus WPH. Preparation of monoclonal antibodies specific for the subclasses of canine IgG. *Res Vet Sci*. 1994;57:140-5.
36. McDade JE. Ehrlichiosis - a disease of animals and humans. *J Infect Dis*. 1990;161:609-17.
37. Mendes CO, De Sousa EP, Borja-Cabrera GP, Batista LM, Santos MA, Parra LE, *et al*. IgG1/IgG2 antibody dichotomy in sera of vaccinated or naturally infected dogs with visceral leishmaniasis. *Vaccine*. 2003;21:2589-97.
38. Mettler M, Grimm F, Capelli G, Camp H, Deplazes P. Evaluation of Enzyme-Linked Immunosorbent Assays, an Immunofluorescent-antibody test, and two rapid tests (Immunochromatographic-Dipstick and Gel Tests) for serological diagnosis of symptomatic and asymptomatic *Leishmania* infections in dogs. *J Clin Microbiol*. 2005;43:5515-9.
39. Ministério da Saúde. Manual de vigilância e controle da leishmaniose visceral. Brasília: Ministério da Saúde, 2006. 120 p.
40. Ministério da Saúde. Manual de vigilância e controle da leishmaniose tegumentar americana. Brasília: Ministério da Saúde; 2007. 182 p.
41. Molina R, Amela C, Nieto J, San-Andrés M, González F, Castillo JA, *et al*. Infectivity of dogs naturally infected with *Leishmania infantum* to colonized *Phlebotomus perniciosus*. *Trans R Soc Trop Med Hyg*. 1994;88:491-3.

42. Moreira SM, Bastos CV, Araújo RB, Santos M, Passos LMF. Retrospective study (1998-2001) on canine ehrlichiosis in Belo Horizonte, MG, Brazil. *Arq Bras Med Vet Zootec.* 2003;55:141-7.
43. Nakhaee A, Taheri T, Taghikhani M, Mohebbali M, Salmanian AH, Fasel N, *et al.* Humoral and cellular immune responses against Type I cysteine proteinase of *Leishmania infantum* are higher in asymptomatic than symptomatic dogs selected from a naturally infected population. *Vet Parasitol.* 2004;119:107-23.
44. Quinnell RJ, Courtenay O, Garcez LM, Kaye PM, Shaw MA, Dye C, *et al.* IgG subclass responses in a longitudinal study of canine visceral leishmaniasis. *Vet Immunol Immunopathol.* 2003;91:161-8.
45. Reis AB, Teixeira-Carvalho A, Vale AM, Marques MJ, Giunchetti RC, Mayrink W, *et al.* Isotype patterns of immunoglobulins: hallmarks for clinical status and tissue parasite density in Brazilian dogs naturally infected by *Leishmania (Leishmania) chagasi*. *Vet Immunol Immunopathol.* 2006;112:102-16.
46. Ribeiro FC, Schubach AO, Mouta-Confort E, Schubach TM, Madeira MF, Marzochi MC. Use of ELISA employing *Leishmania (Viannia) braziliensis* and *Leishmania (Leishmania) chagasi* antigens for the detection of IgG and IgG1 and IgG2 subclasses in the diagnosis of American tegumentary leishmaniasis in dogs. *Vet Parasitol.* 2007;148:200-6.
47. Rikihisia Y. The tribe *Ehrlichiae* and ehrlichial diseases. *Clin Microbiol Rev.* 1991;4:286-308.
48. Sanchez Visconti G, Tesouro Diez MA, editors. Ehrlichiosis. *Canis et felis* (San Román, Madrid). 2001; 51:49-56.
49. Solano-Gallego L, Riera C, Roura X, Iniesta L, Gallego M, Valladares JE, *et al.* *Leishmania infantum*-specific IgG, IgG1 and IgG2 antibody responses in healthy and ill dogs from endemic areas. Evolution in the course of infection and after treatment. *Vet Parasitol.* 2001;96:265-76.
50. Souza MA, Da Silva AG, Afonso-Cardoso SR, Favoreto Júnior SF, Ferreira MS. Perfil de isotipos de imunoglobulinas e subclasses de IgG na leishmaniose tegumentar americana. *Rev Soc Bras Med Trop.* 2005;38:137-41.
51. Tesh R. Control of zoonotic visceral leishmaniasis: is it time to change strategies? *Am J Trop Med Hyg.* 1995;52:287-92.
52. Tizard IR. Anticorpos forma solúvel de RBC. In: Tizard IR. *Imunologia veterinária, uma introdução.* 5ª ed. São Paulo: Roca; 1998. p. 160-171.
53. Vercammen F, Fernandez-Perez FJ, Amo CD, Alunda JM. Follow-up of *Leishmania infantum* naturally infected dogs treated with allopurinol: immunofluorescence antibody test, ELISA and Western blot. *Acta Trop.* 2002;84:175-81.
54. Voller A, Bartlett A, Bidwell DE, Clark MF, Adams AN. The detection of viruses by enzyme-linked immunosorbent assay (ELISA). *J Gen Virol.* 1976;33:165-7.
55. Woody BJ, Hoskins JD. Ehrlichial diseases of dogs. *Vet Clin North Am Small Anim Pract.* 1991;21:75-98.

Received: 24 August 2009

Accepted: 5 August 2011

Revista do Instituto de Medicina Tropical de São Paulo *on line.*

Publications from 1990 to the present data are now available on:

<http://www.scielo.br/rimtsp>
PAST ISSUES 1959-1989 (PDF)
www.imt.usp.br/portal/

Revista do Instituto de Medicina Tropical de São Paulo - Home page - Windows Internet Explorer

http://www.scielo.br/rimtsp?script=sci_serial&pid=0036-4665&lng=en&nrm=iso

Arquivo Editar Exibir Favoritos Ferramentas Ajuda

issues articles search

all previous current next author subject form alpha

português
español

Update on
July 02, 20

▶ about the journal
▶ editorial board
▶ instructions to authors
▶ subscription
▶ statistics

Indicadores SCImago

Indicator	2000-2007	Value
SJR		0.1
Cites per doc		0.8
Total cites		205

www.sciimagojr.com

REVISTA
DO
INSTITUTO
DE
MEDICINA
TROPICAL
DE
SÃO PAULO

JOURNAL OF THE SÃO PAULO
INSTITUTE OF TROPICAL MEDICINE

Print ISSN 0036-4665

Publication of
Instituto de Medicina Tropical

Mission
To publish articles dealing with research on tropical diseases and related sciences

© 2008 Instituto de Medicina Tropical de São Paulo

Av. Dr. Enéas de Carvalho Aguiar, 470
05403-000 São Paulo SP - Brazil
Tel. / Fax: +55 11 3062-2174

revimtsp@edu.usp.br

SciELO – The Scientific Electronic Library OnLine - SciELO is an electronic virtual covering a selected collection of Brazilian scientific journals.

The library is an integral part of a project being developed by FAPESP – Fundação de Amparo à Pesquisa do Estado de São Paulo, in partnership with BIREME – the Latin American and Caribbean Center on Health Sciences Information.

SciELO interface provides access to its serials collection via an alphabetic list of titles or a subject index or a search by word of serial titles, publisher names, city of publication and subject.

The interface also provides access to the full text of articles via author index or subject index or a search form on article elements such as author names, words from title, subject and words from full text.

FAPESP/BIREME Project on Scientific Electronic Publications
Latin American and Caribbean Center on Health Sciences Information
Rua Botucatu 862 – 04023-901 São Paulo, SP – Brazil
Tel. (011) 5576-9863
scielo@bireme.br