

## Soil management of sugarcane fields affecting CO<sub>2</sub> fluxes

Rose Luiza Moraes Tavares<sup>1\*</sup>, Diego Silva Siqueira<sup>2</sup>, Alan Rodrigo Panosso<sup>3</sup>, Guilherme Adalberto Ferreira Castioni<sup>1</sup>, Zigomar Menezes de Souza<sup>1</sup>, Newton La Scala Jr.<sup>2</sup>

<sup>1</sup>University of Campinas/FEAGRI – Dept. of Soils, Av. Cândido Rondon, 501 – 13083-875 – Campinas, SP – Brazil.

<sup>2</sup>São Paulo State University/FCAV, Via de Acesso Prof. Paulo Donato Castellane, s/n – 14884-900 – Jaboticabal, SP – Brazil.

<sup>3</sup>São Paulo State University/FEIS, Al. Rio de Janeiro, 266 – 15385-000 – Ilha Solteira, SP – Brazil.

\*Corresponding author <[roseluziamt@gmail.com](mailto:roseluziamt@gmail.com)>

Edited by: Carlos Eduardo Pellegrino Cerri

Received April 23, 2015

Accepted December 21, 2015

**ABSTRACT:** The harvesting system of green sugarcane, characterized by mechanized harvesting and no crop burning, affects soil quality by increasing the remaining straw left on the soil surface after harvesting, thus, contributing to the improvement of physical, chemical, and microbiological soil attributes, influencing CO<sub>2</sub> fluxes. This study aimed to evaluate CO<sub>2</sub> fluxes and their relation to soil properties in sugarcane crops under different harvesting managements: burned (B), Green harvesting for 5 years (G-5) and Green harvesting for ten years (G-10). For this, a 1 ha sampling grid with 30 points was installed in each area, all located in the Northeast of São Paulo State, Brazil. In each point, CO<sub>2</sub> fluxes were measured and the soil was sampled to analyze the microbial biomass, physical (soil moisture and temperature, mean weight diameter, bulk density, clay, macroporosity and microporosity) and chemical characterization (pH, organic C, base saturation and P). The CO<sub>2</sub> fluxes were divided into four quantitative criteria: *high*, *moderate*, *low* and *very low* from the Statistical Division (mean, first quartile, median and third quartile) and the other data were classified according this criterion. The Principal Component Analysis (PCA) was used to identify the main soil attributes that influence CO<sub>2</sub> fluxes. The results showed that G-10 CO<sub>2</sub> fluxes were 28 and 41 % higher than those in the G-5 and B treatments, respectively. The PCA analysis showed that macroporosity was the main soil attribute that influenced the *high* CO<sub>2</sub> fluxes.

**Keywords:** *Sacharum officinarum*, principal component analysis, porosity, biomass

### Introduction

The practice of burning sugarcane residues prior to harvesting aims to facilitate manual cutting, but the temperature during sugarcane burning is around 160-200 °C on the soil surface, causing nutrients loss by volatilization such as phosphorus, sulfur and nitrogen (Ball et al., 1993) and may lead to a great decline in soil C stocks (Song et al., 2013). The "harvesting of green sugarcane" is a system without burning that leaves biomass waste in the field after harvesting, positively influencing soil quality by increasing the deposited residual straw (mean 10 to 30 Mg ha<sup>-1</sup>) allowing carbon accumulation in the soil, which implies in a positive CO<sub>2</sub> balance as described by Razafimbelo et al. (2006).

Soil CO<sub>2</sub> fluxes from areas of sugarcane cultivation were studied by Brito et al. (2009) that found greater fluxes in areas with greater soil macroporosity. Panosso et al. (2009) compared the soil CO<sub>2</sub> in pre-harvesting burned crop with a green harvesting system and found that soil cations were the main soil attribute to explain the CO<sub>2</sub> fluxes mainly in the burned area.

Soil CO<sub>2</sub> fluxes result from physical and biological processes that affect CO<sub>2</sub> production and transport from the soil to the atmosphere. In addition, production is related to root respiration and the action of microorganisms during OM decomposition (Jenkinson and Ladd, 1981; Brito et al., 2009). Transport of soil gases is influenced by the physical structure parameters, such as porosity, which drive the gas flow. Saturation of soil pores also determines CO<sub>2</sub> fluxes. According to the literature, the main soil attributes that influence CO<sub>2</sub> fluxes

include temperature and content of soil water (Xu and Qi, 2001; Epron et al., 2004, 2006; Kosugi et al., 2007; La Scala et al., 2010; Leon et al., 2014), attributes with great influence on microbial activities that promote soil respiration.

The principal component analysis (PCA) of CO<sub>2</sub> fluxes (Panosso et al., 2012) showed that water filled pore space, and total porosity and macroporosity were the main components to explain the variance of CO<sub>2</sub> fluxes. Another study, about soil CO<sub>2</sub> efflux in a water limited ecosystem (Leon et al., 2014), showed that the most important attributes were root biomass, soil volumetric water content and total porosity.

Mitigating CO<sub>2</sub> fluxes in sugarcane cultivation still requires further studies aiming to assess their viability and enhancing their applicability for environmental purposes. More specifically, there is a need to study the main factors responsible for high soil CO<sub>2</sub> fluxes, which can assist in the challenge of achieving stability of soil carbon through improved decision-making managements. This study aimed to evaluate CO<sub>2</sub> fluxes and their relation to soil properties in sugarcane areas under different harvesting managements.

### Materials and Methods

#### Experimental site

This study was conducted on a farm with more than 30 years of sugarcane (*Saccharum* spp.) cultivation history. The land belongs to a sugar and ethanol mill located in the Pradópolis, São Paulo State, Brazil (21°19'8" S, 48°7'24" W), approximately 500 m above sea level

(Figure 1). The soil was classified as Haplustox (USDA Soil Taxonomy) (Latossolo Vermelho Eutroférrico, according to Brazilian Soil Classification System) with a clayey texture (561 g kg<sup>-1</sup> to B, 517 g kg<sup>-1</sup> to G-5 and 531 g kg<sup>-1</sup> to G-10), and the topography of the area is flat and undulating. The regional climate is classified as B<sub>2</sub>rB'4a' by the Thornthwaite system (Rolim et al., 2007), indicating a mesothermal region with rainy summers and dry winters. The average precipitation is 1425 mm yr<sup>-1</sup> and is concentrated between Oct and Mar. The average annual temperature over the last 30 years was 22.2 °C.

In 2011, three plots were chosen in areas with different systems and management times (Figure 1): the burned sugarcane (B) area has been managed under residue burning since the 1980s and the other areas were harvested under the green sugarcane system (G) with different starting times of green sugarcane adoption [5 years (G-5) that started in 2006 and another area with ten years (G-10) that started in 2001]. At the time of renewing the plantations, which occurred in every six ratoons in B and in G-10 in 2007) with mechanical removal of the ratoon of the previous crop and subsoiling to 0.45 m deep in the planting furrows. Afterward, 2 t ha<sup>-1</sup> of dolomitic limestone and 480 kg ha<sup>-1</sup>

of NPK fertilizer at 10-25-20 formulation were also applied. On average, 100 m<sup>3</sup> ha<sup>-1</sup> of vinasse (by-product of biomass distillation of the sugarcane fuel industry) and 300 kg ha<sup>-1</sup> of urea or 200 kg ha<sup>-1</sup> of ammonium nitrate were applied to the areas after 5-7 months of the first fertilization.

In 2011-2012 (experimental evaluation period), treatment B with the sugarcane variety CTC4 (average maturity and high agricultural productivity) was in its 5<sup>th</sup> ratoon with average yield of 67 t ha<sup>-1</sup>. Treatment G-5 with the planted variety RB85 5453 (early maturity, erect growth, high productivity with no limitation of soil water) was in its 4<sup>th</sup> ratoon with average yield of 80 t ha<sup>-1</sup>. Treatment G-10 with the sugarcane variety CTC 20 (early maturity, high tillering and high productivity along the cuts) was in its 5<sup>th</sup> ratoon with an average yield of 75 t ha<sup>-1</sup>. In this experiment, no manure or fertilizers were applied between the years 2010 and 2011 (before and during the field experiment) to control interferences of these factors on the CO<sub>2</sub> fluxes. In each area, a sampling irregular grid of 1 ha was installed with 30 sampling points spaced at intervals with minimum of 2 m and maximum of 100 m. All points were georeferenced with the aid of a total station and a DGPS (Model L1/L2 Hiper Lite Plus).




Figure 1 – Description of experimental locations and relief maps with sampling grid details. B = burned sugarcane; G-5 = green sugarcane with 5 years of implementation; G-10 = green sugarcane with 10 years of implementation.

### Measurement of CO<sub>2</sub> fluxes

Measurements of soil CO<sub>2</sub> fluxes were simultaneously performed in the three sugarcane areas in the same period (10 d in Aug-Sept 2011) and on the same day time (7:00-11:00 a.m.) after harvesting, for standardization. For that, three portable systems (1 system/area) were used to monitor the changes in CO<sub>2</sub> concentration inside the chamber using an infrared gas analyzer. The soil chamber has an internal volume of 854.2 cm<sup>3</sup> with a circular soil contact area at the base of 83.7 cm<sup>2</sup>, which was placed on PVC collars previously inserted at each sampling point to 3 cm deep keeping is distant from the ratoon plant (approximately 30 cm) to decrease its influence on the CO<sub>2</sub> fluxes. Once the chamber is set to the measurement mode, it takes around 1.5 min to run the time-change interpolation of CO<sub>2</sub> concentration inside the chamber. The chambers were previously calibrated for this work.

Soil temperature (Ts) and soil water content (Ms) were measured simultaneously with CO<sub>2</sub> concentration through a temperature sensor coupled with the system, and Ms was registered with a portable Hydrosense system.

### Soil sampling and evaluation

For the microbial biomass (Biom) analysis, soil samples were collected at each point in the grid on the same day and time of CO<sub>2</sub> measurement, but only for 2 d of each collection period (the first and last day of CO<sub>2</sub> measurement) due to the large number of samples to be analyzed in 30 d (recommendation for the microbiological analysis). In the field, samples were kept in plastic bags inside Styrofoam boxes and transferred immediately to a refrigerator at 4 °C. The biomass analysis was performed according to the fumigation-extraction method proposed by Jenkinson and Powlson (1976).

For the other soil analysis, the samples were collected once at each point before the CO<sub>2</sub> analysis. Disturbed soil samples were collected from the first 20 cm of soil to evaluate organic carbon (C) (Nelson and Sommers, 1982), pH in CaCl<sub>2</sub> and phosphorus (P) by resin procedure (Raij et al., 2001), clay and mean weight diameter of soil aggregates (MWD).

Samples were exposed to air for 24 h, kept moist for aggregate preservation and then placed on a sieve set of 6.35 and 2 mm mesh diameter. Aggregates were obtained from samples retained by the 2 mm mesh, whereas those that passed through were again exposed to air until constant weight was achieved. Undisturbed soil

samples were collected with aluminum rings and used to analyze macroporosity (Ma), microporosity (Mi) and bulk density (Bd). These physical analyses were carried out according to Brazilian Agricultural Research Corporation methodologies – Embrapa (1997).

### Statistical analyses

Mean daily CO<sub>2</sub> fluxes were evaluated by the *t* test for comparison between management areas (*p* < 0.05), using the program Minitab 14. These values were integrated to calculate the CO<sub>2</sub> accumulated during 10 d.

Quantitative criteria of CO<sub>2</sub> fluxes were defined by the distribution of CO<sub>2</sub> data in each area (Table 1), defined as: *very low* (VL) fluxes, which included values lower than the first quartile (Q1); *low* fluxes (L), between Q1 and median values; *moderate* (M) fluxes, between median and third quartile (Q3); and *high* (H) fluxes, values greater than Q3.

This criterion was used to identify the influence of soil attributes on different CO<sub>2</sub> concentrations, mainly by high fluxes and if the amount of soil attributes followed the same trend of CO<sub>2</sub> criterion. When the values were different for the three sugarcane systems, discussion was made separately for each area.

The multivariate structure in the original data set was evaluated by the PCA that condensed the relevant information into a smaller set of orthogonal latent variables called principal components (PC-eigenvectors). Each pair of principal components (PCs) generates a two-dimensional representation of the original sample space, known as a biplot. The biplot explains the structure of variables directing beams of variable regions of maximum variability. In this work, we considered the principal components whose eigenvalues were greater than a unity (Kaiser, 1958). The sign and relative size of the linear function coefficients, which define the PC scores were used as an indication of the weight to be assigned to each variable in the different experimental plots (Johnson and Wichern, 2002). The correlation between soil attributes with PCs to explain the management types were compared with the mean values of soil CO<sub>2</sub> fluxes.

## Results and Discussion

### CO<sub>2</sub> total fluxes

Considering the CO<sub>2</sub> total fluxes, G-10 showed higher values (mean 2.71 μmol CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>), representing a significant difference (*p* < 0.05) of 28 % compared

Table 1 – Summary of soil CO<sub>2</sub> flux (μmol m<sup>-2</sup> s<sup>-1</sup>) distribution statistics of burned and green cane areas.

Area	Mean	Variance	Minimum	Q1	Median	Q3	Maximum	K-S
B	1.58 b	0.21	0.54	1.37	1.62	1.85	2.47	0.10
G-5	1.93 b	0.34	1.06	1.56	1.78	2.34	3.32	0.12
G-10	2.71 a	0.91	1.12	2.12	2.39	3.21	4.76	0.14

B = Burned cane; G-5 = Green harvest sugarcane for 5 years; G-10 = Green harvesting sugarcane for 10 years; Q1 = first quartile; Q3 = third quartile; K-S: Kolmogorov-Smirnov test. Means followed by the same letter in column do not differ from each other by the *t* test at 5 % probability level.

with G-5 (1.93 μmol CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>) and 41 % compared with B (1.58 μmol CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>) (Table 1). The G-5 and B did show statistical difference, the G-5 area is in a transition stage, considering the recent conversion to the green sugarcane system. This effect can be seen in Figure 2 that shows the CO<sub>2</sub> total emission accumulation in the three management systems.

Most CO<sub>2</sub> fluxes in G-10 may be associated with greater microbial activity in areas with major plant residue deposition on soil surface. Minimum soil tillage, like the green sugarcane management, provides favorable conditions for the development of microorganisms in the soil surface layer, which increases microbial biomass and CO<sub>2</sub> production (Matias et al., 2009).

Biom results showed similar trends of fluxes in the three experimental areas (Table 2), with higher values of Biom falling in the *high* and *moderate* CO<sub>2</sub> flux groups and lower values in the *low* and *very low* CO<sub>2</sub> flux groups. This indicates a direct participation of the microbial biomass in CO<sub>2</sub> production during soil organic matter (SOM) decomposition (Jenkinson and Ladd, 1981). However, this process is not always favorable for storing soil carbon, and for some authors, the reduction in SOM can be a result of CO<sub>2</sub> fluxes (Cerri et al., 2007).

The physical and chemical soil attributes showed no clear relation with the different groups of CO<sub>2</sub> fluxes, with the exception of macro- and microporosity, where macroporosity was the soil attribute of high frequency in all three areas, mainly to explain the *high* CO<sub>2</sub> fluxes group (Figures 3A, B, C). This can be attributed to gas transport in the soil, because according to Fick's gas diffusion Law, macroporosity provides better conductivity for the CO<sub>2</sub> molecule in the soil, facilitating gas fluxes (Alvenäs and Jansson, 1997; Brito et al., 2009). In turn, microporosity provides a more irregular path hindering CO<sub>2</sub> fluxes in the soil. Thus, G-10 presented greater macroporosity than B and G-5 (Table 2), meaning that despite greater bulk density in G-10 due the mechanized traffic, soil porosity did not impair the fluxes compared to B and this aspect can be attributed to the straw left on the soil that promotes more soil aggregation and porosity.




Figure 2 – Accumulated soil CO<sub>2</sub> emission on soil during Aug-Sept 2011 in burned sugarcane (B), green harvest sugarcane for 5 years (G-5) and 10 years (G-10).

Table 2 – Descriptive statistics of soil microbiological, physical and chemical attributes across experimental areas in accordance to CO<sub>2</sub> flux criterion (high, moderate, low and very low).

	FCO <sub>2</sub>		Biom		Ms		Ts		MWD		Bd		Clay		Ma		Mi		C		V%		pH		P		SD		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
Burned cane (B)	High	2.11	0.25	226.44	21.91	11.87	0.92	24.78	0.3	1.48	0.52	1.17	0.1	557.94	18.34	0.2	0.02	0.37	0.02	2.82	0.41	77.32	3.51	5.26	0.16	14.38	2.92		
	Moderate	1.72	0.07	224.34	35.93	11.58	0.66	24.38	0.37	1.68	0.58	1.19	0.09	568.08	17.12	0.19	0.03	0.38	0.04	2.98	0.29	74.8	2.16	5.17	0.12	18.00	5.73		
	Low	1.48	0.08	215.00	34.83	11.66	1.10	24.31	0.60	1.78	0.58	1.26	0.19	531.50	16.22	0.19	0.03	0.35	0.03	2.90	0.32	77.1	4.22	5.29	0.19	19.63	11.24		
	Very low	0.96	0.27	63.61	12.68	11.00	0.9	25.00	0.30	1.00	0.45	1.00	0.12	536.00	19.11	0.19	0.02	0.38	0.02	3.01	0.22	73.95	3.47	5.13	0.14	16.14	3.02		
Green cane with 5 years (G-5)	High	2.78	0.27	345.92	29.87	24.44	1.79	18.84	0.66	1.69	0.57	1.35	0.18	527.57	13.09	0.26	0.06	0.32	0.03	2.59	0.37	60.29	4.02	4.86	0.10	28.29	7.06		
	Moderate	2.03	0.17	346.57	27.06	25.05	1.82	18.88	0.67	1.17	0.36	1.31	0.11	524.31	19.42	0.19	0.04	0.33	0.02	3.5	0.81	57.71	4.60	4.81	0.16	29.50	11.95		
	Low	1.66	0.06	291.57	17.26	24.29	1.79	18.92	0.23	1.42	0.32	1.35	0.16	505.25	14.66	0.2	0.02	0.33	0.04	3.18	0.74	58.72	5.42	4.76	0.09	36.63	17.74		
	Very low	1.27	0.17	301.79	16.24	23.88	1.65	18.93	0.10	1.30	0.36	1.26	0.14	501.79	11.78	0.18	0.04	0.39	0.09	3.30	0.72	59.71	5.33	4.79	0.12	51.71	13.07		
Green cane with 10 years (G-10)	High	4.02	0.24	342.58	31.24	20.66	1.87	21.58	0.84	1.85	0.57	1.33	0.18	541.60	13.6	0.27	0.11	0.26	0.14	2.66	0.23	62.4	11.58	5.00	0.4	39.29	12.76		
	Moderate	2.84	0.27	355.80	23.44	20.21	1.37	26.03	0.85	1.62	0.46	1.35	0.18	410.63	19.06	0.23	0.03	0.26	0.04	2.65	0.82	57.42	11.68	4.87	0.23	30.71	17.23		
	Low	2.18	0.06	332.61	12.76	19.39	1.42	24.47	0.93	1.61	0.49	1.36	0.13	434.75	19.89	0.25	0.03	0.24	0.08	2.55	0.26	56.66	7.84	4.85	0.19	31.38	19.09		
	Very low	1.56	0.26	314.94	14.55	20.08	0.77	21.65	0.52	1.75	0.55	1.37	0.14	464.20	13.85	0.22	0.03	0.27	0.02	2.46	0.35	58.91	11.94	4.95	0.32	42.50	17.88		

FCO<sub>2</sub> = fluxes of CO<sub>2</sub> (μmol CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>); Biom = microbial biomass (μg C g<sup>-1</sup> d<sup>-1</sup>); Ms = soil moisture (%); Ts = soil temperature (°C); MWD = mean weight diameter (mm); Bd = bulk density (kg m<sup>-3</sup>); Clay (g kg<sup>-1</sup>); Ma = macroporosity (m<sup>3</sup> m<sup>-3</sup>); Mi = microporosity (m<sup>3</sup> m<sup>-3</sup>); C = organic carbon (g kg<sup>-1</sup>); V% = base saturation; P = phosphorus (mg dm<sup>-3</sup>); SD = Standard deviation.

The PCA results showed that the first two principal components, PC1 and PC2, explained respectively 50 and 30 % of the variance for all areas and jointly was responsible for more than 80 % of the variance. A similar result was found in a study by Panosso et al. (2011) on CO<sub>2</sub> fluxes, where the PCs together explained 70 % of the variability of soil attributes (physical and chemical), with PC1 explaining 52 % and PC2, 18 %. This means that the soil attributes included in the two principal components are sufficient to explain the CO<sub>2</sub> flux variations in the soil. This is because the soil attributes used in this study promote the CO<sub>2</sub> flux, such as porosity that makes gas transportation in the soil viable (Xu and Qi, 2001; Kosugi et al., 2007; Brito et al., 2009) and the microbiological attribute that produces CO<sub>2</sub> by microbial respiration during OM decomposition increasing the CO<sub>2</sub> flux (Fang et al., 1998).

### Burned sugarcane

In the burned area, the attributes contributing most to PC1 by order of influence represented by the correlation coefficient were Bd (-0.99), Biom (-0.97) and Mi (0.94) (Table 3). Bd and Biom influenced the *low* CO<sub>2</sub> fluxes (mean 1.48  $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) (Figure 3A), Bd described 15 % of the variability and Biom in 14 %. The positive and negative correlation could indicate that in places with low bulk density and microbial biomass, there was a greater incidence of *low* CO<sub>2</sub> fluxes (Figure 3A), reinforcing the relation between biomass and CO<sub>2</sub>, in areas where biomass was low, there were more *low* CO<sub>2</sub> fluxes. This is because the microorganisms promote the CO<sub>2</sub> flux, as cited previously.

Bulk density influences soil porosity in general, increasing Mi, and the prevalence of more Mi than Ma can hinder gas transportation in the soil and the emission to the surface, resulting in more *low* category of CO<sub>2</sub> fluxes. This effect can be confirmed by the Mi analysis that described 13 % of the variability in *very low* CO<sub>2</sub> fluxes (mean 0.96  $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) (Figure 3A), because Mi hinders the soil gas circulation through the less rectilinear and more irregular paths, diminishing *high* CO<sub>2</sub> fluxes and promoting *very low* CO<sub>2</sub> fluxes.

The PC2 showed that the attributes explaining most of the variance were Ma (-0.91), C (0.84) and Ms (-0.78). Ms explained 14 % of the *high* CO<sub>2</sub> flux class (2.11  $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) (Figure 3A). Similar results were found by Ryu et al. (2009) in soils in California (U.S.A.), where Ms explained 14 % of the CO<sub>2</sub> flux variability and showed a negative correlation with CO<sub>2</sub>. Epron et al. (2006) also found a negative correlation between CO<sub>2</sub> and Ms in a study on CO<sub>2</sub> fluxes from forest soils in French Guiana.

Still, the negative relationship between soil respiration and soil water content contrasted with results that highlighted soil moisture controlling temporal soil respiration (Panosso et al., 2008; Maier et al., 2010; Goutal et al., 2012), but not spatial variation and, in our study, only the spatial variability of CO<sub>2</sub> was analyzed, which explained the non-direct effect of soil moisture on the CO<sub>2</sub> flux.




Figure 3 – Two-dimensional representation of the principal components 1 and 2 (biplot) in areas of B (A), G-5 (B) and G-10 (C), with the description of *high* (H), *moderate* (Md), *low* (L) and *very low* (VL) CO<sub>2</sub> fluxes. Biom = microbial biomass; Ms = Soil water content; Ts = soil temperature; MWD = mean weight diameter; Bd = bulk density; Ma = macroporosity; Mi = microporosity; C = organic carbon content; V% = base saturation; P = phosphorus.

Table 3 – Variance data of the principal components PC1 and PC2 with correlation and ranking of importance of the microbiological, physical and chemical soil attributes.

		PC1		PC2		
		Burned cane (B)				
Variance	Total	53		34		
	Accumulative	53		88		
Variable		Correlation	Ranking	Correlation	Ranking	VTE
Microbiology	Biom	-0.97	2 <sup>nd</sup> (14 %)	0.12	12 <sup>th</sup> (0.3 %)	7 %
	Ms	-0.54	9 <sup>th</sup> (4 %)	-0.78	3 <sup>rd</sup> (14 %)	
	Ts	0.67w	8 <sup>th</sup> (6 %)	-0.63	6 <sup>th</sup> (9 %)	
Physical	MWD	-0.85	4 <sup>th</sup> (11 %)	0.37	9 <sup>th</sup> (3 %)	
	Bd	-0.99	1 <sup>st</sup> (15 %)	0.14	11 <sup>th</sup> (0 %)	58 %
	Clay	0.32	12 <sup>th</sup> (1 %)	-0.44	8 <sup>th</sup> (4 %)	
	Ma	0.41	11 <sup>th</sup> (2 %)	-0.91	1 <sup>st</sup> (19 %)	
	Mi	0.94	3 <sup>rd</sup> (13 %)	0.19	10 <sup>th</sup> (0 %)	
Chemical	C	0.53	10 <sup>th</sup> (4 %)	0.84	2 <sup>nd</sup> (16 %)	
	V	-0.69	7 <sup>th</sup> (7 %)	-0.71	4 <sup>th</sup> (12 %)	33 %
	pH	-0.81	5 <sup>th</sup> (10 %)	-0.57	7 <sup>th</sup> (7 %)	
	P	-0.73	6 <sup>th</sup> (8 %)	0.65	5 <sup>th</sup> (10 %)	
Green cane with 5 years (G-5)						
Variance	Total	54		33		
	Accumulative	54		87		
Variable		Correlation	Ranking	Correlation	Ranking	VTE
Microbiology	Biom	-0.11	12 <sup>th</sup> (0.2 %)	-0.95	2 <sup>nd</sup> (22 %)	8 %
	Ms	-0.45	10 <sup>th</sup> (3 %)	0.86	3 <sup>rd</sup> (18 %)	
	Ts	0.93	2 <sup>nd</sup> (13 %)	-0.12	12 <sup>th</sup> (0.3 %)	
Physical	MWD	-0.74	8 <sup>th</sup> (8 %)	-0.6	5 <sup>th</sup> (8 %)	
	Bd	-0.71	9 <sup>th</sup> (7 %)	0.21	10 <sup>th</sup> (1 %)	58 %
	Clay	-0.85	3 <sup>rd</sup> (11 %)	0.33	8 <sup>th</sup> (2 %)	
	Ma	-0.95	1 <sup>st</sup> (13 %)	-0.3	9 <sup>th</sup> (2 %)	
	Mi	0.81	6 <sup>th</sup> (10 %)	-0.45	7 <sup>th</sup> (4 %)	
Chemical	C	0.78	7 <sup>th</sup> (9 %)	0.6	4 <sup>th</sup> (9 %)	
	V	-0.3	11 <sup>th</sup> (1 %)	-0.95	1 <sup>st</sup> (22 %)	33 %
	pH	-0.82	5 <sup>th</sup> (10 %)	-0.16	11 <sup>th</sup> (0.6 %)	
	P	0.85	4 <sup>th</sup> (11 %)	-0.52	6 <sup>th</sup> (6 %)	
Green cane with 10 years (G-10)						
Variance	Total	50		33		
	Accumulative	50		84		
Variable		Correlation	Ranking	Correlation	Ranking	VTE
Microbiology	Biom	0.64	8 <sup>th</sup> (6 %)	-0.68	4 <sup>th</sup> (11 %)	8 %
	Ms	-0.74	7 <sup>th</sup> (9 %)	0.48	6 <sup>th</sup> (5 %)	
	Ts	0.9	4 <sup>th</sup> (13 %)	0.15	11 <sup>th</sup> (0.5 %)	
Physical	MWD	-0.91	3 <sup>rd</sup> (13 %)	-0.42	7 <sup>th</sup> (4.4 %)	
	Bd	0.15	11 <sup>th</sup> (0.3 %)	0.97	2 <sup>nd</sup> (22 %)	58 %
	Clay	-0.75	6 <sup>th</sup> (9 %)	0.26	10 <sup>th</sup> (1 %)	
	Ma	0	12 <sup>th</sup> (0.0004 %)	-0.99	1 <sup>st</sup> (23 %)	
	Mi	-0.64	9 <sup>th</sup> (6 %)	0.32	9 <sup>th</sup> (2 %)	
Chemical	C	0.24	10 <sup>th</sup> (0.9 %)	-0.76	3 <sup>rd</sup> (14 %)	
	V	-0.78	5 <sup>th</sup> (10 %)	-0.61	5 <sup>th</sup> (9 %)	33 %
	pH	-0.93	2 <sup>nd</sup> (14 %)	-0.36	8 <sup>th</sup> (3 %)	
	P	-0.97	1 <sup>st</sup> (15 %)	0.12	12 <sup>th</sup> (0.3 %)	

Biom = microbial biomass; Ms = soil moisture; Ts = temperature of the soil; MWD = mean weight diameter; Bd = bulk density; Ma = macroporosity; Mi = microporosity; C = organic carbon content; V% = base saturation; P = phosphorus content; VTE = total variance explained.

Soil C content showed a positive correlation with PC2 and an opposite trend with Ma for the group *high* CO<sub>2</sub> fluxes (Figure 3A), which suggests that in places with high CO<sub>2</sub> fluxes, the C trend is decreased, that is, more fluxes and less C storage (Cerri et al., 2007). A similar result was found by Panosso et al. (2012) where the CO<sub>2</sub> results showed a positive correlation with PC2 (0.77) and negative with carbon stock (-0.31). Intense

OM decomposition tends to consume the C available in the soil with increasing CO<sub>2</sub> released by microorganisms. On the other hand, low organic carbon content can be understood as protected and stabilized inside microaggregates (Lenka and Lal, 2013). Moreover, corroborating the results of our study, Fang et al. (1998) detected more CO<sub>2</sub> associated with the low C content in pine soils.

### Green sugarcane with 5 years

For the G-5 treatment, the attributes with greater correlation to PC1 were Ma (-0.95), Ts (0.93), and clay (-0.85) (Table 3). Ma and clay were grouped in the *high* CO<sub>2</sub> class (2.78 μmol CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>) (Figure 3B), where the Ma explained 13 % and clay 11 % of variance. Panosso et al. (2011) showed correlation of -0.73 between clay and PC1 in a study of CO<sub>2</sub> fluxes, with greater clay in the green than in the burned cane, however, the CO<sub>2</sub> showed more significance in PC2, indicating little interaction of clay with CO<sub>2</sub>.

A possible explanation of negative correlation of clay with PC1 to explain the *high* CO<sub>2</sub> flux is that clay promotes a type of carbon protection in the soil, that is, the adsorption of OM with mineral particles (mainly clay minerals and oxides) protects OM from microbial decomposition and prevents C loss as CO<sub>2</sub>, however, if the clay content is low, as shown in our study, protection is smaller, promoting more *high* CO<sub>2</sub> flux situations.

Soil temperature Ts explained 13 % of the variance in *low* CO<sub>2</sub> fluxes (mean of 1.66 μmol CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>) for the G-5 treatment. It is justifiable because soil temperature in G-5 was lower than B and G-10 and low temperature can promote low CO<sub>2</sub> fluxes due the slow microorganism activity. The great performance of soil microorganisms occurred around 30 °C (Kononova, 1975) and in G-5, soil temperature was 18 °C (Table 2). The correlation between CO<sub>2</sub> and Ts was also found in studies of Carbonell-Bojollo et al. (2012), Lenka and Lal (2013), Shrestha et al. (2013) and Song et al. (2013).

In PC2, the attributes that showed higher correlation were V (-0.95), Biom (-0.95) and Ms (0.86). The V and Biom were grouped in the *very low* CO<sub>2</sub> flux class (mean 1.27 μmol CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>) (Figure 3B), indicated by the negative correlation coefficient, with places of low V and Biom with more incidences of *very low* CO<sub>2</sub> fluxes. This is because low base saturation and soil acidity affected the soil microbial activity resulting in lower CO<sub>2</sub> fluxes.

### Green sugarcane with 10 years

The attributes that mostly contributed to the explanation of PC1 on the G-10 were, in order of influence, P (-0.97), pH (-0.93) and MWD (-0.91) (Table 3). In addition, we observed that pH and MWD influenced the *high* CO<sub>2</sub> fluxes group (mean 4.02 μmol CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>) in 14 % and 13 % respectively, with negative correlations with PC1. This indicates that the greater incidence of *high* CO<sub>2</sub> fluxes can be explained by the low values of pH and MWD.

The pH values ranged from 4-5 (Table 2), probably due to the more intense OM decomposition than in the other areas due the greater amount of straw that stimulates microbial decomposition and this process may decrease the pH during the nitrification stage. Xu and Qi (2001) showed negative correlation of CO<sub>2</sub> emission with the pH in a study on spatial variation of soil CO<sub>2</sub> fluxes.

MWD is associated with the physical protection of OM, and soil aggregates play this role by preventing the release of occluded carbon that serves as a source of energy for the microbial biomass. Thus, the negative correlation between MWD with PC1, mainly to explain the *high* CO<sub>2</sub> fluxes, helps to understand that lower MWD can promote less C protection and, thus, *high* CO<sub>2</sub> fluxes.

Some variation (15 %) of CO<sub>2</sub> fluxes within *very low* CO<sub>2</sub> (1.56 μmol CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>) was explained by the P content, with a negative correlation with PC1. This indicates that the increased amount of P in the G-10 area influenced the smaller incidence of *very low* fluxes. The P content stimulates the production of the phosphatase enzyme produced by specific phosphate solubilizing microorganisms (Barroso and Nahas, 2006), resulting in greater CO<sub>2</sub> by microorganisms during decomposition, thus, explaining the smaller incidence of *very low* CO<sub>2</sub> fluxes.

PC2 presented a greater explanation by the attributes Ma (-0.99), Bd (0.97) and C (-0.76). The Ma trend was the same in B and G-5 areas in which *high* CO<sub>2</sub> fluxes were explained by Ma (Table 3). Panosso et al. (2012) studied CO<sub>2</sub> soil fluxes in sugarcane management and also found significant correlation of PC1 with CO<sub>2</sub> (0.77) and with Ma (0.75).

In a study by Leon et al. (2014), the PCA showed that the principal soil attribute responsible for *High* CO<sub>2</sub> fluxes season was root biomass. In our study, Ma was always present and explained the significant variance of *high* CO<sub>2</sub> flux, despite the negative correlation with PC1 (Figure 3B) and PC2 (Figures 3A, C). Ma explained 19 % in B, 13 % in G-5 and 23 % in G-10 of *high* CO<sub>2</sub> fluxes.

Ma has an important relationship with soil CO<sub>2</sub> fluxes, as the greater number of macropores enables soil gas circulation (Brito et al., 2009), furthermore, an increased number of Ma enables greater concentrations of soil O<sub>2</sub>, which stimulates the activity of microbial decomposition and consequent soil CO<sub>2</sub> fluxes. In a study by Goutal et al. (2012), changes in soil macroporosity percentages affected plant roots and associated microbial activities. Thus, soil Ma and Mi influenced possible flux trajectories of soil gases, which affected O<sub>2</sub> and CO<sub>2</sub> fluxes (Brito et al., 2009). According to the Fick's law (Alvenäs and Jansson, 1997), the relations between Ma and CO<sub>2</sub> are controlled by several factors including total porosity, soil water content and tortuosity coefficient.

Other soil attributes interpreted in isolation can cause double interpretation, for example, Bd and C were attributes that presented both positive and negative correlations with the principal components (Table 3). A study of Panosso et al. (2011) concluded that the bulk density associated with the humification index relates better than other properties with soil CO<sub>2</sub> emission, as this property is the most important to understand the emission variability in the area of burned cane.

In some studies, CO<sub>2</sub> and C showed positive correlations (La Scala et al., 2000; Lenka and Lal, 2013; Medeiros et al., 2011) mainly related to the supply of

the substrate to microbial activity. Negative correlations (Fang et al., 1998) associated with the limitation of decomposition due to adverse soil and climate conditions have also been observed, although their influence was not always significant (Epron et al., 2004). According to Song et al. (2013), the increase in CO<sub>2</sub> fluxes associated with soil carbon is complex and may involve both positive and negative feedbacks. This requires further studies of the CO<sub>2</sub> flow and soil carbon dynamics.

## Conclusions

The harvesting of green sugarcane presented higher CO<sub>2</sub> total fluxes than the burned sugarcane did. This effect is associated with greater microbial activity in areas with greater plant waste deposition on the soil surface. On the other hand, the CO<sub>2</sub> fluxes based on *high-low* criteria showed that macroporosity explained the *high* CO<sub>2</sub> fluxes, this is because the greater number of macropores improved soil gas circulation and enabled greater concentrations of soil O<sub>2</sub>, which stimulate the activity of microbial decomposition and consequent soil CO<sub>2</sub> fluxes.

## Acknowledgments

To the São Paulo State Foundation for Research Support (FAPESP - 2010/18.979-5; 2011/04.842-0) for financial support and to the São Martinho ethanol mill for providing the study area.

## References

- Alvenäs, G.; Jansson, P.E. 1997. Model for evaporation, moisture and temperature of bare soil: calibration and sensitivity analysis. *Agricultural and Forest Meteorology* 88: 47-56.
- Ball, B.C.; Tiessen, H.; Stewart, J.W.B.; Salcedo, I.H.; Sampaio, E.V.S.B. 1993. Residue management effects on sugarcane yield and soil properties in northeastern Brazil. *Agronomy Journal* 85: 1004-1008.
- Barroso, C.B.; Nahas, E. 2006. Solubilization of iron phosphate by free or immobilized pellets and spores of *Aspergillus niger*. *Research Journal of Microbiology* 1: 210-219.
- Brito, L.F.; Marques Júnior, J.; Pereira, G.T.; Souza, Z.M. 2009. Soil CO<sub>2</sub> emission of sugarcane fields as affected by topography. *Scientia Agricola* 66: 77-83.
- Carbonell-Bojollo, R.; Torres, M.A.R.R.; Rodriguez-Lizana, A.; Ordóñez-Fernández, R. 2012. Influence of soil and climate conditions on CO<sub>2</sub> emissions from agricultural soils. *Water Air and Soil Pollution* 223: 3425-3435.
- Cerri, C.E.P.; Sparoveki, G.; Bernoux, M.; Easterling, W.E.; Melillo, J.M.; Cerri, C.C. 2007. Tropical agriculture and global warming: impacts and mitigation options. *Scientia Agricola* 64: 83-99.
- Brazilian Agricultural Research Corporation [Embrapa]. 1997. Elisabeth, M., Claessen, C. Manual methods of soil analysis = Manual de análise de solos. 2ed. Embrapa, Rio de Janeiro, RJ, Brazil (in Portuguese).
- Epron, D.; Bosc, A.; Bonal, D.; Freycon, V. 2006. Spatial variation of soil respiration across a topographic gradient in a tropical rain forest in French Guiana. *Journal of Tropical Ecology* 22: 565-574.
- Epron, D.; Nouvellon, Y.; Rounsard, O.; Mouvondy, W.; Mabilia, A.; Saint-Andre, L.; Joffre, R.; Jourdan, J.; Bonnefond, J.M.; Berbigier, P.; Hamel, O. 2004. Spatial and temporal variations of soil respiration in a eucalyptus plantation in Congo. *Forest Ecology and Management* 202: 149-160.
- Fang, C.; Moncrieff, J.B.; Gholz, H.L.; Clark, K.L. 1998. Soil CO<sub>2</sub> efflux and its spatial variation in a Florida slash pine plantation. *Plant and Soil* 205: 135-146.
- Goutal, N.; Parent, F.; Bonnaud, P.; Demaison, J.; Nourisson, G.; Epron, D.; Ranger, J. 2012. Soil CO<sub>2</sub> concentration and efflux as affected by heavy traffic in forest in northeast France. *European Journal of Soil Science* 63: 261-271.
- Jenkinson, D.S.; Ladd, J.N. 1981. Microbial biomass in soil: measurement and turnover. *Soil Biology and Biochemistry* 5: 415-471.
- Jenkinson, D.S.; Powlson, D.S. 1976. The effects of biocidal treatments on metabolism in soil. V. Method for measuring soil biomass. *Soil Biology and Biochemistry* 8: 209-213.
- Johnson, R.A.; Wichern, D.W. 2002. *Applied Multivariate Statistical Analysis*. 5ed. Prentice Hall, Upper Saddle River, NJ, USA.
- Kaiser, H.F. 1958. The varimax criterion for analytic rotation in factor analysis. *Psychometrika* 23: 178-200.
- Kononova, M.M. 1975. Humus of virgin and cultivated soils. In: Gieseking, J.E., ed. *Soil components*. Springer, Berlin, Germany.
- Kosugi, Y.; Mitani, T.; Itoh, M.; Noguchi, S.; Tani, M.; Matsuo, N.; Takahashi, S.; Ohkubo, S.; Nik, A.R. 2007. Spatial and temporal variation in soil respiration in a southeast Asian tropical rainforest. *Agricultural and Forest Meteorology* 147: 35-47.
- La Scala, N.; Mendonça, E.S.; Souza, J.J.; Panosso, A.R.; Simas, F.N.B.; Schaefer, C.E.G.R. 2010. Spatial and temporal variability in soil CO<sub>2</sub>-C emissions and relation to soil temperature at King George Island, Maritime Antarctica. *Polar Science* 4: 4479-487.
- La Scala, N.; Marques Júnior, J.; Pereira, G.T.; Corá, J.E. 2000. Carbon dioxide emission related to chemical properties of a tropical bare soil. *Soil Biology and Biochemistry* 32: 1469-1473.
- Lenka, N.K.; Lal, R. 2013. Soil aggregation and greenhouse gas flux after 15 years of wheat straw and fertilizer management in a no-till system. *Soil and Tillage Research* 126: 78-89.
- Leon, E.; Vargas, R.; Bullock, S.; Lopez, E.; Panosso, A.R.; La Scala, N. 2014. Hot spots, hot moments, and spatio-temporal controls on soil CO<sub>2</sub> efflux in a water-limited ecosystem. *Soil Biology and Biochemistry* 77: 12-21.
- Maier, M.; Schack-Kirchner, H.; Hildebrand, E.E.; Holst, J. 2010. Pore-space CO<sub>2</sub> dynamics in a deep, well-aerated soil. *European Journal of Soil Science* 61: 877-887.
- Medeiros, J.C.; Silva, A.P.; Cerri, C.E.P.; Fracetto, F.J.C. 2011. Linking physical quality and CO<sub>2</sub> emission under long-term no-till and conventional-till in a subtropical soil in Brazil. *Plant and Soil* 338: 5-15.



- Nelson, D.W.; Sommers, L.E. 1982. Total C, organic C and OM. In: Page, R.H.; Kenny, D.R., eds. *Methods of soil analysis. II. Chemical and microbiological properties*. 2ed. Soil Science Society of America, Madison, WI, USA.
- Panosso, A.R.; Perillo, L.I.; Ferraudo, A.S.; Pereira, G.T.; Miranda, J.V.G.; La Scala, N. 2012. Fractal dimension and anisotropy of soil CO<sub>2</sub> emission in a mechanically harvested sugarcane production area. *Soil and Tillage Research* 124: 8-16.
- Panosso, A.R.; Marques Júnior, J.; Milori, D.M.B.P.; Ferraudo, A.S.; Barbieri, M.; Pereira, G.T.; La Scala, N. 2011. Soil CO<sub>2</sub> emission and its relation to soil properties in sugarcane areas under Slash-and-burn and Green harvest. *Soil and Tillage Research* 111: 190-196.
- Panosso, A.R.; Marques Júnior, J.; Pereira, G.T.; La Scala, N. 2009. Spatial and temporal variability of soil CO<sub>2</sub> emission in a sugarcane area under green and Slash-and-burn managements. *Soil and Tillage Research* 105: 275-282.
- Panosso, A.R.; Pereira, G.T.; Marques Junior, J.M.; La Scala, N. 2008. Spatial variability of CO<sub>2</sub> emission on Oxisol soils cultivated with sugarcane under different management practices = Variabilidade espacial da emissão de CO<sub>2</sub> em Latossolos sob cultivo de cana-de-açúcar em diferentes sistemas de manejo. *Engenharia Agrícola* 28: 227-236 (in Portuguese, with abstract in English).
- Raij, B. van; Andrade, J.C.; Cantarella, H.; Quaggio, J.A. 2001. *Chemical Analysis to Evaluate the Fertility of Tropical Soils = Análise Química para Avaliação da Fertilidade de Solos Tropicais*. Instituto Agronômico, Campinas, SP, Brazil (in Portuguese).
- Razafimbelo, T.; Barthès, B.; Larré-Larrouy, M.C.; De Luca, E.F.; Laurent, J.Y.; Cerri, C.C.; Feller, C. 2006. Effect of sugarcane residue management (mulching versus burning) on OM in a clayey Oxisol from southern Brazil. *Agriculture, Ecosystems and Environment* 115: 285-289.
- Rolim, J.S.; Camargo, M.P.B.; Lania, D.G.; Moraes, J.F.F. 2007. Climatic classification of Koppen and Thornthwait systems their applicability in the determination of agroclimatic zoning for the state of São Paulo, Brazil. *Bragantia* 66: 711-720 (in Portuguese, with abstract in English).
- Ryu, S.; Concilio, A.; Chen, J.; North, M.; Ma, S. 2009. Prescribed burning and mechanical thinning effects on belowground conditions and soil respiration in a mixed-conifer forest, California. *Forest Ecology Management* 257: 1324-1332.
- Shrestha, R.K.; Lal, R.; Rimal, B. 2013. Soil carbon fluxes and balances and soil properties of organically amended no-till corn production systems. *Geoderma* 197-198: 117-185.
- Song, Z.; Yuan, H.; Kimberley, M.O.; Jiang, H.; Zhou, G.; Wang, H. 2013. Soil CO<sub>2</sub> flux dynamics in the two main plantation forest types in subtropical China. *Science of the Total Environment* 444: 363-368.
- Xu, M.; Qi, Y. 2001. Soil-surface CO<sub>2</sub> efflux and its spatial and temporal variations in a young ponderosa pine plantation in northern California. *Global Change Biology* 7: 667-677.