

ORIGINAL ARTICLE

Delivery and postpartum care in Rio Branco in the northern state of Acre, Brazil: a population-based survey



Andréa Ramos da Silva Bessa¹, Leila Maria Geromel Dotto¹, Margarida de Aquino Cunha¹, Pascoal Torres Muniz², Suelen de Oliveira Cavalcante³

¹Enfermeira, Doutora, Universidade Federal do Acre - Rio Branco - AC - Brasil

²Nutricionista, Doutor, Universidade Federal do Acre - Rio Branco - AC - Brasil

³Enfermeira, Mestre, Hospital das Clínicas - Rio Branco - AC - Brasil

Corresponding author:
andreaufac@bol.com.br

Manuscript received: Setember 2017

Manuscript accepted: November 2017

Version of record online: March 2018

Abstract

Introduction: Pregnancy can be associated with health risks for both the mother and infant, and specialised care during the pregnancy, delivery and puerperium periods can help reduce complications for the mother-infant binomial.

Objective: To assess the demographic, social and reproductive aspects of delivery and the postpartum period in Rio Branco, Acre, Brazil.

Methods: A population-based cross-sectional survey was conducted via an interview with 552 mothers with children aged 0 to 5 years between 2007 and 2008. Cluster sampling of the population was performed in two stages (census sectors and domiciles).

Results: The majority of the 552 mothers were multiparous (79.6%). Around 70% of pregnancies occurred in teenage mothers. The proportion of caesarean section deliveries was 38.4% in mothers from urban zones and 28.5% in those from rural areas, and was higher in mothers who attended private clinics, who self-reported as being Caucasian, and who had a higher educational level. With regard to puerperal complications, mothers who underwent caesarean section had a higher prevalence of hypertension (71.1%) with an adjusted prevalence ratio of 3.90 (95% CI [2.00, 7.61]).

Conclusions: The results revealed a high rate of teenage pregnancy, and arterial hypertension was the leading complication during the postpartum period experienced by women who had undergone caesarean section. These findings merit further attention, and should be used to improve the care provided to Rio Branco.

Keywords: natural childbirth, cesarean section, postpartum period.

Suggested citation: Bessa ARS, Dotto LMG, Cunha MA, Muniz PT, Cavalcante SO. Delivery and postpartum care in Rio Branco in the northern state of Acre, Brazil: a population-based survey. *J Hum Growth Dev.* 2018; 28(1):69-76. DOI: <http://dx.doi.org/10.7322/jhgd.118010>

■ INTRODUCTION

It has been increasingly recognised that childbirth and the immediate postpartum period are vulnerable periods for both the mother and newborn. It is estimated that between 25% and 45% of neonatal deaths and 45% of maternal deaths occur during the first 24 hours after birth^{1,2}.

Attention to childbirth in Brazil is characterised by high intervention rates, especially caesarean section, ranging from 44%³ to 51.9%⁴ of all births in the country. The socioeconomic level of the mother and the resulting level of care provided in the private or public sectors determine the degree of intervention during childbirth⁵.

Adequate care for women during childbirth plays a key role in ensuring good outcomes during this process, besides reducing the occurrence of severe maternal morbidity and death⁶.

Most pregnancies and births occur without incident. However, all pregnancies present a risk. About 15% of all pregnant women experience some form of life-threatening complication requiring skilled care, and in some cases, successful and safe obstetric intervention can save their lives⁷.

A study that investigated 4560 births that occurred in a public maternity centre in Goiânia, Goiás, Brazil, reported 86 transfers of women (pregnant and postpartum) to intensive care units. Of these women, 52.3% were nulliparous, 73.3% belonged to the age group of 19–35 years, 72.4% underwent a caesarean birth, 81.4% were transferred during the puerperal period, with 82.6% of the transfer indications related to obstetric causes. Gestational hypertension disorders accounted for more than half (57.7%) of obstetric indications, with eclampsia being the cause in half of these cases⁸.

The greatest challenge for maternal care is its unpredictability, especially at the time of childbirth. Complications that can put a woman and child at risk may

arise suddenly, so the presence of a qualified professional is essential to ensure that complications are properly managed⁹.

In Brazil, 89% of deliveries are attended by physicians and 8.3% by obstetric nurses. However, there is still a significant percentage of women without access to skilled care. These include almost 8% of mothers in the North of Brazil who had a live birth within the past 5 years, and more than 13% of those with no schooling. In the North and Northeast regions, deliveries attended by lay midwives represent 5.8% and 3.6% of live births, respectively³.

Data from the Brazilian National Hospital Based Survey, “Nascer no Brasil”, conducted in 2011/2012, revealed that good practices during labour occurred in less than 50% of women, being less frequent in less developed areas in the North and Northeast, in which obstetric and perinatal indicators are considered the worst in the country⁴.

Therefore, recognising the reality of childbirth care in Rio Branco, the capital city of the northern Brazilian state of Acre, is extremely important to identify where and with whom women are giving birth in this municipality, a region with very different health indicators to other regions of the country. We aimed to identify the prevalence of caesarean delivery and the demographics of women who undergo this procedure. This study fills the gap on the characterisation of childbirth care in this municipality. It is also important to identify the complications that affect women during the postpartum period, a critical period in which the risk of mortality is significant.

Thus, the objective of this study was to analyse the characteristics of attention to childbirth and the postpartum period in the city of Rio Branco considering the sociodemographic and reproductive aspects.

■ METHODS

This cross-sectional, population-based study was carried out in the city of Rio Branco, Acre, using data from the Health and Nutrition Survey of Children and Adults of Rio Branco.

Data collection was carried out between 2007 and 2008, in which information on all pregnancies among women with children aged 0 to 5 years was obtained. To that end, one of the inclusion criteria was that the person who responded to the survey was the biological mother.

The sampling model adopted was clustering with two selection stages (census tracts and domiciles). The census tracts used were the 250 sectors used by the Brazilian Institute of Geography and Statistics (IBGE) for the 2000 census, which included the urban and rural areas of the municipality of Rio Branco. Of these, 35 census tracts, 31 urban and four rural sectors, were selected at random.

In each census tract, a total of 875 households were randomly selected to be surveyed, and mothers of children aged 0 to 5 years who lived in the selected household and accepted to participate in the study by providing their free

and informed consent were eligible for participation in the survey. However, it was observed that only 40% of the households had children under the age of 5, leading to the need to draw two more samples of equal size.

A total of 2622 households were randomly selected, in which 723 children under the age of 5 were eligible. However, 3% of the mothers of these children refused to participate in the study. Thus, 552 mothers were interviewed with a total of 648 pregnancies and births, that is, the births of 648 children were analysed.

The data collection instrument included a structured and precoded questionnaire which was used to interview the mothers of children under 5 years old in the participating households.

For this study, the following variables were selected for inclusion in the survey:

- Sociodemographic variables related to the mother: age, schooling, marital status, living area, skin colour, paid activity and number of children;

- Pregnancy-related variables: number of prenatal consultations, woman’s age at first pregnancy and at the

most recent pregnancy, and child death;

-Variables related to childbirth: type of delivery, type of professional who performed the delivery and place of birth (public or private hospital or home);

-Postpartum variables: postpartum complications (haemorrhage, fever, seizure, psychic disorder, hypertension and hospitalisation).

The conditions associated with delivery in Rio Branco were initially analysed in a descriptive way, then the main factors associated with caesarean delivery were investigated. We believed that the attention provided to childbirth would differ between the urban and rural areas of Rio Branco, therefore, we chose to perform the descriptive analysis by area of housing.

Descriptive data analysis was performed to determine the frequency of the studied events. This was followed by estimation of the prevalence ratios of interest

using Poisson regression. Initially, the univariate Poisson regression model was used to select the variables that should compose the multiple model according to the level of statistical significance of entry in the model, which was a p-value of <0.20, introduced by the ENTER method in the model. Those that remained presented higher statistical significance ($p < 0.05$). In the data analysis, the Svy tool or STATA survey module were used to correct the results obtained according to the sampling process performed by clusters.

This study was conducted according to the ethical principles of research established by the National Health Council in resolution no. 196/96. The research only began after approval of the project under process no. 23107.00115/ 2007-22 by the Committee of Ethics in Research at the Federal University of Acre.

RESULTS

Concerning the sociodemographic characteristics, the mean age of the mothers interviewed was 26.7 (standard deviation 6.8) years, with a predominance of women in the 25–34 (45.8%) and 14–24 (42.6%) year age groups. About 73% lived with a partner and 65.8% were housewives. Regarding schooling, 42.7% of the women had between 5–9 years of schooling, 32.8% between 10–15 years, and 24.6% with less than 5 years of schooling. In regard to skin colour, 67.5% defined themselves as brown, 18.0% as black and 14.5% as white.

Regarding the reproductive history of the mothers interviewed, the majority were multiparous (79.6%). The mean age at which women had their first pregnancy was 18.9 (standard deviation 6.3) years in urban areas and 18.0 (standard deviation 3.6) years in rural areas ($p = 0.321$).

The prevalence of home births in Rio Branco was 1.7%, including 3.0% of women living in rural areas and 5.9% of those with lower levels of schooling. The accomplishment of labour according to the type of professional evidences the accomplishment of the same with doctors in 70% of deliveries in the urban area (64.6% in the rural area).

Caesarean delivery was performed in 38.4% of deliveries in urban areas and 28.5% of those in rural areas. The number of deliveries assisted by midwives was 12.4% and 23.5% in urban and rural areas, respectively, and 17.6% and 11.9% of deliveries were assisted by nurses, respectively (Table 1).

In the urban area, about 70% of pregnancies occurred in females under 20 years of age, and 10.4%

Table 1: Distribution of variables related to pregnancy and delivery, according to housing area, in the municipality of Rio Branco - Acre, 2007-2008.

Variables	Urban Zone		Rural zone		p value
	n	(%)	n	(%)	
Birth Local					
Public hospital	540	90,0	45	96,9	
Private hospital	58	9,3	-	-	0,066
Home	04	0,7	01	3,1	
Birth type					
Normal	361	61,6	33	71,5	
Ceasarean	240	38,4	13	28,5	0,008
Who made the delivery					
Doctor	419	70,0	27	64,6	
Nurse	117	17,6	06	11,9	0,171
Midwife	65	12,4	13	23,5	
Activity					
Not payed	384	63,8	42	94,3	
Payed	217	36,2	04	5,7	0,015
Children					
1-2	340	55,0	25	55,6	
3-4	190	33,6	10	17,5	0,039
≥5	72	11,4	11	26,9	
Age at first pregnancy					

07-19	411	69,9	32	65,2	
20-28	174	28,1	14	34,8	0,309
29-37	15	2,0	-	-	
Prenatal appointments					
1-3	42	7,8	06	10,2	
4-5	120	19,4	20	44,7	0,001
≥6	440	72,8	20	45,1	
Child death ≤5 years					
Yes	59	10,4	09	20,8	
No	541	89,6	37	79,2	0,003
Age at last pregnancy					
13-20	168	26,7	14	34,2	
21-28	277	46,4	20	42,8	0,008
29-39	144	26,0	08	18,4	
40-46	10	0,9	02	4,6	

*The differences in absolute values of frequencies correspond to the losses

of the interviewees reported the occurrence of death in a child younger than 5 years. These prevalence rates were 65.0% and 20.9%, respectively, in rural areas.

The prevalence of caesarean delivery (Table 2) in women with less than 5 years of schooling was 25.6%,

which was increased to 48.3% in those with 10–15 years of schooling ($p = 0.001$). Similarly, caesarean deliveries differed according to strata, performed in 32.1% of women who underwent 1–3 prenatal consultations and in 37.5% who had six or more consultations. In regard to the parity

Table 2: Prevalence of normal and caesarean deliveries according to the selected maternal variables in the city of Rio Branco - Acre, 2007-2008.

Variables	Categories	Normal delivery n	Normal delivery (%)	Caesarean delivery n (%)	RP Caesarean delivery	RP adjusted*	
Maternal schooling (years)	0-4	84	74,4	27	25,6	0,53 (0,35-0,80)	0,85 (0,60-1,21)
	5-9	186	71,9	77	28,1	0,58 (0,42-0,79)	0,86 (0,60-1,24)
	10-15	124	51,7	149	48,3	1,00	1,00
Maternal age (years)	14-24	177	67,3	96	32,7	0,91 (0,77-1,09)	0,63 (0,47-0,83)
	25-34	171	64,4	116	35,6	1,00	1,00
	≥35	46	66,5	41	33,5	0,94 (0,59-1,49)	1,00 (0,79-1,25)
Appointments	1-3	36	67,9	12	32,1	0,85 (0,36-2,01)	1,03 (0,49-2,17)
	4-5	96	72,1	44	27,9	0,74 (0,42-1,29)	0,97 (0,50-1,85)
	≥6	262	62,5	197	37,5	1,00	1,00
Skin colour	Black	47	69,3	23	30,7	0,54 (0,35-0,82)	0,71 (0,39-1,27)
	Mixed	289	69,9	171	30,1	0,53 (0,34-0,82)	0,62 (0,38-1,01)
	White	58	43,4	58	56,6	1,00	1,00
Home zone	Rural	33	71,5	13	28,5	0,74 (0,60-0,90)	1,04 (0,69-1,57)
	Urban	361	61,6	240	38,4	1,00	1,00
Place of birth	Public hospital	376	67,8	208	32,2	1,00	1,00
	Home	5	100	-	-	-	-
	Private hospital	13	20,4	45	79,6	2,47 (2,03-3,00)	1,77 (1,41-2,21)
Children	1-2	193	56,5	171	43,5	1,00	1,00
	3-4	137	70,3	62	29,7	0,68 (0,49-0,93)	0,62 (0,44-0,87)
	≥5	64	88,1	19	11,9	0,27 (0,82-0,91)	0,25 (0,77-0,86)

*Prevalence ratio adjusted for age, education, number of prenatal appointments, maternal skin colour, home zone, place of birth and number of children.

distribution, caesarean deliveries were reported in 43.5% of mothers with one or two children, 29.7% of those with three or four children, and 11.9% of mothers with five or more children ($p < 0.001$). The adjusted prevalence ratio for mothers with five or more children versus those with one or two children was 0.25 (95% CI [0.77, 0.86]). The proportion of caesarean deliveries in private institutions was 79.6%, with 32.2% in public hospitals, with an adjusted prevalence ratio of 1.77 (95% CI [1.41, 2.21]).

The prevalence of caesarean delivery in mothers with less than 5 years of schooling was 25.6%, performed in 28.1% of women with 5-9 years and 48.3% of those with 10 or more years of schooling ($p < 0.001$; Table 2).

Table 3 shows that arterial hypertension during puerperium was observed in 71.1% of caesarean deliveries and 28.9% of normal deliveries ($p = 0.001$).

The requirement for hospitalisation during puerperium was reported by 9.5% of women under 25

Table 3: Distribution of frequencies of selected postpartum complications, according to type of delivery, in the city of Rio Branco - Acre, 2007-2008.

Complications	Normal Delivery*		Caesarean Delivery*		p value
	n	%	n	%	
Haemorrhage					
Yes	28	70,1	22	29,9	0,614
No	360	65,4	228	34,6	
Fever					
Yes	43	63,6	28	36,4	0,549
No	348	66,4	221	33,6	
Convulsion					
Yes	04	48,3	04	51,7	0,337
No	386	65,9	246	34,1	
Psychic dist.					
Yes	54	63,7	46	36,3	0,642
No	336	66,2	204	33,8	
Hypertension					
Yes	21	28,9	34	71,1	0,001
No	370	68,1	216	31,9	
Hospitalization					
Yes	18	64,1	31	35,9	0,912
No	372	66,0	218	34,0	

*The differences in the absolute values of the frequencies correspond to the losses

years of age, 4.2% of those aged 25 to 34 years, and 21.2% of mothers aged 35 years or older ($p = 0.005$; Table 4).

Postpartum hypertension was 71.1% in mothers who underwent caesarean delivery, with an adjusted

prevalence ratio of 3.90 (95% CI [2.00, 7.61]). Hypertension also affected 60.3% of women between the ages of 14 and 24 years, with an adjusted prevalence ratio of 2.2 (95% CI [1.01, 4.80]; Table 5).

Table 4: Frequency distribution of postpartum complications, according to the selected maternal variables, in the city of Rio Branco - Acre, 2007-2008

Variables	Fever*		Haemorrhage *		Hypertension*		Psychic dist.*		Hospitalization*	
	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)
Schooling (years)										
0-4	15	9,6	10	3,7	09	3,3	12	7,6	10	3,5
5-9	28	6,5	16	6,3	18	2,9	52	16,6	19	13,4
10-15	29	8,9	24	8,3	28	11,1	37	17,6	20	5,6
p value	0,617		0,357		0,102		0,283		0,092	
Age (years)										
14-24 years	33	10,3	20	8,9	25	8,1	37	15,1	16	9,5
25-34 years	35	7,5	24	4,5	23	3,7	47	13,8	23	4,2
≥35 years	03	2,1	05	4,3	07	4,7	17	17,1	10	21,2
p value	0,066		0,241		0,186		0,796		0,050	
Children										
1-2	36	7,9	24	7,0	30	6,4	51	14,5	22	8,8
3-4	27	9,2	19	6,8	22	7,5	35	13,2	20	6,6
≥5	08	6,7	07	3,7	03	1,0	15	17,6	07	10,1
p value	0,807		0,609		0,037		0,742		0,522	

*The differences in the absolute values of the frequencies correspond to the losses

Table 5: Prevalence of postpartum hypertension according to the selected maternal variables in the municipality of Rio Branco - Acre, 2007-2008

Variables	Categories	Hypertension		RP Hypertension	RP adjusted* Hypertension
		n	(%)		
Birth type	Normal	21	28,9	1,00	1,00
	Cesárea	34	71,1	4,73 (1,77-12,59)	3,90 (2,00-7,62)
Maternal age	14-24 anos	25	60,3	2,18 (0,76-6,24)	2,20 (1,01-4,80)
	25-34 anos	23	30,1	1,00	1,00
	≥ 35 anos	07	9,6	1,26 (0,39-4,05)	0,88 (0,28-2,73)
Maternal skin colour	Negra	06	5,8	0,11 (0,11-1,07)	0,20 (0,24-1,66)
	Parda	35	51,5	0,25 (0,51-1,29)	0,40 (0,12-1,29)
	Branca	14	42,7	1,00	1,00
Home zone	Rural	01	23,9	1,00	1,00
	Urbana	54	76,1	2,45 (0,80-7,49)	2,06 (0,60-7,03)

*Prevalence ratio adjusted for age, type of delivery, skin colour and home area

DISCUSSION

The sample included in this study can be considered representative of the population of mothers of children under 5 years old in Rio Branco as it was performed as a population-based survey, supported by the low rate of loss and refusals.

The findings of the present study indicate that there is a higher probability of performing a caesarean delivery in women who give birth at private hospitals or in maternity hospitals, in those with a higher level of educational, in women who define themselves as white, those who live in urban areas and who are older than 25 years.

The prevalence of caesarean deliveries in public and private institutions was high in relation to the 15% prevalence reported by the World Health Organization (WHO). However, a higher prevalence has also been reported in other studies. In Santa Catarina, the prevalence of caesarean section was found to be 61.3% in public institutions and 100% in private institutions^{10,11}.

A national survey⁴ carried out with 23,894 puerperae supports the findings of the present investigation, in which it was evidenced that caesarean section was less frequent among non-white public sector users with lower levels of education and who were multiparous. This reinforces a pattern previously described in Brazil¹².

The average number of children per woman in Rio Branco is higher than the national level and that of the North region, which is 1.8 and 2.28 children, respectively³.

It is worth highlighting the universality of hospital delivery, especially public hospitals. Furthermore, we found a smaller proportion of births in Acre compared to that reported in the North region of 7.5%³.

The vast majority of deliveries were performed by a formally qualified professional (physician or nurse). However, a significant proportion of women (12.4% in urban areas and 23.5% in rural areas) had their births attended by traditional birth attendants, especially those

with lower levels of education, those living in rural areas of the municipality and those who had undergone fewer prenatal consultations. This is higher than the prevalence of births performed by traditional midwives in the North and Northeast of 5.8% and 3.6%, respectively³.

As for complications experienced by the studied women during the postpartum period, psychiatric disorders (mental/emotional problems, depression, loss of judgment) had the highest prevalence. According to the literature, the pregnancy-puerperal period is associated with a high incidence of psychiatric disorders, affecting 10–20% of puerperal women¹³. These disorders, known as maternal or postpartum depression, involve an emotional, humoral and reactive disorder during the puerperium period¹⁴.

The prevalence of reports of hypertension during puerperium was higher in women who underwent caesarean delivery. Although the presence of hypertension in pregnancy does not necessarily require a caesarean delivery, several studies have shown that hypertension is one of the most cited reasons for performing caesarean deliveries as hypertensive syndromes are the leading cause of maternal death in Brazil, accounting for 15% of deaths¹⁵.

The association between advanced age and caesarean section reported in several studies, such as in Santa Catarina^{10,16}, was not found in this survey. We observed a higher prevalence of caesarean section in young women.

The results of the “Nascer no Brasil” study showed a high proportion of caesarean sections among adipose primiparas (40%). The factors most strongly associated with caesarean section being considered a safer delivery method included childbirth funded by the private sector, the same health professional attending prenatal consultations and childbirth, and presentation of clinical indications of risk and intercurrents during pregnancy. Adolescent pregnancy remains a topic of discussion in the

area of reproductive health, and the proportion of surgical delivery in these women is worrying due to early exposure to the effects of caesarean¹⁷. The findings of this survey also indicate that, according to the geographic region analysis, the chance of a pregnancy resulting in caesarean section is higher in the North region when compared to other regions of the country, probably due to the low coverage provided by health plans and the high caesarean rate in the public and private services of this region⁴.

It was found that a higher prevalence of births were attended by traditional midwives in women aged 35 years or over. This is worrying as it is known that pregnancies in this age group are considered riskier because they are associated with a higher incidence of maternal complications, such as greater weight gain, obesity, gestational diabetes, hypertension and preeclampsia¹⁵.

The association between a higher education level and caesarean section found in the present study is in line with other studies in the literature conducted in Chile and Santa Catarina^{10,16}.

Most births took place in public institutions. The groups with the highest number of childbirths in public institutions included pregnant women aged 14 to 24 years, those with lower levels of schooling, self-reported black skin colour, those living in rural areas of the city and having had fewer prenatal consultations.

It is worth noting the observed prevalence of home births in the sample, which were more frequent in women in the rural areas of the municipality and in those with lower schooling.

Taken together, the results observed in this study reveal the need to expand the debate on caesarean deliveries, the prevalence of which are much higher in Rio Branco than that recommended by the WHO. This high rate of caesarean section deliveries are associated with important postpartum complications such as prolonged hospitalisation of puerperal, an increased cost of care, and causing controllable damage to the health of the puerperal population, such as those described in this research.

In order to improve assistance during the pregnancy- puerperal cycle, as well as the quality of life of the mother-

baby binomial, interventions in the field of obstetric care should be guided based on scientific evidence in both public and private institutions.

The findings of the present study highlight an issue related to the high number of caesarean deliveries performed in the city of Rio Branco, as well as the repercussions for these women during the puerperal phase, with arterial hypertension occurring in 71.1% of those who underwent caesarean section.

The increase in the incidence of caesarean section represents an important public health problem, especially in Brazil, where caesarean rates are already unacceptably high and may contribute to an increase in the rates of maternal and perinatal morbidity and mortality.

There is a clear need to further study and discuss this public policy issue in Rio Branco, Acre, Brazil, in an attempt to develop strategies to increase access to qualified prenatal care and provide a link between outpatient care and childbirth.

Some limitations of the present investigation should be considered. This study was performed as a survey in which data were collected from pregnancies over the past 5 years, which may generate information and memory bias.

Acknowledgements

Thanks to our heartfelt thanks to Prof. Sérgio Koifman ("In Memoriam"), for his invaluable contribution to the design, planning and construction of the research project, data collection, analysis and interpretation of the data.

The realization of this investigation was made possible through the academic collaboration established between the program of Masters in collective health of the Federal University of Acre and the program of post-graduation in public health and environment of Fundação Oswaldo Cruz, the which has been supported with the resources of the National Council of Scientific Research and development-CNPq (edictal blanket UFAC-FIOCRUZ, Case No. 620024/2008-9); CAPES (Programs PROCAD-NF 1442/2007 and PROCAD-NF 2557/2008) and Ministry of Health.

REFERENCES

1. Lawn JE, Cousens S, Zupan K; Lancet Neonatal Survival Steering Team. 4 million neonatal deaths: When? Where? Why? *Lancet*. 2005;365(9462):891-900. DOI: [http://dx.doi.org/10.1016/S0140-6736\(05\)71048-5](http://dx.doi.org/10.1016/S0140-6736(05)71048-5)
2. World Health Organization (WHO). Health and the millennium development goals. Geneva: World Health Organization, 2005.
3. Brasil. Centro Brasileiro de Análise e Planejamento. Pesquisa Nacional de Demografia e Saúde da Criança e da Mulher: PNDS 2006. Dimensões do processo reprodutivo e da saúde da criança. Brasília: Ministério da Saúde, 2009.
4. Leal MC, Pereira APE, Domingues RMSM, Theme Filha MM, Dias MAB, Nakamura-Pereira M, et al. Intervenções obstétricas durante o trabalho de parto e parto em mulheres brasileiras de risco habitual. *Cad Saúde Pública*. 2014;30(suppl 1):S17-32. DOI: <http://dx.doi.org/10.1590/0102-311X00151513>
5. Roberts CL, Tracy S, Peat B. Rates of obstetric intervention among private and public patients in Australia: population based descriptive study. *BMJ*. 2000;321(7254):137-41. DOI: <https://dx.doi.org/10.1136/bmj.321.7254.137>
6. Cecatti JG. Saúde da mulher: Enfoque da evidência científica para a prevenção da morbidade e mortalidade materna. *Rev Bras Saude Mater Infant*. 2005;5(1):9-11.

7. World Health Organization (WHO). Managing complications in pregnancy and childbirth: a guide for midwives and doctors. Geneva: WHO/ UNFPA/ UNICEF/ World Bank, 2000.
8. Viggiano MB, Viggiano MGC, Souza E, Camano L. Necessidades de cuidados intensivos em maternidade pública terciária. Rev Bras Ginecol Obstet. 2004; 26(4):317-23. DOI: <http://dx.doi.org/10.1590/S0100-72032004000400009>
9. Starrs A. The safe motherhood action agenda: priorities for the next decade. Proceedings. Washington, DC: World Bank, 1997.
10. World Health Organization. Appropriate technology for birth. Lancet. 1985; 2(84532):436-67. DOI: [https://doi.org/10.1016/S0140-6736\(85\)92750-3](https://doi.org/10.1016/S0140-6736(85)92750-3)
11. Bofante TM, Silveira GC, Sakae TM, Sommacal LF, Fedrizzi EN. Fatores associados à preferência pela operação cesariana entre puérperas de instituição pública e privada. Arq Catarin Med. 2009;38(1):26-32.
12. Domingues RMSM, Dias MAB, Nakamura-Pereira M, Torres JA, d'Orsi E, Pereira APE, et al. Processo de decisão pelo tipo de parto no Brasil: da preferência inicial das mulheres à via de parto final. Cad Saude Publica. 2014; 30(suppl1): S101-16. DOI: <http://dx.doi.org/10.1590/0102-311X00105113>
13. Félix GMA, Gomes APR, França PS. Depressão no ciclo gravídico-puerperal. Comun Ciencias Saude. 2008;19(1):51-60.
14. Holmes, DS. Psicologia dos transtornos mentais. 2ed. Porto Alegre: Artmed; 2001.
15. Queiroz MVO, Jorges MSB, Marques JF, Cavalcante AM, Moreira KAP. Indicadores de qualidade da assistência ao nascimento baseado na satisfação de puérperas. Texto Contexto Enferm. 2007;16(3):479-87. DOI: <http://dx.doi.org/10.1590/S0104-07072007000300014>
16. Potter JE, Berquo E, Perpetuo IH, Leal OF, Hopkins K, Souza MR, et al. Unwanted caesarean sections among public and private patients in Brazil: prospective study. BMJ. 2001;323(7322):1155-8. DOI: <https://dx.doi.org/10.1136/bmj.323.7322.1155>
17. Gama SGN, Viellas EF, Schilithz AOC, Theme Filha MM, Carvalho ML, Gomes KRO, et al. Fatores associados à cesariana entre primíparas adolescentes no Brasil, 2011-2012. Cad Saude Publica. 2014; 30(suppl 1): S117-27. DOI: <http://dx.doi.org/10.1590/0102-311X00145513>

Resumo

Introdução: A maternidade pode estar associada a riscos para a saúde da mulher e da criança, de forma que uma assistência especializada durante a gravidez, parto e puerpério contribuem para minimizar as complicações para o binômio mãe-bebê.

Objetivo: Analisar as características do parto e pós-parto em Rio Branco em relação aos aspectos demográficos, sociais e reprodutivos.

Método: Estudo transversal de base populacional realizado durante 2007-2008 através de amostragem populacional por conglomerados em duas etapas (setores censitários e domicílios), sendo entrevistadas 552 mães de crianças de zero a cinco anos.

Resultados: Das 552 mães entrevistadas, a maioria era constituída por múltíparas (79,6%). Cerca de 70% das gestações ocorreram em menores de 20 anos de idade. A prevalência de cesarianas foi de 38,4% nas mães da zona urbana e de 28,5% nas que moravam na zona rural, sendo maior nas mulheres assistidas em instituições privadas, que se auto definiram como de cor branca e com maior nível educacional. Quanto às complicações do período puerperal as mulheres que foram submetidas a partos cesáreas apresentaram uma maior prevalência de hipertensão (71,1%), sendo observada uma razão de prevalência ajustada de 3,90 (IC 95%: 2,00-7,61).

Conclusão: O alto índice de gestação na adolescência e a hipertensão arterial sendo a principal complicação no pós-parto cesárea, dados que merecem atenção e que devem ser observados na assistência prestada no município de Rio Branco.

Palavras-chave: parto normal, cesárea, período pós-parto.

©The authors (2018), this article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.