

A counterexample to the existence of a Poisson structure on a twisted group algebra

Eliana Zoque¹

Department of Mathematics, The University of Chicago, 5734 S. University Avenue, Chicago, Illinois 60637

E-mail address: elizoque@math.uchicago.edu

Abstract. Crawley-Boevey [1] introduced the definition of a noncommutative Poisson structure on an associative algebra A that extends the notion of the usual Poisson bracket. Let (V, ω) be a symplectic manifold and G be a finite group of symplectimorphisms of V . Consider the twisted group algebra $A = \mathbb{C}[V] \# G$. We produce a counterexample to prove that it is not always possible to define a noncommutative poisson structure on $\mathbb{C}[V] \# G$ that extends the Poisson bracket on $\mathbb{C}[V]^G$.

1. Introduction

Crawley-Boevey [1] defined a noncommutative Poisson structure on an associative algebra A over a ring K as a Lie bracket $\langle -, - \rangle$ on $A/[A, A]$ such that for each $a \in A$ the map $\langle \bar{a}, - \rangle : A/[A, A] \rightarrow A/[A, A]$ is induced by a derivation $d_a : A \rightarrow A$; i.e. $\langle \bar{a}, \bar{b} \rangle = \overline{d_a(b)}$ where the map $a \mapsto \bar{a}$ is the projection $A \rightarrow A/[A, A]$. When A is commutative a noncommutative Poisson structure is the same as a Poisson bracket.

Let (V, ω) be a symplectic manifold, with the usual Poisson bracket $\{-, -\}$ on $\mathbb{C}[V]$. Let G be a finite group of symplectimorphisms of V . Consider the twisted group algebra $A = \mathbb{C}[V] \# G$. The algebra of G -invariant polynomials $\mathbb{C}[V]^G$ is contained in $A/[A, A]$. We produce a counterexample to prove that it is not always possible to define a noncommutative poisson structure on $\mathbb{C}[V] \# G$ that extends the Poisson bracket on $\mathbb{C}[V]^G$.

¹The author is grateful to William Crawley-Boevey for a careful review of this paper, and to Victor Ginzburg for posing the question.

2. Twisted group algebra and derivations

From now on, let $A = \mathbb{C}[V] \# G$. (\mathbb{C} can be replaced by any field of characteristic 0.) We use the symbol ${}^g\psi$ to denote the left action of $g \in G$ on $\psi \in \mathbb{C}[V]$. For every $g \in G$ we denote $(-)_g$ the projection $A \rightarrow \mathbb{C}[V]$ into the g -part, i.e., $(\psi h)_g = \psi \delta_{g,h}$ if $\psi \in \mathbb{C}[V], h \in G$. Let $G = C_0 \cup C_1 \cup \dots$ be the conjugacy classes of G , with $C_0 = \{1\}$.

It is proved in [4] that

$$\frac{A}{[A, A]} = HH_0(A) = (HH_0(\mathbb{C}[V], \mathbb{C}[V] \# G))^G,$$

therefore

$$\begin{aligned} \frac{A}{[A, A]} &= \left(\bigoplus_{g \in G} HH_0(\mathbb{C}[V], \mathbb{C}[V]g) \right)^G \\ &= \left(\bigoplus_{g \in G} \frac{\mathbb{C}[V]}{\langle \varphi - {}^g\varphi : \varphi \in \mathbb{C}[V] \rangle} g \right)^G \\ &= \bigoplus_i \left(\frac{\mathbb{C}[V]}{\langle \varphi - {}^{g_i}\varphi : \varphi \in \mathbb{C}[V] \rangle} \right)^{G_{g_i}} g_i \end{aligned}$$

where g_i is an arbitrary element of C_i and $G_g = \{h \in G | gh = hg\}$. The first summand is precisely $\mathbb{C}[V]^G$. Let P_i be the projection

$$A \rightarrow \left(\frac{\mathbb{C}[V]}{\langle \varphi - {}^{g_i}\varphi : \varphi \in \mathbb{C}[V] \rangle} \right)^{G_{g_i}} g_i.$$

The Poisson bracket gives us a family of derivations $d_\psi : \mathbb{C}[V]^G \rightarrow \mathbb{C}[V]^G, \phi \mapsto \{\psi, \phi\}$ for $\psi \in \mathbb{C}[V]^G$; and we want to extend it to a larger family. The following Lemma restricts the possibilities.

Lemma 1. Let $d : A \rightarrow A$ be any derivation. If $x \in \mathbb{C}[V]^g \neq \mathbb{C}[V]$ then $(d(x))_g = 0$.

Proof. Let $y \notin \mathbb{C}[V]^g$. The equality $d(xy) = d(yx)$ implies $d(x)y + xd(y) = d(y)x + yd(x)$. The g -part of this equality is

$$(d(x))_g g y + x(d(y))_g g = (d(y))_g g x + y(d(x))_g g$$

or

$$(d(x))_g {}^g y + x(d(y))_g = (d(y))_g {}^g x + y(d(x))_g.$$

Since ${}^g x = x, {}^g y \neq y$ we conclude $(d(x))_g ({}^g y - y) = 0$, so $(d(x))_g = 0$ \square

Therefore if the action of G on V is faithful and $g \neq 1$, the g -part of the derivative an element of $\mathbb{C}[V]^g$ is zero. This implies that for every $\psi \in \mathbb{C}[V]^G$, $d(\psi) \in \mathbb{C}[V] \subset A$.

The condition $\langle \overline{\psi g}, \overline{\phi h} \rangle = -\langle \overline{\phi h}, \overline{\psi g} \rangle$ implies $\overline{d_{\psi g}(\phi h)} = -\overline{d_{\phi h}(\psi g)}$. Consider the case $\phi, \psi \in \mathbb{C}[V]^G$, $h = 1$ and $g \in C_i, i \neq 0$. Since $P_i(d_{\psi g}(\phi)) = 0$, we must have $0 = P_i(d_\phi(\psi g)) = P_i(d_\phi(\psi)g + \psi d_\phi(g))$. The only terms that must be taken into account are $d_\phi(\psi)g + \psi \sum (d_\phi(g))_{hgh^{-1}} hgh^{-1}$. Modulo $[A, A]$ this is equal to

$$\left(d_\phi(\psi) + \sum_h^{h^{-1}} \left(\psi (d_\phi(g))_{hgh^{-1}} \right) \right) g = (d_\phi(\psi) + \psi \sigma_{\phi,g}) g$$

where $\sigma_{\phi,g} = \sum_h^{h^{-1}} \left((d_\phi(g))_{hgh^{-1}} \right)$ does not depend on ψ .

We want $0 = P_i((d_\phi(\psi) + \psi \sigma_{\phi,g}) g) = P_i((\{\phi, \psi\} + \psi \sigma_{\phi,g}) g)$ since we want a Poisson structure extending the usual Poisson bracket on $\mathbb{C}[V]^G$. Therefore a necessary condition for the existence of the Poisson structure is the existence of $\sigma_{\phi,g} \in \mathbb{C}[V]$ so that

$$P_i((\{\phi, \psi\} + \psi \sigma_{\phi,g}) g) = 0 \tag{1}$$

for every $\psi \in \mathbb{C}[V]$. We will see that this is not always possible.

3. The counterexample

Let $V = \mathbb{C}^4$ with linear coordinates $\{x_1, x_2, x_3, x_4\}$ and the symplectic form $\omega = dx_1 \wedge dx_2 + dx_3 \wedge dx_4$, so $\mathbb{C}[V] = \mathbb{C}[x_1, x_2, x_3, x_4]$, and

$$\{\phi, \psi\} = \frac{\partial \phi}{\partial x_1} \frac{\partial \psi}{\partial x_2} - \frac{\partial \phi}{\partial x_2} \frac{\partial \psi}{\partial x_1} + \frac{\partial \phi}{\partial x_3} \frac{\partial \psi}{\partial x_4} - \frac{\partial \phi}{\partial x_4} \frac{\partial \psi}{\partial x_3}.$$

Let $G = \mathbb{Z}_2 \times (\mathbb{Z}_2 \oplus \mathbb{Z}_2)$. Let e, b, c be the generators of the three copies of \mathbb{Z}_2 (in that order). G acts on V as follows: b and c act as $diag(-1, -1, 1, 1)$ and $diag(1, 1, -1, -1)$, respectively, on $\{x_1, x_2, x_3, x_4\}$ and e interchanges $x_1 \leftrightarrow x_3, x_2 \leftrightarrow x_4$. Clearly $\mathbb{C}[V]^G$ is the set of all polynomials $\sum \lambda_{i_1, i_2, i_3, i_4} x_1^{i_1} x_2^{i_2} x_3^{i_3} x_4^{i_4}$ such that $\lambda_{i_1, i_2, i_3, i_4} \neq 0$ implies $i_1 + i_2, i_3 + i_4$ are even and $\lambda_{i_1, i_2, i_3, i_4} = \lambda_{i_3, i_4, i_1, i_2}$.

Using Magma (<http://magma.maths.usyd.edu.au>) we find that the ring of invariant polynomials is generated, as an algebra, by $f_1 = x_1^2 + x_3^2, f_2 = x_2^2 + x_4^2, f_3 = x_1^4 + x_3^4, f_4 = x_2^4 + x_4^4, h_1 = x_1 x_2 + x_3 x_4, h_2 = x_1^2 x_2^2 + x_3^2 x_4^2, h_3 = x_1^2 x_3 x_4 + x_1 x_2 x_3^2, h_4 = x_1 x_2 x_4^2 + x_2^2 x_3 x_4$; with relations

$$\begin{aligned}
& -f_1f_2h_1 + f_1h_4 + f_2h_3 - h_1^3 + 2h_1h_2, \\
& \frac{1}{2}f_1^2f_2 + \frac{1}{2}f_1h_1^2 - \frac{1}{2}f_1h_2 - \frac{1}{2}f_2f_3 - h_1h_3, \\
& \frac{1}{2}f_1f_2^2 - \frac{1}{2}f_1f_4 + \frac{1}{2}f_2h_1^2 - \frac{1}{2}f_2h_2 - h_1h_4, \\
& -\frac{1}{2}f_1^2f_4 + f_1f_2h_2 - \frac{1}{2}f_2^2f_3 + f_3f_4 - h_2^2, \\
& -\frac{1}{2}f_1^2h_4 + \frac{1}{2}f_1f_2h_3 + \frac{1}{2}f_1h_1h_2 - \frac{1}{2}f_2f_3h_1 + f_3h_4 - h_2h_3, \\
& \frac{1}{2}f_1f_2 * h_4 - \frac{1}{2}f_1f_4h_1 - \frac{1}{2}f_2^2h_3 + \frac{1}{2}f_2h_1h_2 + f_4h_3 - h_2h_4, \\
& \frac{1}{2}f_1^3f_2 + \frac{1}{2}f_1^2h_1^2 - f_1^2h_2 - \frac{1}{2}f_1f_2f_3 - \frac{1}{2}f_3h_1^2 + f_3h_2 - h_3^2, \\
& \frac{1}{2}f_1^2f_2^2 - 3/4f_1^2f_4 + \frac{1}{2}f_1f_2h_1^2 - 3/4f_2^2f_3 + f_3f_4 - \frac{1}{2}h_1^2h_2 - h_3h_4, \\
& \frac{1}{2}f_1f_2^3 - \frac{1}{2}f_1f_2f_4 + \frac{1}{2}f_2^2h_1^2 - f_2^2h_2 - \frac{1}{2}f_4h_1^2 + f_4h_2 - h_4^2.
\end{aligned}$$

Proposition 2. The Poisson bracket on $\mathbb{C}[V]^G$ cannot be extended to a Poisson structure on $\mathbb{C}[V]\#G$ for V and G as defined above.

Proof. Take $\phi = x_1^2 + x_3^2$, $\psi = x_1x_2 + x_3x_4 \in \mathbb{C}[V]^G$ and $g = b$. In this case $\{\phi, \psi\} = 2x_1^2 + 2x_3^2$, $\langle \varphi - {}^g\varphi : \varphi \in \mathbb{C}[V] \rangle = \langle x_1, x_2 \rangle$ and $G_b = \{1, b, c, bc\}$. Hence,

$$\left(\frac{\mathbb{C}[V]}{\langle \varphi - {}^g\varphi : \varphi \in \mathbb{C}[V] \rangle} \right)^{G_g} = \mathbb{C}[x_3, x_4]^{\{1, b, c, bc\}} = \mathbb{C}[x_3^2, x_3x_4, x_4^2],$$

so $P_i((\{\phi, \psi\})b) = 2x_3^2b$

On the other hand, $P_i((\psi\sigma_{\phi, g})b) = P_i(((x_1x_2 + x_3x_4)\sigma_{\phi, g})b) = P_i(((x_3x_4)\sigma_{\phi, g})b)$ and none of the terms here can be equal to $-2x_3^2$ since they all contain x_4 . This contradicts (1). \square

References

- [1] W. Crawley-Boevey. *A note on commutative Poisson structures*, AirXiv: math.QA/0506268.
- [2] A. Cannas da Silva. *Lectures on symplectic geometry*, Lecture Notes in Mathematics, 1764, Springer-Verlag, Berlin, 2001.
- [3] J.J. Cannon, C. Playoust, *An Introduction to Algebraic Programming in Magma*, Sydney: School of Mathematics and Statistics, University of Sydney, 1996.

- [4] V. Dolgushev, P. Etingof, *Hochschild cohomology of quantized symplectic orbifolds and the Chen-Ruan cohomology*, Int. Math. Res. Not. 2005, no. 27, 1657-1688, also math.QA/0410562.