

doi: <http://dx.doi.org/10.11606/issn.1679-9836.v97i1p12-17>

Medical education assessment

Avaliação em educação médica

Cristina P. Camargo¹, Rolf Gemperli², José Otavio Costa Auler Junior³

Camargo CP, Gemperli R., Auler Junior JOC. Medical education assessment / *Avaliação em educação médica*. Rev Med (São Paulo). 2018 jan.-fev.;97(1):12-7.

Abstract: One of the most critical steps for medical education is the assessment. The assessment can be divided into short-term memory, long-term memory, and retrieval memory. If the student acquires all these memories steps, the surgical skill will be mind incorporate for years. As a healthcare provider, the medical community needs to transform training and learning to a valid and reliable activity. Most of the medical evaluations are subjective; an objective assessment is difficult but most desirable. The authors described some objective surgical skill assessment based on OSAST, dexterity and global rating scale. Moreover, we discussed the formative and summative assessment roles to the medical learning process.

Keywords: Education, medical/methods; Evaluation/methods; General surgery/education; Learning; Training.

RESUMO: Um dos aspectos mais críticos na educação em medicina é a avaliação de novos conhecimentos e habilidades. Essa avaliação deve testar memórias curta, longa e de recuperação. Quando o aluno obtiver todas essas etapas, a memória perdurará por anos permitindo a realização de atividades cirúrgicas eficientes e seguras a longo prazo. A maioria das avaliações são subjetivas, embora difícil de se realizar a avaliação objetiva é modalidade de ensino ideal. Esse artigo descreveu algumas opções de avaliações objetivas como OSAST, destreza e escala de taxa global. Além disso, foi abordado os valores da avaliação somativa e formativa no processo de aprendizado do estudante de medicina.

Descritores: Educação médica/métodos; Avaliação/métodos, Cirurgia geral/educação, Ensino; Aprendizagem.

INTRODUCTION

One of the most important steps for medical learning is the assessment. As a final goal, an appropriate assessment allows a better medical service for the community¹⁻³.

Surgical competence assessment evaluate technical skills, decision making and communication skills as well. The assessment can be done during the learning process (formative assessment) and at the end of the training

(summative assessment). The former assessment will provide data that enables the faculty interference to improve learning. The summative process will show if the student achieves the competence¹⁻³.

Nowadays, competence is difficult to measure because most of surgical evaluations are subjective, lacks validity and reliability. In the medical learning process an objective evaluation is desirable. Only with an objective evaluation the clerkship is able to correct and train the students⁴ (Table 1).

1. Disciplina de Cirurgia Plástica, Faculdade de Medicina FMUSP, Universidade de São Paulo, São Paulo, SP, BR. Division of Plastic Surgery, Department of Surgery, School of Medicine, Laboratory of Microsurgery and Plastic Surgery (LIM-04), University of São Paulo, SP, Brazil. <https://orcid.org/0000-0002-3134-0003>

2. Disciplina de Cirurgia Plástica, Faculdade de Medicina FMUSP, Universidade de São Paulo, São Paulo, SP, BR. Full Professor and Head, Division of Plastic Surgery, Department of Surgery, School of Medicine USP, São Paulo, SP, Brazil. Email: rolf.gemperli@fm.usp.br; <https://orcid.org/0000-0001-9913-6079>.

3. Faculdade de Medicina FMUSP, Universidade de São Paulo, São Paulo, SP, BR. Dean of FMUSP, Full Professor and Head, Division of Anesthesiology, School of Medicine, Laboratory of Anesthesiology (LIM-08), University of São Paulo, SP, Brazil. Email: jose.otavio@fm.usp.br; <https://orcid.org/0000-0002-3919-1743>.

Correspondence: Cristina Camargo. Av. Brigadeiro Luis Antonio, 4161. CEP: 01401-001 São Paulo, SP, Brazil. Email: consultoriocrisrina@gmail.com

Table 1. Examples of subjective and objective assessment in surgical training⁵

Subjective	Objective
Operative log book	Checklists
Examination	Global rating scales
Complication data (morbidity and mortality)	Error score rating
Time expenditure by each task	Dexterity analysis

Dexterity analysis - this analysis can be used for any manual skill evaluation. In surgery a device detects all the surgeon movements and up load in a computer system to measure the range motion. Low range motion more movement accuracy.

Reliability - The skill can be tested by two or more different student. The result must be similar (inter-rater reliability).

Because all safety and quality political changes for patient assistance, and the decrease of surgical rotation period due to knowledge overload, an effective and objective assessment is necessary to achieve competence

The aim of this article is to review surgical training assessment.

It is interesting to comment that these tools can be applied for education undergraduate students learning and for continued medical education (CME).

One of the most frequent assessment formats is OSCE- objective structures clinical examination. The Toronto group adapt this model for surgical skills evaluation: OSATS – objective structured assessment for technical skills^{6,7,8}.

Moreover other objective type of assessments, such as dexterity, global rating scale can be used in the association of OSATS to improve surgical training evaluation⁸.

ASSESSMENT METHODS

OSATS

The objective structured assessment of technical skills is based on stations where the students perform a specific surgical ability in a fixed period of time. Depend on the aim of the evaluation this assessment can be divided in 6-8 different stations. Each station mimics a technical skill or part of a particular ability, in the case of complex abilities.

The advantages of this type of assessment are task

flexibility, several models can be used: animal, simulators, cadaver, objective evaluation. The disadvantages are cost, schedule complexity, human resources (number of monitors and education and time expenditure) to apply this evaluation^{4,8}.

Dexterity

A computer program can be used to evaluate human movement assessment and simulation.

Human movement assessment is a kinesiology tool to study human movements. This evaluation allows studying purposeful hand direction, depth perception, finger coordination, movement speed, dexterity and movement precision⁸.

Some programs are available in the market (ADEP⁹, ICSAD¹⁰). Despite their technical characteristics all the software aim is to measure surgeon accuracy.

Another interesting simulation model is virtual reality simulator. The 3-D software provides a “real world” simulation, but the available models are considered as a low-fidelity simulator. Some new softwares are in testing phase and included haptic properties (force and touch skills) to promote a more realistic model of assessment⁴.

Global Rating Scale

This scale is based on seven questions related to a particular surgical ability evaluated by a Likert scale (5-point scale)¹¹.

For an illustrative example see knots and suture assessment section and Table 2.

Another methods to evaluate surgical skills

We will describe some assessment related to surgical skills.

Ultra-violet / fluorescein analysis in surgical scrubbing and hand hygiene

All the students will scrub their hand with a mixture of surgical soap and fluorescein (BODE® Visirub dissolved in Sterillium, Schülke Optics, B. Braun Fluo-Rub®, and Ecolab Magic Blue®)¹².

The next step the student will expose their hands in a ultra-violet lamp and a digital photographic will be done. This photography will be digitalized and by an Image J Pro®¹³ software the no fluorescent parts of the hands will be measure. If we found an area greater than 5% of the total hand area a new training will be necessary¹⁴.

Knots and suture

Global rating scale

The most used tool to evaluate knots and suture is the Global rating score. Although the authors claim objectivity some aspects continue to be considered subjective¹¹.

Tabela 2. Global rating scale¹¹

Rating	1	2	3	4	5
Respect for tissue	Frequently used unnecessary force on tissue or causal damage by inappropriate use of instruments		Careful handling of tissue but occasionally caused inadvertent damage		Consistently handled tissues appropriately with minimal damage
Time and motion	Many unnecessary moves		Efficient time/motion but some unnecessary moves		Clear economy of movement and maximum efficiency
Instrumental handling	Repeatedly makes tentative or awkward moves with instruments by inappropriate use of instruments		Competent use of instruments but occasionally appeared stiff or awkward		Fluid moves with instruments and no awkwardness
Knowledge of instruments	Suture and knots with inadequate tension		Competent suture, appropriate tension and, adequate knots		Excellent suture, with adequate tension and knots
Flow of operation	Frequently stopped operating and seemed unsure of next move		Demonstrated some forward planning with reasonable progression of procedure		Obviously planned course of operation with effortless flow from one move to the next
Use of assistants	Consistently placed assistants poorly or failed to use assistants		Appropriate use of assistants most of the time		Strategically used assistants to the best advantages at all time
Knowledge of specific procedure	Deficient knowledge. Needed specific instruction at most steps		Knew all important steps of operation		Demonstrated familiarity with all aspects of operation
Overall results	Fail	Fail	Pass	Pass	Pass

Pender et al.¹⁵, showed an interesting tool to evaluate knots and suture skills. The paper studied 12 different knots and suture¹⁵ (Table 3).

Table 3. Knots and suture training assessment

Task	Category	Competence	Aim
Two handed, under tension, suture's knot	Accuracy, gap, slippage, breakage (suture)	Time ≤ 26 sec, no errors	Two consecutives repetitions (upper limit=80)
Two handed, under tension, slip knot	Accuracy, gap, slippage, breakage (suture)	Time ≤ 30 sec, no errors	Two consecutives repetitions (upper limit=80)
One handed, under tension, slip knot	Accuracy, gap, slippage, breakage (suture)	Time ≤ 30 sec, no errors	Two consecutives repetitions (upper limit=80)
Suture, simple, interrupted	Accuracy, gap, slippage, breakage (suture)	Time ≤ 36 sec, no errors	Two consecutives repetitions (upper limit=80)
Suture, Suture, interrupted, horizontal mattress	Accuracy, gap, slippage, breakage (suture)	Time ≤ 60 sec, no errors	Two consecutives repetitions (upper limit=80)
Suture, interrupted, vertical mattress	Accuracy, gap, slippage, breakage (suture)	Time ≤ 36 sec, no errors	Two consecutives repetitions (upper limit=80)

Another method to assess knots and suture was proposed by Goova et al.¹⁷ The authors assessed six different types of knots and suture. The evaluation was based on the following criteria:

- ✓ Proficiency training goal (number of consecutive procedure repetitions);

- ✓ error rate (accuracy, gap in the knot's sequence, slippage and breakage of the knots and suture);
- ✓ proficiency score (time to perform the task with no errors)^{16,17} (Table 4).

Errors - see Table 5.

Table 4. Knots and suture training assessment

Task	Cutoff time (sec)	Proficiency score
Palm needle driver	60	53 (7 sec), no errors
Two-handed knot, without tension	60	50 (10 sec), no errors
One handed knot, without tension	60	50 (13 sec), no errors
Two-handed knot, with tension, surgen's knot	60	47 (15 sec), no errors
Two-handed knot, with tension, slip knot	60	45 (15 sec), no errors)
One-handed knot, with tension, slip knot	60	45 (15 sec), no errors
Simple interrupted suture	120	102 (18 sec), no errors
Suture, Suture, interrupted, horizontal mattress	120	89 (31 sec), no errors
Suture, interrupted, vertical mattress	120	89 (31 sec), no errors
Simple running suture	600	435 (165 sec), no errors
Subcuticular running suture	600	396 (204 sec), no errors
Subcuticular interrupted suture	120	87 (33 sec), no errors

Table 5. Errors description related to knots and suture assessment

Task	Error
Palm needle driver	No closure of needle drives (3 clicks), accuracy, instrument drop, non-palming
One and two-handed knots and suture	No closure of needle drives (3 clicks), accuracy, instrument drop, non-palming, knot gap (bigger than 3 mm)

Each one of the table errors descriptions correspond to a minus one point in the equation.

However some errors demand a higher punctuation:

- ✓ Slippage bigger than 3 mm, (- 10 points);
- ✓ Total slippage or knot break (- 20 points);
- ✓ Protocol violation – fatal flaw.

According to the previous information, the assessment score can be calculated by the following equation^{16,17}:

- ✓ Cutoff time – completion time – 10X (sum of the errors);
- ✓ The cutoff score are showed in the Table 4.

RESEARCH IN EDUCATION

Many other surgical skills do not have an objective assessment. This type of assessment must be based on Myller's pyramid for declarative and procedural knowledge

acquisition. Basically, the student has to know how this work and show it.

Moreover, there is a correlation between student self-confidence and competence^{1,2,18}. According to Clanton et al.¹⁸ the correlation was 0.88 for knot training in third year medical students.

The confidence evaluation questionnaire was based on the Likert scale and some suggestions for the confidence evaluation are disposable in the medical literature¹⁸.

As an example we showed the following questionnaire extract from Clanton et al.¹⁸, paper.

Confidence questionnaire of surgical skills

Please rate your confidence to complete the following tasks on a scale of 1 to 6.

I feel confident that I can:

1. Tie 10 complete 2-handed square knots using proper surgical technique;

2. Tie 10 complete 1-handed square knots using proper surgical technique;
3. Tie a surgeon's knot using proper surgical technique;
4. Complete a 2-handed with appropriate tension;
5. Complete a 2-handed square knot without the knot breaking;
6. Complete a 2-handed square knot without the knot unraveling;
7. Complete a 2-handed square knot with appropriate speed and efficiency;
8. Tie a series of 4 instrument tied square knots using proper surgical technique;
9. Perform an instrument tied square knot with appropriate tension;
10. Perform an instrument tied square knot while everting the skin edges;
11. Perform an instrument tied square knot without the knot breaking;
12. Perform an instrument tied square knot without the knot unraveling;
13. Perform an instrument tied square knot with appropriate speed and efficiency;
14. Demonstrate proper needle loading onto needle driver;

15. Demonstrate a simple interrupted stitch using proper surgical technique;
16. Demonstrate a simple running stitch using proper surgical technique;
17. Demonstrate suturing technique without excessively damaging the tissue;
18. Suture a wound with a good cosmetic result.

Covariates in learning assessment

Another relevant issue to be considered in medical education is some factors that can interfere in the learning process.

Some of these the variables have been studied: gender, handedness, scholar background, previous experience, psychological traits.

Although, the covariate factors influence in the learning process, the increment of all those covariates can work as confounders factors¹⁹⁻²⁰.

For this reason, and according to literature data, the most relevant factors are: gender, handedness and previous experience related to the studied skill^{7,19,20}.

CONCLUSION

The authors described some objective surgical skill assessment based on OSAST, dexterity and global rating scale. These tools were important to analyze the learning effectiveness regarding long-term memory and retrieval memory.

4. REFERENCES

1. Agha RA, Papanikitas A, Baum M, Benjamin IS. The teaching of surgery in the undergraduate curriculum. Part II e Importance and recommendations for change. *Int J Surg.* 2005;3:151e-57. doi: 10.1016/j.ijssu.2005.03.016.
2. Stefanidis D. Optimal acquisition and assessment of proficiency on simulators in surgery. *Surg Clin North Am.* 2010;90(3):475-89. doi: 10.1016/j.suc.2010.02.010.
3. Milburn JA, Khera G, Hornby ST, Malone PS, Fitzgerald JE. Introduction, availability and role of simulation in surgical education and training: review of current evidence and recommendations from the Association of Surgeons in Training. *Int J Surg.* 2012;10(8):393-8. doi: 10.1016/j.ijssu.2012.05.005.
4. Moorthy K, Munz Y, Sarker SK, Darzi A. Objective assessment of technical skills in surgery. *Br J Med.* 2003;327(1):1032-7. doi: 10.1136/bmj.327.7422.1032.
5. Davies J, Khatib M, Bello F. Open surgical simulation - a review. *J Surg Educ.* 2013;70(5):618-27. doi: 10.1016/j.jsurg.2013.04.007.
6. Acton RD. The evolving role of simulation in teaching surgery in undergraduate medical education. *Surg Clin N Am.* 2015;95:739-50.
7. Tavakol M, Mohagheghi MA, Dennick R. Assessing the skills of surgical residents using simulation. *J Surg Educ.* 2008;65(2):77-83. doi: 10.1016/j.jsurg.2007.11.003.
8. Martin JA, Regehr G, Reznick R, MacRae H, Murnaghan J, Hutchison C, et al. Objective structured assessment of technical skill (OSATS) for surgical residents. *Br J Surg.* 1997;84:273-8. doi: 10.1046/j.1365-2168.1997.02502.x.
9. Francis NK, Hanna GB, Cuschieri A. The performance of master surgeons on the advanced Dundee endoscopic psychomotor tester: contrast validity study. *Arch Surg.* 2002;137:841-4. doi: 10.1001/archsurg.137.7.841.
10. Datta V, Mackay S, Mandalia M, Darzi A. The use of electromagnetic motion tracking analysis to objectively measure open surgical skill in the laboratory-based model. *J Am Coll Surg.* 2001;193:479-85. doi: https://doi.org/10.1016/S1072-7515(01)01041-9.
11. Reznick R, Regehr G, MacRae H, Martin J, McCulloch W. Testing technical skill via an innovative "bench station" examination. *Am J Surg.* 1997;173(3):226-30. doi: https://doi.org/10.1016/S0002-9610(97)89597-9.
12. Fichtner A, Haupt E, Karwath T, Wullenk K, Pohlmann C, Jatzwauk L. A single standardized practical training

- for surgical scrubbing according to EN1500: Effect Quantification, value of the standardized method and comparison with clinical reference groups. *GMS Zeitschrift für Medizinische Ausbildung*. 2013;30(2):3-21. doi: 10.3205/zma000867.
13. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods*. 2012;9(7):671-5. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3555454/>.
 14. World Health Organization. Patient Safety. A World Alliance for Safer Health Care. WHO guidelines on hand hygiene in health care. First global patient safety challenge clean care is safer care [cited 2017 March 20]. Available from: http://apps.who.int/iris/bitstream/10665/44102/1/9789241597906_eng.pdf.
 15. Pender C, Kiselov V, Yu Q, Mooney J, Greiffenstein P, Paige JT. All for knots: evaluating the effectiveness of a proficiency-driven, simulation-based knot tying and suturing curriculum for medical students during their third-year surgery clerkship. *Am J Surg*. 2017;213(2): 362-70. doi: 10.1016/j.amjsurg.2016.06.028.
 16. Scott DJ, Goova MT, Tesfay ST. A cost-effective proficiency-based knot-tying and suturing curriculum for residency program. *J Surg Res*. 2007;141:7-15. doi: 10.1016/j.jss.2007.02.043
 17. Goova MT, Hollet RN, Tesfay ST, Gala RB, Puzifferri N, Kehdy FJ, Scott DJ. Implementation, construct validity, and benefit of a proficiency-based knot-tying and suturing curriculum. *J Surg Educ*. 2008; 65(4): 309-15. doi: 10.1016/j.jsurg.2008.04.004.
 18. Clanton J, Gardner A, Cheung M, Mellert L, Evancho-Chapman M, George RL. The relationship between confidence and competence in the development of surgical skills. *J Surg Educ*. 2014; 71(3): 404-12. doi: 10.1016/j.jsurg.2013.08.009.
 19. Ferguson E, James D, Madeley L. Factors associated with success in medical school: systematic review of the literature. *BMJ : Brit Med J*. 2002;324(7343):952-957.
 20. Gilligan J, Watts C, Welsh F, Treasure T. Square pegs in round holes: has psychometric a place in choosing a surgical career? A preliminary report of work in progress. *Ann R Coll Surg Engl*. 1999;81:73e9. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2503198/>.

Recebido em: 22.11.17

Aceito em: 11.01.18